

858

BEFORE THE HON'BLE NATIONAL GREEN TRIBUNAL

PRINCIPAL BENCH AT NEW DELHI

OBJECTION BY APPLICANT

In

ORIGINAL APPLICATION NO. 511 OF 2023

IN THE MATTER OF

Priyank Bharati

-----APPLICANT IN PERSON

Versus

State of Uttar Pradesh through its Chief Secretary and others


-----RESPONDENTS

INDEX

S.No	Particulars	Page No
1.	Objection by Applicant	01-10
2.	Affidavit	11
3.	Annexure 1	12-13
4.	Anex. 2	14
5.	Anex. 3	15-16
6.	Anex. 4	-46
7.	Anex. 5	47-48

Place :- Meerut

Date : 09.10.2025



Priyank Bharati

(Applicant In Person)

**BEFORE THE HON'BLE NATIONAL GREEN TRIBUNAL
PRINCIPAL BENCH AT NEW DELHI**

OBJECTION BY APPLICANT

In

ORIGINAL APPLICATION NO. 511 OF 2023

IN THE MATTER OF

Priyank Bharati

-----APPLICANT IN PERSON

Versus

State of Uttar Pradesh through its Chief Secretary and others

-----RESPONDENTS

**Affidavit-cum-Objection by Applicant to the Report Submitted by the
Chief Engineer (Ganga), Irrigation and Water Resources Department,
Meerut, Government of Uttar Pradesh dated 11 July 2025**

MOST RESPECTFULLY SHOWETH;

I, Priyank Bharati R/O Jagriti Vihar Meerut-250004, UP hereby solemnly affirm and declare as under:

1. That the River (Budhi Ganga), which carries within its course the cherished legacy and historical reminiscence of the Mahabharata era, represents an invaluable element of India's natural and cultural heritage - a heritage which, regrettably, the institutions of independent India have failed to preserve and protect in accordance with their constitutional and statutory obligations.
2. That in compliance with the directions issued by the Hon'ble Tribunal, the Flood Plain of River Budhi Ganga has been demarcated by the Irrigation Department (as most of the data given by Irrigation

department itself to NIH) through the National Institute of Hydrology (NIH), Roorkee.

3. That the Applicant has repeatedly communicated through official emails requesting the KML files pertaining to the said demarcation; however, several essential files, including the alignment of the river along with its cross-sections, have not been furnished to the Applicant till date. The copy of the email is enclosed herewith and marked as

ANNEXURE 1

4. That, according to the report page no 671, 672, 673, 678 and 680 the **identification of the origin point of River Budhi Ganga** has been **erroneously** determined, thereby rendering the entire delineation of its floodplain scientifically untenable and legally unsustainable. The Report stands vitiated **ab initio**, as it is founded upon an incorrect and defective point of commencement.

5. That, according to the report contradictory statements are made regarding the number of river reaches (Pages 660 and 669)

On **page no. 660**, it is stated that:

*"During the field visit by NIH team along with Irrigation and Water Resources Department, Govt. of U.P., it was also found that at present there are **four river** reaches."*

Whereas, on **page no. 669**, it is stated that:

*"The Budhi Ganga River, a tributary, is having very less width and water flows mainly during the rainy season and joins the Ganga **River at 3 to 4 locations**"*

Both **statements are contradictory** in nature. The report is thus ambiguous even regarding the fundamental geomorphic characterization of the river. The Irrigation Department to ascertain the actual number and continuity of river reaches clearly reflects verification and lack of hydro-geomorphic clarity. It proves that till date,

the concerned departments do not possess the actual delineation or verified stretches of River Budhi Ganga.

6. That on **page 659, point no. 7** of the report, it is stated that the three issues earlier faced by NIH, as mentioned in the previous affidavit dated 12.03.2025 (at page 627), have been rectified, and the details of such rectifications are provided on **page 660** of the report. There are some objections on issues rectified which are as follows:

Issues (pg 660)	How it is addressed in the report (pg 660)	Objection by applicant
The Digital Elevation Model (DEM) from Survey of India (SOI) doesn't properly represent the river course. The elevation in the river course are higher than surrounding area . This might have happened as vegetations in the river channel not properly accounted for while generating the DEM.	The river cross-sections at 200 m interval was provided to NIH by the Irrigation and Water Resources Department, Govt. of UP. These surveyed river cross-section data is used in the modelling study and outside the river cross-section area the 0.5 m DEM data is used	That it is pertinent to note that on page 632, point 2 of the report clearly mentions that the said cross-sections already existed prior to the identification of the problem. Therefore, it remains unclear how the same pre-existing cross-sectional data, as referred to on page 660, could have been utilized to resolve the issue arising from the inaccurate representation of the river course in the DEM. The DEM does not accurately represent the river course, and

		reliance on cross-sections at 200 m intervals is insufficient to ensure scientific accuracy. Therefore, a re-survey using high-resolution DEM, detailed bathymetry, and closely spaced cross-sections is essential for a precise and legally sustainable floodplain delineation of River Budhi Ganga.
The surveyed river cross-section's locations at few sections is not on the river course alignment of DEM	The Irrigation and Water Resources Department, Govt. of UP. provided the KML file of alignment of river along with cross-sections. These alignments used as river centreline in this report	The mere use of departmental KML data, without cross-checking renders the delineation scientifically questionable and legally unsustainable. As this KML file of alignment of river not provided to Applicant.

7. That, the statement regarding the operation of canal is factually incorrect (Page 677)

On page no. 677, NIH Roorkee mentioned that:

"As a conservative approach, the Runoff Coefficient of 0.9 is used as during rainy season and the canal is in operation..."

This statement is not accepted by the applicant. The Madhya Ganga Canal system remains **non-operational during the rainy/monsoon season** or in heavy floods, as confirmed by field verification and photographic evidence by applicant. The assumption that the canal is operational during monsoon has resulted in erroneous discharge estimation and hydrological misrepresentation in the floodplain modelling. The geotagged photos are annexed herewith and marked as **ANNEXURE 2**

8. **That in the Hastinapur region**, there exists a natural stream of river which originates locally from Kaurvan region (as also visible in toposheet page no 682) and subsequently confluences with the main channel of River Budhi Ganga at Dudhli Khadar/Shimla; however, the said stream has been completely omitted from consideration in the Flood Plain Delineation Report (pg 671, 672, 673, 678, 682 and 685), thereby rendering the report incomplete and scientifically inaccurate. This constitutes a serious **omission of factual and environmental assessment**. The Google Earth image is annexed herewith and marked as **ANNEXURE 3**.

9. That on **page no 676 point no 5.1** under the head of **Flood Frequency Analysis**

Statement in Report	Objection
Historical discharge series of for flood analysis is not available. Hence the discharge corresponding	That to represent the catchment and sub-catchment areas very small , the gauge sites have been selected several kilometres away

to 100 year return period is estimated from the rainfall data. The estimated value of catchment characteristics are given in table 5.1. It is noted that the catchment and sub catchment area are very small. The nearest three rain gauge station namely, Bijnor, Mawana and Hasanpur are selected based on the sub catchment area.	from the actual river stream. However, the report does not contain any map or detailed information delineating the catchment and sub-catchment areas of River Budhi Ganga. It is pertinent to mention that the river flows through Districts Muzaffarnagar, Meerut, and Hapur, yet the rain gauge stations considered in the report are not situated within the actual catchment or sub-catchment area of River Budhi Ganga, thereby rendering the hydrological analysis and results unreliable.
--	--

10. That in **Assumption and Limitations, on page 686** of the report,

According to report page no 686	Objection by Applicant
<i>Limitation of river bathymetry below the water spread area in the SOI DEM and accounted using available cross section only</i>	The Flood Plain Delineation Report suffers from significant methodological limitations, as the river bathymetry below the water spread area in the SOI DEM has not been properly represented and has been accounted for merely through limited cross-sectional data.
<i>Evaporation, infiltration/groundwater interaction and diversion losses neglected.</i>	The omission of critical hydrological parameters such as groundwater interaction

undermines the accuracy of the hydrodynamic modelling.

Floodplain delineation is generally incomplete without considering surface-groundwater interactions. Groundwater and surface water interaction plays a significant role in floodplain hydrology because floodplains act as the interface zones where exchanges occur between streams and groundwater systems. Refer ANNEXURE 4

Then the flood plain delineation **cannot be considered scientifically accurate or complete**. Furthermore, Hence, the delineation prepared on such incomplete data and assumptions cannot be treated as scientifically sound or legally sustainable.

11. That **the energy equation** used for calculating the water surface profile, as mentioned on **page 674** of the report, differs from the formulation provided in the *HEC-RAS River Analysis System, Hydraulic Reference Manual*, Version 6.5, March 2024. If any modification in the formula has been made, it is necessary to specify the reasons for such alteration and clarify why the variables or alphabets have been rearranged. A copy of the relevant pages of the HEC-RAS Hydraulic Reference Manual is attached herewith and marked as **ANNEXURE 5**
12. That, **until proper and verified floodplain demarcation is completed, all activities must be stopped on River Budhi Ganga and its floodplain**. It is respectfully submitted that **no construction work, land sale, or allotment activities** should be permitted within the proposed or disputed floodplain stretch of Budhi Ganga in **District**

Muzaffarnagar, District Meerut and District Hapur till a scientifically verified demarcation is approved by the Hon'ble Tribunal after public disclosure of KML, DEM, and ground-truth data. In certain areas, including Deval in District Muzaffarnagar and Hastinapur in District Meerut and several others, construction activities are still being carried out within the floodplain of River Budhi Ganga, in contravention of the environmental safeguards and regulatory provisions governing floodplain zones.

- 13. That, the model parameters and calibration details are missing (Pages 674–678).** No details are provided about **boundary conditions, mesh size, sensitivity analysis, or model validation** of the HEC-RAS 6.5 simulation. Manning's coefficients are arbitrarily assigned without field calibration. Such omissions invalidate the technical credibility of the modelling process.

- 14. That on page 687 it is mentioned in the report that :**

Statement	Objection to the statements
However, at present there is no surface flow from the Ganga River to Budhi Ganga River in normal Condition.	Under normal conditions, the Budhi Ganga River maintains surface water flow throughout the year, except during the very dry months of May and June.
It is recommended that while conducting the Flood Plain Zoning of the River Ganga for 100-year return period flood, these tributaries may be	If it is recommended that tributaries be jointly modelled with the Ganga for 100-year flood studies then in this condition this flood plain is accepted or not ?

modelled/studied together for combined effects.	Clarification is required on how such delineation has been approved in contradiction to the stated recommendation.
---	--

Prayer

In light of the above deficiencies, contradictions, and data concealment, the Applicant most respectfully prays that the Hon'ble Tribunal may kindly:

1. Direct the **Irrigation and Water Resources Department** and **NIH, Roorkee** to submit complete datasets including DEM, KML, KMZ, and cross-section files for verification.
2. Direct **immediate suspension of all construction, sale, or allotment activities** within the suspected floodplain zone till final demarcation is approved.
3. Direct **Irrigation and Water Resources Department** and **NIH** to carry out a **revised and ground-truth floodplain demarcation** integrating hydrology, ecology, ground water and surface water interaction parameters as per Hon'ble NGT and Ganga Rejuvenation guidelines.
4. Pass any such and further orders as this Hon'ble Tribunal may deem fit and proper in the facts and circumstances of the case.

For all the reasons stated above the Hon'ble Tribunal may kindly be pleased to take cognizance of all these facts and to pass appropriate orders to meets the ends of Justice and equity.

AND FOR THIS ACT OF KINDNESS THE APPLICANTS, AS IN DUTY BOUND, SHALL EVER PRAY.


Verification

Verified on this 9th day of October 2025 that the contents of the present Application are true and correct to my knowledge and belief and nothing material is concealed therefrom.



Priyank Bharati

Applicant in Person



Date : 09th Oct 2025

Place : Meerut



ATTESTED
NOTARY

9/10/25

**BEFORE THE HON'BLE NATIONAL GREEN TRIBUNAL
PRINCIPAL BENCH AT NEW DELHI**

ORIGINAL APPLICATION NO. 511 OF 2023

IN THE MATTER OF

Priyank Bharati

-----APPLICANT IN PERSON

Versus

State of Uttar Pradesh through its Chief Secretary and others

-----RESPONDENTS

AFFIDAVIT

I, Priyank Bharati, S/o, Shri. Brahampal Singh aged about 36 years, R/o Jagriti Vihar, District Meerut, Uttar Pradesh do hereby solemnly affirm and declare as under:

That I am the Applicant/Applicant In Person in above mentioned application and I am fully conversant with the facts and circumstances of the case and therefore competent to swear this affidavit.

That the statements made in above paragraphs of this affidavit is true to my knowledge and belief.

That the contents of the Application are true and correct and nothing material has been concealed therefrom.

Sharali

Applicant In Person



VERIFICATION

Verified on this 09th day of October 2025 that the contents of the present Application are true and correct to my knowledge and belief and nothing material is concealed therefrom,

Sharali

Applicant In Person/Deponent

ATTESTED
9/10/25
NOTARY





Priyank Bharati <naturalsciencetrustmrt@gmail.com>

Request for Copy of Flood Plain Delineation Report of River Budhi Ganga in Compliance with Hon'ble Tribunal's Order Dated 14.07.2025 in the matter of OA 511 of 2023.

3 messages

Priyank Bharati <naturalsciencetrustmrt@gmail.com>
To: bhanwar jadon <bhanwar09jadon@gmail.com>

Mon, Jul 14, 2025 at 1:49 PM

To,
14.07.2025**Mr. Bhanwar Pal Singh Jadon,****Standing Counsel for the State of UP****Subject: Request for Copy of Flood Plain Delineation Report of River Budhi Ganga in Compliance with Hon'ble Tribunal's Order Dated 14.07.2025 in the matter of OA 511 of 2023.**

Respected Sir,

I respectfully submit that during the course of hearing dated **14th July 2025**, the **Hon'ble National Green Tribunal** was pleased to direct that the **flood plain delineation report of River Budhi Ganga** be made available to Applicant.

In compliance with the said order, I hereby request that the **entire copy of the floodplain delineation report of River Budhi Ganga**, along with all accompanying maps, annexures and explanatory notes, kindly be provided to me at the earliest.

I shall be grateful for your prompt cooperation in this regard.

With regards,
Priyank Bharati

Applicant

Priyank Bharati <naturalsciencetrustmrt@gmail.com>
To: bhanwar jadon <bhanwar09jadon@gmail.com>, "Consultant Judicial-NGT(P.B.)" <judicial-ngt@gov.in>

Wed, Jul 16, 2025 at 11:07 AM

Respected Sir,

I acknowledge receipt of your email dated 14th July 2025. **However, the KML file of the flood plain appears to be missing from the attachments.**

I kindly request you to provide the KML file as referred to in Point No. 10 and Annexure A1 of your communication.

Thank you for your attention to this matter.

Warm regards,
Priyank

[Quoted text hidden]

Priyank Bharati <naturalsciencetrustmrt@gmail.com>
To: bhanwar jadon <bhanwar09jadon@gmail.com>, cegangaidupme-up@nic.in

Fri, Sep 5, 2025 at 9:04 AM

Respected Sir,
Kindly provide the KML file at the earliest so that I may submit my objections in OA 511/2023 within the stipulated time.

Warm Regards,

Priyank Bharati
(Applicant In Person)
Mobile : 09411823914

871


13

cc:

1. **Shri Gyan Prakash Srivastava Ji , Chief Engineer-II (Ganga), Meerut**

[Quoted text hidden]





Hastinapur

Write a description for your map.

16
Legend


874

NULL

Google Earth

Image © 2025 Airbus


2 km





Contents lists available at ScienceDirect

Journal of Hydrology

journal homepage: www.elsevier.com/locate/jhydrol

Research papers

Groundwater-Surface water interactions research: Past trends and future directions

Dylan J. Irvine^{a,b,*}, Kamini Singha^c, Barret L. Kurylyk^d, Martin A. Briggs^e, Yakub Sebastian^a, Douglas R. Tait^f, Ashley M. Helton^g^a Research Institute for the Environment and Livelihoods and Faculty of Science and Technology, Charles Darwin University, Casuarina, Northern Territory, Australia^b National Centre for Groundwater Research and Training, Bedford Park, South Australia, Australia^c Hydrologic Science and Engineering Program, Colorado School of Mines, Golden, CO, USA^d Department of Civil and Resource Engineering and Centre for Water Resources Studies, Dalhousie University, Halifax, Canada^e U.S. Geological Survey, Observing Systems Division, Hydrologic Remote Sensing Branch, Storrs, CT, USA^f Faculty of Science and Engineering, Southern Cross University, Lismore, New South Wales, Australia^g Department of Natural Resources and the Environment, and the Center for Environmental Sciences and Engineering, Storrs, University of Connecticut, Connecticut, USA

ARTICLE INFO

Keywords:

Rivers
Streams
Lakes
Hyporheic zone
Baseflow
Groundwater-dependent ecosystems
Numerical modelling
Climate change
Submarine groundwater discharge
UAVs

ABSTRACT

Interactions between groundwater and surface water sustain groundwater-dependent ecosystems and regulate river temperature and biogeochemical cycles, amongst many other processes. These interactions occur in freshwater environments including rivers, springs, lakes, and wetlands, and in coastal environments via tidal pumping, submarine groundwater discharge, and seawater intrusion. Here, we explore groundwater-surface water interactions research using bibliometric analyses of titles, abstracts, and keywords from 20,275 journal papers published between 1970 and 2023 extracted from Scopus. Analyses show that research into groundwater-surface water interactions is highly multi-disciplinary, with growing contributions from the social and biological sciences. The number of groundwater-surface water interactions papers is rapidly increasing with over 1200 papers published per year since 2020. Drawing on our data-driven approach and expert knowledge, we synthesise current research trends and identify critical future research directions. Despite the thousands of papers on groundwater-surface water interactions, important processes are still difficult to quantify or predict at meaningful spatial scales to inform water-resources management. We see benefits in future groundwater-surface water interactions research focusing on: (1) using new technologies including internet-of-things-based sensors, uncrewed vehicles, and remote-sensing approaches for data collection to inform groundwater-surface water interactions at large scales, (2) seeking approaches to upscale site-specific findings to better inform management, and (3) continuing the movement towards multi-disciplinary investigations to better inform the understanding of groundwater-surface water interactions and processes that will enable better management outcomes.

1. Introduction

Groundwater and surface water are widely recognised as interconnected hydrologic systems (e.g., Boano et al., 2014; Brunke and Gonser, 1997; Winter et al., 1998). Modern research into groundwater-surface water (GW-SW) interactions dates to the 1930 s, with multi-disciplinary GW-SW interactions research widely published across diverse fields of study including hydrogeology, hydrology, limnology, amongst many others. This multi-disciplinary focus is necessary given the broad range of processes that influence and are influenced by GW-

SW interactions. For example, GW-SW exchange is a hydrological process that plays important biogeochemical and ecological roles (Ward, 2016), including regulating surface water temperature regimes (Benz et al., 2024; Hare et al., 2021), controlling processes including denitrification and the redistribution of oxygen in the hyporheic zone (Boano et al., 2014; DelVecchia et al., 2022), and sustaining riparian zone vegetation (Harvey and Gooseff, 2015). Understanding GW-SW interactions is also important in studies of baseflow (Arnold and Allen, 1999; Ladson et al., 2013; McCallum et al., 2010), bank storage (Cartwright and Irvine, 2020; Doble et al., 2012), flooding (e.g., Blöschl

* Corresponding author at: Research Institute for the Environment and Livelihoods and Faculty of Science and Technology, Charles Darwin University, Casuarina, Northern Territory, Australia.

E-mail address: dylan.irvine@cdu.edu.au (D.J. Irvine).

<https://doi.org/10.1016/j.jhydrol.2024.132061>

et al., 2007; Miguez-Macho and Fan, 2012), and stream depletion (e.g., Huang et al., 2018; Hunt, 2014). GW-SW exchange is driven by factors including hydraulic gradients, geologic heterogeneity (e.g., Tonina and Buffington, 2009; Wohl, 2021), and channel morphology (Cardenas and Zlotnik, 2003a; Stonedahl et al., 2010). The wide-ranging drivers and effects of GW-SW interactions intersect with many broader fields, from geology, hydrology, and fluvial geomorphology to biogeochemistry and ecohydrology.

Effective water-resources management is one of humanity's greatest priorities (e.g., Jakeman et al., 2016) given the expanding demand for water by the increasing global population and a rapidly changing climate punctuated by extreme events. Despite its importance, conjunctive management of groundwater and surface water resources is challenging. Challenges include the complexity of GW-SW systems, measuring and summarising those system dynamics at management-relevant scales and the often-fragmented decision-making process caused by the involvement of multiple agencies (Howe, 2002). Natural and human-driven changes to GW-SW systems cause a myriad of important effects, resulting in aquifers and groundwater-connected systems that function as 'socioecological systems' (Huggins et al., 2023). Therefore, a detailed understanding of the natural processes that GW-SW interactions support, and the effects of future climate change and further extraction of water resources could help inform the management of groundwater and surface water resources.

With many thousands of papers on the topic, GW-SW interactions are now a well-established research field. Here we take the opportunity to provide an overview of research trends and consider potential future research trajectories. We use the Scopus database (Elsevier, 2024) to explore GW-SW interactions research and assess publishing trends and trends in the focus of GW-SW interactions research. Open questions include how the field of GW-SW interactions has evolved since research began in earnest in the 1970 s, where the field is headed, and what are the future research needs and opportunities. To address these questions, we present a bibliometric analysis of GW-SW interactions studies published between 1970 and 2023. After providing an overview of GW-SW

interactions processes (Section 2), we detail our literature review methodology (Section 3), we then focus on publishing trends through time (Sections 4.1 to 4.3), the journals and countries publishing in the field (Sections 4.4 and 4.5), and the relationships between related research areas (Section 4.6). We then investigate the current state of the field, including an attempt to identify recent work that is projected to have a major future impact (Section 5.1), and a summary of recent research trends (Section 5.2). Finally, we discuss future GW-SW interactions-research needs and opportunities (Section 5.3).

2. An overview of GW-SW interactions

2.1. GW-SW interactions, processes, and zones

Subsurface waters exchange with rivers, lakes, streams, springs, wetlands, and the ocean (Fig. 1), where dynamic hydraulic head gradients between surface waters and the adjacent groundwater control the directionality of exchange (Fig. 1b). A 'gaining' system occurs where the surface water body increases in volume through seepage, fracture flow, and macropore discharge. Conversely, a 'losing' system occurs where surface water is routed to either the saturated or unsaturated zone in the subsurface. The relationship between groundwater and surface water is dynamic in both time and space; for example, (1) rivers can gain and lose water along their length, possibly with nested gaining/losing reaches (Payn et al., 2009), (2) heavy rainfall can temporarily cause a gaining section of river to transition to losing (e.g., Fig. 1b, Doble et al., 2012), and (3) groundwater extraction can cause surface water bodies (or at least part of a river system) to switch from gaining to losing (e.g., Hunt, 2014; Zipper et al., 2022), with various processes leading to diurnal variations in GW-SW exchange (Wu et al., 2021). The difference in hydraulic head between surface water and groundwater and flow directions is consistent across all GW-SW interactions settings, including in the coastal zone (e.g., Fig. 1a, 1e) where fluid density variations influence hydraulic heads and tides induce 'pumping' exchange (e.g., Ma and Zhang, 2020; Sekar et al., 2022), and in 'disconnected' systems




Fig. 1. Diagrams of groundwater-surface water (GW-SW) interactions throughout (a) a hydrological system including groundwater, a river and floodplain, lake, spring, and the ocean. SGD denotes submarine groundwater discharge. (b) Cross section showing the riparian zone, a floodplain wetland, the processes of bank storage, the direction of GW-SW exchanges at two river stages (blue denotes gaining conditions, red denotes losing conditions), and connections to the critical zone, i. e., from treetops to the water table (Sawyer et al., 2016b; Singha and Navarre-Sitchler, 2022). (c) Lacustrine-groundwater interactions, flow through conditions shown. (d) Hyporheic exchange (white arrows) caused by river geometry, (e) bedforms, and (f) Ocean-aquifer exchanges, including seawater intrusion, SGD, ocean water circulation, and tidal pumping.

where an unsaturated zone forms beneath surface water bodies (e.g., Brunner et al., 2009; Xie et al., 2014).

Bed materials have a strong control on GW-SW interactions, with the call for streambed science to be recognised as a scientific discipline (Constantz, 2016). GW-SW interactions are influenced by heterogeneities in aquifers and bed materials, often focusing GW-SW exchange to preferential zones of particularly strong hydrologic connectivity (e.g., Briggs and Hare, 2018; Covino, 2017). Measurements or estimates of streambed hydraulic properties can be obtained from a wide range of techniques, including hydraulic methods (e.g., Cardenas and Zlotnik, 2003a; Chen, 2000), the paired use of temperature time series and hydraulic gradients (Hatch et al., 2010), and equivalent porous media/modelling methods (Cardenas and Zlotnik, 2003a). The application of methods to estimate bed hydraulic properties is challenging in streams, creeks, and rivers, which are dynamic due to erosion and deposition events, relative to lentic settings. Brunner et al. (2017) and references therein discuss the challenges associated with the application of hydraulic methods due to the presence of thin biofilms at the water-sediment interface, and suggest that freeze-coring approaches (e.g., Ulrich et al., 2015) can help facilitate the quantification of streambed properties. Investigations into the role of streambed properties on stream-aquifer interactions include studies of chemistry and microbial composition (e.g., Knapp et al., 2017; Nelson et al., 2019) and their effect on river-aquifer disconnection (e.g., Schilling et al., 2017). Overviews of the role of the bed materials in regulating GW-SW interactions are discussed at length in reviews by Brunner et al. (2017) and Tripathi et al. (2021).

Groundwater discharge plays a vital role in sustaining surface water bodies (Hayashi and Rosenberry, 2002). For example, groundwater discharge generates river baseflow, maintaining flows through seasonal low flow and drought periods. Similarly, groundwater discharge sustains spring-fed wetlands and groundwater flow-through lakes (i.e., termed lacustrine-groundwater discharge, e.g. Meinikmann et al., 2013). In both instances, a reduction in groundwater head, through either reduction in rainfall or groundwater extraction, will reduce the rates of groundwater discharge. In the case of springs, should the reduced groundwater head fall below the spring geomorphic threshold (the land surface, or lip of a spring vent, e.g., Currell et al., 2017), spring flow will cease, which could have major ecological or cultural impacts (Currell et al., 2020; Davis et al., 2013). However, rainfall or anthropogenic modifications can influence GW-SW interactions, switching gaining conditions to losing conditions via temporary increases in river stage (e.g., Ziliotto et al., 2021). Major rainfall events that generate streamflow can also lead to large groundwater-recharge events. Recharge routing and timing depend strongly on hydrogeologic setting, including whether groundwater flow is through primary pore spaces, or secondary pore spaces in the case of karst or fractured rock aquifers. Additionally, hydropeaking from hydropower operations can influence bank-storage processes, temporarily driving surface water into the adjacent aquifer (Fig. 1b). The return to pre-event conditions is influenced by the flood wave height, hydraulic conductivity, riverbank slope, and regional hydraulic gradients (Cartwright and Irvine, 2020; Doble et al., 2012). Despite the importance of source groundwater attributes for evaluating the baseflow resiliency of a given surface water system, baseflow generation characteristics are difficult to predict with confidence, necessitating novel model calibration and validation techniques (Barclay et al., 2022, 2020).

GW-SW interactions play an important role in sustaining aquatic and terrestrial groundwater-dependent ecosystems. For example, in arid and semi-arid zones, groundwater-fed springs can provide habitat for endemic species (Currell et al., 2020; Davis et al., 2013), provide access to water for migratory birds (Kløve et al., 2011), and sustain tree stands (Doody et al., 2015). Comprehensive reviews of spring types and their functions include those by Alfaro and Wallace (1994), Springer and Stevens (2009), and Keegan-Treloar et al. (2022). Groundwater discharge can also temporally modulate stream and river temperatures

(Briggs et al., 2018; Burkholder et al., 2008), thereby increasing ecosystem niches and resilience during extreme high or low temperatures (Kurylyk et al., 2015). GW-SW exchange can also provide an important source of nutrients for aquatic ecosystems, particularly in nutrient-poor oligotrophic systems where groundwater provides the nutrients that drive patterns of primary productivity, biofilm growth that supports aquatic food webs, and even biodiversity (Naranjo et al., 2019; Valett et al., 1994).

The hyporheic zone (Fig. 1c, d) is defined as the subsurface region underlying/adjacent to lotic waters where surface water enters the sediments, mixes with groundwater, and a portion of the original surface water discharges back to the channel, often with altered chemistry, temperature, and gas content (e.g., Boano et al., 2014). The hyporheic zone has received increased research attention from disciplines including freshwater ecology, aquatic biogeochemistry, hydrogeology, geomorphology, terrestrial ecology and hydrology (Boulton et al., 2010; Harvey and Gooseff, 2015). Although no strict hydrologic definition exists for hyporheic exchange, the spatiotemporal window of relevant hyporheic flow paths and fraction of returned surface water is best judged relative to hyporheic-influenced processes of interest (Harvey et al., 2013). The hyporheic zone spans several spatial extents (Poole et al., 2008), ranging from permeable alluvial floodplains (Helton et al., 2014) to meander bends (Boano et al., 2006; Malenda et al., 2019), bedforms (Cardenas et al., 2008; Fox et al., 2014; Gomez-Velez et al., 2015) and shallow, coarse sediments beneath rivers and streams (Boano et al., 2014). The spatial extent of hyporheic zones often varies over time because of connections that influence the co-evolution of hydrodynamic, geomorphological, and hydrological processes. The hyporheic zone has been extensively reviewed, including broad reviews of hyporheic zone processes (e.g., Ward, 2016) and reviews that focus on multi-disciplinary work (e.g., Krause et al., 2011), the ecological role of the hyporheic zone (e.g., Boulton et al., 2010; Harvey and Gooseff, 2015; Stubbington, 2012), biogeochemical processes (Boano et al., 2014), and the influence of climate change on processes in the hyporheic zone (Zhou et al., 2014).

Lastly, the exchange between groundwater and the ocean is an important area of GW-SW interactions, and a major component of the global water cycle (e.g., Moore, 2010; Post et al., 2013). The interactions between fresh groundwater and saline ocean water play numerous important roles in the coastal zone. For example, submarine groundwater discharge (Fig. 1a, 1e) provides local sources of freshened water at the seafloor (Burnett et al., 2003; Taniguchi et al., 2019) and influences thermal and biogeochemical conditions important to the health of coastal wetlands, marshes (Wilson and Morris, 2012), lagoons, and estuaries (Befus et al., 2013; KarisAllen et al., 2022). Submarine groundwater discharge can be circulated (brackish) or fresh and is driven by processes across timescales ranging from seconds to millennia. These processes include waves, tides, seasonal oscillations in sea-level or water table elevation, and long-term sea-level changes (Taniguchi et al., 2019). One major difference between GW-SW interactions in freshwater and marine settings is the massive extent of the discharge area to the ocean, complicating efforts to measure submarine groundwater discharge directly and increasing reliance on tracers such as radium and radon (Garcia-Orellana et al., 2021). The role of submarine groundwater discharge has been extensively reviewed, including reviews focusing on biogeochemical processes (e.g., Santos et al., 2021) and methods to measure submarine groundwater discharge rates (e.g., Duque et al., 2020).

2.2. Anthropogenic factors influencing coupled GW-SW systems

The effects of both climate change (Ali et al., 2012; Uhl et al., 2022) and groundwater extraction (Vainu and Terasmaa, 2016; Yin et al., 2021) on GW-SW interactions have been documented and are expected to continue. For example, groundwater extraction and climate change have led to declines in the abundance of and biodiversity within

groundwater-dependent ecosystems such as springs (Gorelick and Zheng, 2015). Groundwater extraction can lead to stream depletion, with pumped 'groundwater' possibly including intercepted water that would otherwise discharge to a river and water directly from the river (Barlow and Hess, 1993; Hunt, 2014). Many analytical solutions to quantify stream depletion have been compiled for a range of conceptual models and hydrological settings as outlined in reviews by Hunt (2014), Huang et al. (2018) and Zipper et al. (2022). Additionally, the projected influence of climate change on GW-SW-related processes have been explored across a range of contexts, and continents, including studies in Australia (Zhou and Cartwright, 2021; Zhu et al., 2020), North America (Huntington and Niswonger, 2012; Levy et al., 2020), Asia (Karthé et al., 2015; Taft and Evers, 2016), Africa (Kahsay et al., 2018; Watson et al., 2019), South America (Gomez et al., 2022; Viola et al., 2015), and Europe (Chen et al., 2018; Pulido-Velazquez et al., 2015).

Groundwater and surface waters can either be a source or the receiving environment for contaminants (Tran et al., 2015). These contaminants include excess nutrients (Santos et al., 2021), contaminants such as the 'forever chemicals' per- and polyfluoroalkyl substances (PFAS) (Briggs et al., 2020; Pétré et al., 2021), trace organic contaminants (e.g., Hamdhani et al., 2020; Heeb et al., 2012), pharmaceuticals (Lapworth et al., 2015), heavy metals (Luo et al., 2022), and pesticides (Welch et al., 2019), amongst others. Concentrations of many contaminants in groundwater can be orders of magnitude higher than in surface waters; thus even small groundwater inputs can result in large fluxes (Conant, 2004). For example, submarine groundwater discharge can be a major contributor of nutrients to the ocean, and, in some cases, far exceed the nutrient loads discharged to coastal waters by rivers (Tait et al., 2023). Similarly, phosphorous transported via groundwater can lead to risk of lake eutrophication (e.g., Kazmierczak et al., 2021; Meinenmann et al., 2015). The contribution of contaminants can be diffuse, or via preferential pathways (Moore et al., 2023; Sawyer et al., 2016a). One particular concern is that contaminants can be stored in soils or groundwater systems for decades (or longer) before being discharged to surface waters, leading to long lag times between a pollution event occurring and when negative effects are seen in surface waters (Sanford and Pope, 2013; Tait et al., 2014; Tesoriero et al., 2013). The management of contaminated groundwater discharge is challenging because once contaminants enter the subsurface, the timing, potential attenuation through the sediment profile, and pathway of discharges are complex and highly heterogeneous (Cartwright and Irvine, 2020; Conant et al., 2004; Shishaye et al., 2021). Lagged delivery of nutrients from groundwater to surface waters is particularly widespread and prevalent across agricultural regions, where groundwater-delivered nutrients can significantly delay effects of nutrient management actions on water quality (Sanford and Pope, 2013; Van Meter et al., 2018).

Nutrients may be significantly transformed or attenuated within groundwater systems (Rivett et al., 2008). The GW-SW interface is particularly important for nutrient attenuation because the high reactivity of near-surface sediments surrounding streams, rivers, lakes, and wetlands reduces excess nutrient loads and pharmaceutical inputs at the distal end of groundwater flow paths (e.g., Hillebrand et al., 2015; Lewandowski et al., 2015), though preferential flow paths may bypass reactive sediments (Hester and Fox, 2020). Nitrogen dynamics within the hyporheic zone have been studied extensively (as reviewed by Zhao et al., 2021), and it is well established that hyporheic exchange can regulate nitrogen removal by stream and river ecosystems (e.g., Hall et al., 2009; Hill and Lymburner, 1998). The distribution of available carbon for microbial nitrogen removal and retention pathways as well as hydrologic residence times within the hyporheic zone control the magnitude and spatial pattern of GW-SW-driven nutrient removal (Zarnetske et al., 2011). Given the importance of hyporheic exchange in nitrogen removal, stream restoration research and practices have evolved to restore hyporheic exchange to enhance nitrogen removal (Merill and Tonjes, 2014).

2.3. Quantifying GW-SW interactions and further data interpretation

A wide range of methods are available to quantify GW-SW exchange rates and the key parameters that control processes, over point, reach and basin scales (Hammett et al., 2022; Kalbus et al., 2006; Ma et al., 2024). The breadth of techniques and approaches to quantify GW-SW exchange is due to the wide variability of settings where GW-SW interactions occur and the broad range of disciplines that study GW-SW interactions. For example, in some instances, GW-SW exchange can be measured directly at the point scale using seepage meters (Murdoch and Kelly, 2003; Rosenberry and Morin, 2004), or at the reach scale in riverine contexts using differential gauging (McCallum et al., 2014; Ruehl et al., 2006). Other approaches are based on Darcy's law (Cardenas and Zlotnik, 2003b; Chen et al., 2009a), isotope-based tracer methods (e.g., Cranswick et al., 2014; Gilfedder et al., 2015), naturally occurring or injected noble gasses (Blanc et al., 2024; Brennwald et al., 2022, 2016), reactive tracers (e.g., Höhne et al., 2022; Knapp et al., 2017), or the use of heat as a tracer (e.g., Banks et al., 2018; Koch et al., 2016; Ruehl et al., 2006; Schneidewind et al., 2016). In addition to the broad overview of techniques to quantify GW-SW exchange rates by Kalbus et al. (2006) and Ma et al., (2024), many techniques are further detailed in dedicated reviews (e.g., Cook, 2013; Irvine et al., 2017; Kalbus et al., 2006; Rau et al., 2014), and the methods-selection tool of Hammett et al. (2022). Recent advances have produced a new suite of tracing techniques to inform water origins and GW-SW exchange rates. These advances include the ability to measure noble gasses (e.g., He, Kr, Xe) in situ using portable mass spectrometers (Brennwald et al., 2016). This has allowed the use of noble gasses in injected gas tracer tests to inform a range of processes (e.g., Blanc et al., 2024; Brennwald et al., 2022; Weber et al., 2019). Microbial source tracking is another relatively recent addition to the suite of approaches that can identify the origins of contaminants and has been used to identify bacteria sources in GW-SW systems (e.g., Duvert et al., 2019; Neave et al., 2014).

Geophysical methods are one of the more recently developed tools for exploring GW-SW interactions, where electrical resistivity methods are probably the most widely used. For instance, in one of the earliest published studies, Acworth and Dasey (2003) used electrical methods to image the naturally occurring exchange of seawater with a freshwater sand aquifer in a tidal stream. Crook et al. (2008) mapped subsurface sediment deposits in two streams using electrical resistivity, which they used to estimate hyporheic exchange based on the distribution of alluvial deposits. Interactions (recharge and salinisation) between surface seawater flooding and shallow coastal aquifers have also been mapped using electrical resistivity instruments (Cantelon et al., 2022). Electrical resistivity has been used as a process-based tool for mapping GW-SW exchanges. Singha et al. (2008) demonstrated numerically that time-lapse electrical resistivity could be used to quantify hyporheic exchange processes by comparing bulk electrical resistivity and in-stream fluid conductivity measurements during conductive tracer tests. Where tracer-laden surface water exchanges with groundwater, there is an increased hyporheic zone fluid conductivity, which persists longer than that in the stream and is imageable with electrical resistivity. Since that synthetic test, many field studies have paired electrical resistivity and in-stream tracer experiments (Cardenas and Markowski, 2011; Doetsch et al., 2012; Ward et al., 2014, 2010). We note that the resolution of electrical resistivity surveys and the subsequent influence of the reliability on our understanding of hyporheic zone processes are rarely explored in detail. For example, Zhang et al. (2023a) demonstrate how well electrical inversions could be expected to resolve various changes in a GW-SW system using tracer tests coupled with electrical resistivity in a series of numerical experiments.

Remote sensing tools have also been widely used in GW-SW interactions research (Brunner et al., 2017), and the field is expanding rapidly. For example, thermal infrared sensors mounted on drones (e.g., Watts et al., 2023), helicopters (e.g., Tamborski et al., 2015), airplanes (e.g., Torgersen et al., 2001), and satellites (e.g., Jou-Claus et al., 2021)

have been used to identify zones of focused groundwater discharge that, depending on the season, manifest as cold- or warm-water plumes in surface water bodies. The spatial coverage and spatial resolution of such surveys are often inversely related. While drones provide a new, inexpensive alternative for mapping small aerial footprints at high resolution, there are ongoing challenge of thermal drift that limit sensor accuracy (O'Sullivan and Kurylyk, 2022). Remote-sensing studies often focus on mapping springs and seeps to identify thermal refuges for aquatic species (e.g., Dugdale et al., 2013), or to identify potential locations of focused contaminant loading to surface waters (Johnson et al., 2008). Other remotely sensed data types indicate vegetation health and flooding extent (Doody et al., 2014), and surface water elevation (Chen et al., 2020), which can be influential to, and influenced by, GW-SW interactions.

Field data collected to inform GW-SW interactions processes are often coupled to numerical models of groundwater, surface water, or coupled GW-SW processes to allow further interpretation of system behaviour. The widely used MODFLOW codes (e.g., Langevin et al., 2017) include various packages that can represent GW-SW interactions. For example, the RIV package can be used to represent rivers in a simplistic way using a conductance term and head differences to determine if groundwater 'discharges' to a river (i.e., is removed from the model). Surface water models including MIKE SHE (DHI, 2023), HEC-RAS (Brunner, 2020), and SWAT+ (Bailey et al., 2020; Chawanda et al., 2020) include representations of groundwater and GW-SW interactions processes. Additionally, there are approaches to couple surface and groundwater models, as well as model codes that simulate both surface water and groundwater processes in a fully coupled way, including HydroGeoSphere (Aquanty, 2015), GSFLOW (Markstrom et al., 2008) and ParFlow (Maxwell et al., 2023). However, not all investigations require numerical models, and various analytical solutions exist to assist with the additional interpretation of field data. For example, analytical solutions are widely used to interpret exchange fluxes from temperature (Hatch et al., 2006; Lin et al., 2022; McCallum et al., 2012; Schneidewind et al., 2016; Shi et al., 2024) or to quantify impacts of groundwater extraction on surface water bodies (e.g., Hunt, 2014; Zipper et al., 2022). Machine-learning approaches are also becoming more widely applied to investigate GW-SW interactions (e.g., Basu et al., 2022a; Husic et al., 2022; Yang et al., 2019).

3. Literature review methods

We used the Scopus database to search for GW-SW interactions-related research. The search criteria included Article title, Abstract and Keywords to identify research outputs from 1970 to 2023. The study period was restricted to 1970 to facilitate analyses by decade, and because only a small number of articles per year (i.e., 1–6) were available in the Scopus database prior to 1970. There is substantial heterogeneity in the terms used to describe GW-SW interactions research. As such, three groups of terms were used in the search; (1) terms that relate to groundwater or aquifers, (2) terms that relate to surface water bodies, and (3) terms that relate to "interactions" or associated processes (Fig. 2). The search was intended to be broadly representative, rather than exhaustive.

Following the original search, results were restricted to academic journal papers (i.e., excluding book chapters, conference abstracts, corrections and retracted papers). Bibliometric studies often omit papers published in languages other than English (Jackson and Kuriyama, 2019). However, the outputs of the Scopus search typically had abstracts and keywords in English even for papers published in other languages, and these papers were retained for further analysis. De-duplication was run on each output file. In addition to the main data extraction from Scopus, summary files were extracted using the "Summary Filter Count" option.

Bibliometric analyses were performed using a range of analytical approaches. Raw Scopus outputs were analysed using Python (Python

Groundwater terms

(groundwater* OR aquifer* OR "ground water*" OR "ground-water*")

AND

Surface water terms

("surface water*" OR river* OR stream* OR wetland* OR floodplain* OR "flood plain*" OR spring* OR lake*)

AND

Interactions terms and/or further context

(interaction* OR hyporhe* OR riparia* OR "submarine groundwater discharge" OR "sub-marine groundwater discharge" OR "SGD" OR "baseflow*" OR "base flow*" OR "base-flow*" OR "exchange*" OR "flux*").



Results of search

Fig. 2. Search terms used in the Scopus Article Title, Abstract, Keyword search on the 1st of January 2024. The search was limited to papers published between 1970–2023, retaining the following paper types: Article, Review, Note, Editorial, Data Paper, and Letter.

Software Foundation, 2024), based on either publication year, authors, affiliations, or countries of origin. In addition to these simpler analyses, a user-driven Boolean keyword-based classification approach was used. This process used Author Keywords and Index Keywords, reclassified against user-generated lists of key terms (see Supporting Information Table S1). This process was necessary given the broad range of analogous Author Keywords used (e.g., groundwater-surface water interactions, surface water-groundwater interactions).

Citespace (Chen and Song, 2019) was used to conduct further bibliometric analyses based on relationships between papers by authors, keyword, or citation patterns. These investigations include Betweenness Centrality analyses to identify important papers that link subject areas and to show clusters of research, and Structural Variation Analysis to project which recently published papers are likely to be influential. Further details on the application of these techniques are provided below, and in the Supporting Information.

The Betweenness Centrality analysis was used to identify papers that indicate influence and control over the flow of information in a network of papers. This approach is predicated on the assumption that the conceptual associations between research papers can be modelled as a network of clustered papers with co-citation links between them. The analyses identify influential papers that serve as important links between different research areas within GW-SW interactions research. The process automatically groups the co-cited papers into clusters using an algorithm that clusters papers that share more co-citation links (Blondel et al., 2008). Betweenness Centrality approaches have been previously used to identify critical research papers in various areas of environmental studies (e.g., Shen et al., 2023; Wen et al., 2023; Zhang et al., 2023b).

Structural Variation Analysis is used to measure the intellectual values of a new research paper based on the amount and type of changes it exerts on the current research landscape at the time of its publication.

Structural Variation Analysis scores have been shown to be positively correlated with the paper's future high citation count (Chen, 2012) and have provided useful early indicators of future Nobel Prize-winning papers (Sebastian and Chen, 2021). Here, Structural Variation Analysis was used to identify "promising" recent papers before they accrue a substantial number of citations. The extent to which a paper is considered promising is measured by the new connections generated between previously unlinked research areas over timeframes of one and five years prior to the publication of the paper. Although not initially designed as filtering mechanisms, these analyses present one approach to help researchers navigate through the proliferation of GW-SW papers by highlighting important or promising papers.

The methods and results of additional CiteSpace analyses into 'author bursts', whereby authors have a 'burst' of citations over time periods of 1–5 years is presented in the Supporting Information.

4. Analyses of GW-SW interactions research

The original Scopus search returned a total of 23,077 documents. After removing non-academic journal articles, corrections, and retracted papers, restricting outputs between 1970 and 2023, and de-duplication, 20,275 journal articles remained for further analysis.

4.1. Trends in GW-SW interactions research

The number of GW-SW interactions papers published has steadily increased through time (Fig. 3), as similarly observed in other disciplines (e.g., Gxokwe et al., 2020; Jägerbrand et al., 2022; Padilla et al., 2018, see Fig S1, Table S2). In the case of GW-SW interactions research, the number of papers published approximately tripled from the 1970 s to the 1980 s, and again from the 1980 s to the 1990 s, with 164, 525, and 1566 papers, respectively (Fig. 3). From 2000 onwards, the number of GW-SW interactions papers has continued to expand rapidly, with 3980, 8829, 5211, papers published in the 2000 s, 2010 s, and 2020–2023, respectively. The maximum number of papers published in any given year (1396) was in 2021. While there is year-to-year variability in the number of publications, paper numbers post-2020 were likely impacted by COVID (Abramo et al., 2022), particularly in the case of laboratory and field-based research.

The increasing number of GW-SW interactions papers has resulted in trends in the number of authors and references per paper. For example, the average number of authors per paper increased from 1.5 in 1970, to a peak of 5.6 in 2022 (Fig. 4a). This increase is approximately linear from 1980 onwards, more than doubling throughout the study period. The average number of references per paper has also steadily increased, from less than 10 in 1970 to more than 60 from 2020 onwards (Fig. 4a). This increase is not surprising because of the increasing volume and

accessibility of literature and the increasingly interdisciplinary nature of GW-SW interactions research. The rising number of references per paper follows the trends identified across other disciplines, with an increasing number of references cited per article through time (e.g., Dai et al., 2021).

The citations data (Fig. 4a) show much more complex trends than for authors and reference data. The average citations/paper prior to 1980 is erratic from year to year, because of the low number of papers published during this period (see Fig. 3) and the relatively high percentage of papers that were never cited (Fig. 4b). Between 1980 and 2000, the average citations/paper steadily increased from 22 in 1980 to approximately 60, peaking at 79 in 1998 (Fig. 4a). From approximately 2000 onwards, the number of citations/paper steadily decreased (Fig. 4a). More fundamental studies were published in the 1990 s (e.g., Brunke and Gonser, 1997; Constantz et al., 1994; Harvey and Bencala, 1993; Winter, 1999) and the early-mid 2000 s (Hatch et al., 2006; Kalbus et al., 2006; Ruehl et al., 2006; Woessner, 2000), and a strong case can be made for the recent proliferation of publications reducing the impact per paper of more recent work.

Papers cannot be cited immediately because subsequent works need to make their way through the peer-review process. Accordingly, Fig. 4b presents citations/year data normalised by the number of years since the paper was published. This approach assumes whole years and rounds up, thus a paper published in December of any given year is slightly disadvantaged in this process. The normalised citation values (Fig. 4b) peak in 2019, with a value of ~ 4 citations/paper/year published.

The number of authors per paper has steadily increased since approximately 1990 (Fig. 4a). This pattern is also observed in other hydrologically similar investigations, including investigations into nitrate leaching (Padilla et al., 2018), groundwater remediation (Zhang et al., 2017), and river health (Wen et al., 2023); refer to Supporting Information for comparisons. With increases in authors/paper across multiple research fields, it is likely that increases in the number of authors per paper have more to do with the general trend to conduct research in larger networks within and across disciplines.

4.2. Document type of GW-SW interactions research

Journal articles can be published in a range of formats, with Article and Review formats being the most common and the most cited in GW-SW interactions research (30.3 and 70.0 mean citations per paper, respectively, Table 1). In the case of GW-SW interactions research, most papers follow the Article format (19,529, ~ 96 %, Table 1). Of the remaining papers, only 13 are identified as Data Papers (i.e., the focus of the paper is the collation of a large dataset). These Data Papers appear to follow the recent move towards open science and the increased accessibility of resources to host online datasets (e.g., Hydroshare (CUASHI,




Fig. 3. GW-SW interactions papers by year (bars) and cumulative research outputs (black line) from 1970 to 2023. Colours denote decades.




Fig. 4. Groundwater-surface water interactions research trends using the following details for any given year: Total papers (TP), total references (TR), total authors (TA), citation count (CC) and years since publication (YP). (a) Average citations/paper (CC/TP) and average references/paper (TR/TP) (primary y-axis) and average authors/paper (TA/TP) (secondary y-axis). (b) Normalised citations/paper, calculated as CC/TP/YP (primary y-axis), and uncited papers (%) (secondary y-axis).

Table 1

Groundwater-surface water interactions paper type, by decade. The mean citation column is mean citations by paper type, between 1970 and 2023.

Document Type	1970–1979	1980–1989	1990–1999	2000–2009	2010–2019	2020–2023	Total Papers	Mean citations
Article	162	521	1534	3832	8535	4945	19,529	30.3
Data Paper	0	0	0	0	1	12	13	1.1
Editorial	0	0	0	4	9	9	22	12.6
Letter	0	2	1	2	9	7	21	17.7
Note	0	0	5	13	14	13	45	18.4
Review	2	2	26	129	261	225	645	70.0
Total	164	525	1566	3980	8829	5211	20,275	25.0
Papers/day	0.04	0.14	0.43	1.09	2.42	3.57		

2024), Figshare (2024), ScienceBase (U.S. Geological Survey, 2024a)), with all Data Papers being published in or after 2019. The published Data Papers collated datasets on submarine groundwater discharge, radon and radium data (Duque et al., 2019; Wolfe et al., 2023), hydrochemistry (Ba et al., 2023; Pan et al., 2022; Rumuri, 2020), and stable isotopes (Kalvāns et al., 2020; Tetzlaff et al., 2023), typically with

a local or regional focus.

Although the GW-SW interactions papers collated in this study are admittedly broad in nature, the number of Review papers and the rate of increase in publications through the decades is remarkable. For example, on average, 1.09, 2.42 and 3.57 papers were published per day in the 2000 s, 2010 s and 2020 s, respectively. The fact that there are 645




Fig. 5. Top 10 subject areas of published GW-SW interactions papers, by percentage of papers published by decade. Subject areas as classified by the Scopus database. Subject areas outside of the yearly top 10 are grouped into “OTHER”. Black numbers denote the combined percentage of papers in Environmental Science, Earth and Planetary Sciences, Agricultural and Biological Sciences and Engineering (EEAE) papers for each decade.

Review papers, 225 of which have been published since 2020 (Table 1), further highlights the challenges of keeping up with the literature, even review papers (the irony of making this point in a review paper is not lost on us). The challenge of staying up to date with relevant research has also been documented in smaller, but rapidly growing disciplines (e.g., Tague and Brandt, 2023).

4.3. Scopus subject areas through time

Research in GW-SW interactions is often multi-disciplinary by necessity, given the broad range of processes that influence and are influenced by surface–subsurface hydrologic exchanges. Four fields—Environmental Science, Earth and Planetary Sciences, Agricultural and Biological Sciences, and Engineering (herein the EEAE subject areas)—contribute the largest number of GW-SW interactions papers (Fig. 5). The total contribution of the EEAE subject areas has consistently declined since the 1980 s. For example, the total contribution of the EEAE subject areas decreased from 93 % of papers in the 1980 s to 76 % of papers in the 2020 s (so far). Interestingly, Engineering is not featured in the top four subject areas in the 2020 s, following a large growth in the subject areas of Social Sciences, Biochemistry, Genetics, and Microbiology. Overall, the decadal trends in subject area (Fig. 5) suggest that GW-SW interactions research is becoming increasingly multi-disciplinary.

Many of the subject areas that make up a smaller percentage of GW-SW interactions papers are categorised into somewhat curious subject areas, including medicine. In this example, the research identified as having a medical focus generally studied the transport of various chemicals including antibiotics (e.g., Li et al., 2023) or PFAS (e.g., Tokranov et al., 2021) and legacy contamination events (e.g., Dwivedi et al., 2022). The percentage of studies that are outside of the top 10 (labelled here as “OTHER”, Fig. 5), increase from < 1 % in the 1970 s to 4.6 % in the 2020 s, further demonstrating the increasingly multi-disciplinary nature of GW-SW interactions research.

4.4. Journals publishing GW-SW interactions research

GW-SW interactions research has been published in over 2000

different journals between 1970 and 2023. The top 15 journals, by number of GW-SW interactions papers published (Fig. 6) collectively comprise 7584 (37 %) of the total GW-SW interactions papers considered in this analysis. The *Journal of Hydrology* has published the most articles, with 1543; this count did not include the relatively new *Journal of Hydrology X*. The count for *Groundwater* included both *Ground Water* and *Groundwater*, following the name change of the journal in 2013.

Many of the journals in the top 15 (Fig. 6) were founded during the timeframe of the study (1970–2023). For example, the first edition of *Science of the Total Environment* and *Hydrogeology Journal* was released in 1992. The first edition of *Environmental Earth Sciences* was published in 2009, and the first issue of *Hydrology and Earth System Science* was published in 1997. The most recently added journal from Fig. 6, *Water*, was first released in 2009. Despite its relatively recent release, *Water* ranks seventh on the number of GW-SW interactions publications, publishing a high volume of papers, with an overall lower citation rate per article (9.5 citations/paper) than other journals (e.g., *Journal of Hydrology*, *Water Resources Research* and *Hydrological Processes* receive 39.3, 52.9 and 33.3 citations/paper, respectively). The citations of these papers are at least partially affected by the age of the papers, and the time that they have had to be cited. However, it is worth noting that there is controversy surrounding whether or not *Water* is a predatory journal (MDPI, 2023; Predatory Reports, 2022). In contrast, *Environmental Science & Technology* papers attract higher citations, on average (64.9), than all other journals.

4.5. GW-SW interactions papers published by country

GW-SW interactions papers have been published by authors from at least 159 different countries between 1970 and 2023 (Fig. 7). The top five publishing countries over the study period, by total papers published, are the United States (6568, 23.4 %), China (3176, 11.3 %), Germany (1658, 5.9 %), Canada (1428, 5.1 %) and Australia (1290, 4.6 %). The top five publishing countries account for 50.3 % of the GW-SW interactions papers in the study period. The top 10 countries (also including the United Kingdom, France, India, Italy, and Spain) account for 66.6 % of GW-SW interactions papers from the study period. The number of publications produced per country is influenced by factors




Fig. 6. Publications data for GW-SW interactions research published between 1970 and 2023. (a) Publications by journal (y-axis), year range (x-axis), average number of citations per document (colour) and total number of publications (thickness). In panel (a), the length of the lines denotes the years where papers appeared from that journal in the Scopus search. (b) Boxplots of citations with 5 and 95th percentiles as the whiskers, 25th and 75th percentiles as the edges of the boxes, black lines as the 50th percentile, coloured line as the arithmetic mean citation, and grey numbers denoting the number of papers.




Fig. 7. Country contributions to GW-SW interactions papers published between 1970 and 2023. (a) Total number of GW-SW interactions papers for each country. (b) Total papers for each country per million people. Population data from Jan 2020 was used (United Nations, 2022). Countries with no data are shown in grey. (c) Country contributions (papers/year) over time for the top 10 publishing countries, with the inset panel showing papers from 2005 to 2023.

including population size and level of economic development, amongst others. Although we do not account for economic status, we do present paper outputs normalised by population (Fig. 7b). The top 10 countries, using the population normalised metric of papers per million inhabitants, were Monaco (162.0), Iceland (63.1), Switzerland (59.8), Denmark (51.1), Australia (50.5), New Zealand (49.3), Luxembourg (43.1), Sweden (38.6), Canada (37.8) and Estonia (33.1).

€.

Publishing rates of GW-SW interactions papers have varied

significantly across different countries (Fig. 7c). Until 2021, the US published the most GW-SW papers in any given year, with an approximately linear increase in the papers/year between 1990 and 2020 ($R^2 = 0.97$). In 2021, China passed the US in papers/year, with an almost exponential increase in the number of papers/year between 2000 and 2023 ($R^2 = 0.94$). Incidentally, the rate of research outputs from China seems to be less affected by COVID compared to other countries, i.e., post-2019. For example, there is an observable decline in GW-SW papers/year from the US, post-2020; however, declines over this period

also occurred in Germany and Australia, and research outputs in many other countries have been relatively constant (Fig. 7c). The decline in 2023 can be partially attributed to delays in some papers being listed on Scopus at the time of the search.

4.6. Identifying connections between papers using Betweenness Centrality analyses

To assess the importance of individual research papers beyond their citation counts, we applied a well-established measure in the field of network science called Betweenness Centrality (Freeman, 1977; Newman, 2005). The co-citation network of GW-SW interactions papers from 2000 to 2023 (Fig. 8) shows the output of the Scopus search as a network of research areas (clusters). Each cluster is labelled with the most salient terms automatically extracted by the Latent Semantic Indexing algorithm (Deerwester et al., 1990) from the titles of citing papers, showing distinct research subfields. For example, Cluster #0 *submarine groundwater* denotes the largest cluster, noting the prevalence of research in this area since 2000. Papers labelled by blue text (Fig. 8) are the top-four research papers with the highest Betweenness Centrality scores (see Table 2), with these papers providing strong links between research areas.

The links between individual papers and those that link the research clusters was identified using Citespace and shown in Fig. 8. For example, the large number of links between #2 “Hyporheic exchange” and #3 “Using heat” is clear, given the wide use of heat as a tracer to quantify exchanges in the hyporheic zone. However, some of the classifications differ from what would be the case if this process were done manually.

For example, the Sophocleous (2002) review is broad in nature and could fit in many categories presented here. Similarly, the Boulton et al. (1998) review, focused on the hyporheic zone, and covered concepts well beyond headwater streams (the cluster that it was assigned to). The challenge in categorising these papers, amongst others, is further demonstrated in Table 2, where some papers link large numbers of research clusters. For example, the Sophocleous (2002) review links seven and Boulton et al. (1998) links five research clusters. Although this could be an unsurprising result, the approach presented here demonstrates the ability of some papers, typically reviews, to link various aspects of GW-SW interactions research together.

There are notable exclusions from Table 2, given its focus on academic papers. There are highly influential reports that do not feature in the analyses presented here, including the report *Ground water and surface water – A single resource* by Winter et al. (1998). The report has been widely influential, because it provides overviews of key concepts and several U.S. examples of GW-SW interactions studies. As the title clearly states, the Winter et al. (1998) report makes the case for groundwater and surface water to be managed as a single resource, an approach that was not common at the time the report was published.

High BC score papers (Table 2, Fig. 8) identify opportunities for techniques to be applied to inform multiple processes and/or document impacts on GW-SW interactions systems. These papers are, in most cases, review papers, and are 10 to 25 years old, providing sufficient time to be cited. Review papers are also cited more than Articles, on average (Table 1). Non-reviews in Table 2 include the Cardenas (2008) paper that demonstrates the control of geomorphology on residence times, and the Cardenas (2009) paper that demonstrates the control of




Fig. 8. Betweenness Centrality analysis of GW-SW interactions papers between 2000–2023 for clusters 0 to 14. Betweenness Centrality analyses identify clusters of (nodes) papers, and papers that are important links between clusters. Named/ numbered clusters (as identified by the Citespace software) in descending order of its sizes: 0: submarine groundwater; 1: nutrient cycling; 2: hyporheic exchange; 3: using heat; 4: riparian zone; 5: groundwater quality index; 6: integrated hydrologic model; 7: geomorphology hydrology; 8: subterranean estuary; 9: shallow groundwater; 10: headwater stream; 11: riparian vegetation; 12: using tritium; 13: runoff processes; and 14: chemical component. The blue labelled papers partially identify the top papers by Betweenness Centrality scores and the size of their associated pink nodes indicates strength of Betweenness Centrality score. Table 2 provides further details on the top 10 papers by Betweenness Centrality score.

Table 2

Top 10 papers by Betweenness Centrality (BC) scores (see Fig. 8). These papers provide important links between and within research clusters. For cluster numbers and names, see Fig. 8. In addition to BC score, the square brackets include the number of citations in Google Scholar (GS) and Scopus (S) on June 14, 2024.

Rank: BC score [GS/S]	Paper reference	Title	Linked research clusters
1: 0.22 [165/ 213]	Cardenas (2008)	Surface water-groundwater interface geomorphology leads to scaling of residence times	0, 2, 3, 5, 7, 12
2: 0.20 [901/ 1520]	Anderson, (2005)	Heat as a ground water tracer	0, 2, 3, 5, 7, 10, 14
3: 0.12 [945/ 1546]	Boulton et al. (1998)	The functional significance of the hyporheic zone in streams and rivers	3, 7, 10, 11, 14
4: 0.11 [1417/ 2502],	Sophocleous (2002) ¹	Interactions between groundwater and surface water: the state of the science	2, 3, 4, 5, 7, 8, 10
5: 0.10 [234/ 380]	Fleckenstein et al. (2010)	Groundwater-surface water interactions: new methods and models to improve understanding of processes and dynamics	0, 1, 2, 6, 12
6: 0.08 [372/ 594]	Constantz (2008)	Heat as a tracer to determine streambed water exchanges	0, 1, 2, 3, 4, 6, 7
7: 0.08 [661/ 887]	Boano et al. (2014)	Hyporheic flow and transport processes: Mechanisms, models, and biogeochemical implications.	0, 1, 2, 6, 9
8: 0.08 [1132/ 1835]	Brunke and Gonser (1997)	The ecological significance of exchange processes between rivers and groundwater.	4, 7, 10, 11, 13
9: 0.07 [151/ 218]	Cardenas (2009)	Stream-aquifer interactions and hyporheic exchange in gaining and losing sinuous streams.	0, 1, 2, 3, 5, 6
10: 0.07 [463/ 594]	Santos et al. (2012)	The driving forces of porewater and groundwater flow in permeable coastal sediments: A review.	0, 1, 6

¹ The Sophocleous (2002) article was flagged for plagiarism (e.g., Voss, 2015).

bedforms on in-channel flow fields for gaining and losing conditions. The Fleckenstein et al. (2010) paper, although labelled as an 'Article' in Scopus, is a short review with two main components: (a) an overview of GW-SW interactions research, summarising the influence of reviews by Sophocleous (2002), Winter (1999), and Woessner (2000), and (b) a high-level summary of the content from a special issue into GW-SW interactions, of which it is the first article.

5. Recent trends and future opportunities in GW-SW interactions research

5.1. Projecting important papers using Structural Variation analysis

Here, we seek to identify papers that may substantially influence on the discipline in the near future using the Structural Variation Analyses (SVA) (Table 3). In SVA analyses, papers that create important yet previously absent links between other groups of existing research papers may have a higher probability of receiving more citations and being seen as transformative (Chen et al., 2009b; Sebastian and Chen, 2021). Papers that introduce new co-citation links in ways that significantly altered the prior distribution of Betweenness Centrality scores of all existing papers (see Fig. 8) are likely to be scientifically more promising than those that did not (Sebastian and Chen, 2021). SVA analyses were run using papers published during the last five years (2019–2023), and the keyword grouping (Table S1) that corresponds with the research topics of coastal processes, nutrients and nutrient cycling, the hyporheic

Table 3

Projected future impactful papers from Structural Variation Analysis from selected research clusters from Fig. 8. Keyword groups are documented in Table S1.

Projected high impact papers	Description of research
Keyword group: Coastal zone Claude et al., (2019)	Cluster number: #0 – Submarine groundwater
Xiong et al., (2022)	Use of Radium mass balance to detect SGD in karst settings.
Martínez-Pérez et al. (2022)	Environmental management and seawater intrusion in China.
Keyword group: Nutrients Wallace et al. (2020)	Multi-disciplinary approach to characterising SGD and seawater intrusion.
Ping et al., (2023)	Cluster number #1 – Nutrient fluxes
Ping et al. (2021)	Reactive transport modelling in a coastal GW-SW system and nitrification processes.
Keyword group: Hyporheic zone Singh et al. (2022)	Numerical modelling to investigate the role of particulate organic carbon deposition on nitrate reduction in the hyporheic zone.
Herzog et al. (2019)	The role of bank storage processes in the removal of nitrate from surface waters.
Wilhelmsen et al. (2021)	Cluster number: #2 – Hyporheic exchange
Keyword group: Riparian zone Sterte et al. (2022)	Coupled surface–subsurface model with reactive transport to demonstrate effect of peak flow events on denitrification.
Blaurock et al. (2022)	Numerical modelling analyses to demonstrate nested hyporheic exchange in headwater streams.
Nogueira et al., (2021)	Laboratory and numerical modelling study to demonstrate the influence of log jams and branching on hyporheic exchange.
Ping et al. (2021)	Cluster number: #4 – Riparian zone
	Relating groundwater transit times to dissolved organic carbon and carbon budgets in surface waters.
	The role of riparian microtopography and dissolved organic carbon and dissolved organic matter in a forested catchment.
	The role of transit times and temperature in aerobic respiration and denitrification processes.
	The role of bank storage processes in the removal of nitrate from surface waters.

zone, and the riparian zone.

Where most of the papers identified in Table 2 were reviews, the studies identified in Table 3 are largely studies of processes and field sites. For example, numerical modelling features prominently in the studies in Table 3 (e.g., Herzog et al., 2019; Sterte et al., 2022; Nogueira et al., 2021; Ping et al., 2023; Singh et al., 2022; Wallace et al., 2020; Wilhelmsen et al., 2021). Many papers also describe field-based studies (e.g., Blaurock et al., 2022; Claude et al., 2019; Sterte et al., 2022; Martínez-Pérez et al., 2022; Nogueira et al., 2021; Ping et al., 2021; Wallace et al., 2020).

5.2. GW-SW interactions keyword trends

The review to this point has focused on providing a summary of the research field and publishing trends. From herein, the focus switches to be forward looking. As part of this transition, we first look at research outputs through time, with a focus on recent trends through the assessment of the published papers in key topics. Normalising by the number of GW-SW interactions papers each year facilitates the analysis of research trends and mitigates the influence of the overall annual increase in scholarly output (Fig. 9). The increased breadth of GW-SW interactions research (e.g., Fig. 5) can also make the interpretation of these results challenging, particularly for keywords that become less prevalent through time. Nonetheless, keywords that increase through time do so despite the increasingly broad research focus of GW-SW interactions research.

Most GW-SW interactions studies focus on rivers and streams




Fig. 9. GW-SW interactions research trends identified by grouping Author Keywords (see Table S1 for details on the Keyword grouping approach) from 1980 to 2023. The panels show publishing trends by (a) surface water type, (b) impacts investigated, (c) data collection and data analysis methods, (d) specific research focus (K denotes hydraulic conductivity), and (e) emerging topics (PFAS denotes Per- and polyfluoroalkyl substances, TrOCs denotes Trace Organic Compounds, and POPs denotes Persistent Organic Pollutants). The percentages for any given year do not necessarily add to 100, as papers can include multiple keywords, and/or papers may not include any of the keywords for any given panel. 1970 s papers are omitted because the outputs (as a percentage) are erratic during this period due to the low number of papers published.

(Fig. 9a), with keywords that indicate river-related research increasing since 1995. Studies of the coastal zone show a steady increase until 2005, after which they remain relatively constant. Papers investigating drought and climate have increased over the timeframe of the papers considered here (1970–2023), with a particularly large increase starting in 2016 (Fig. 9b). A noteworthy percentage of papers (~40 %) include some mention of contamination. This has been reasonably consistent since the 1990 s. From the investigation techniques (Fig. 9c), increases in the use of machine learning and remote sensing are evident. For example, the increased use of remote sensing provides opportunities to upscale site-specific findings to broader scales to further inform management. Use of machine-learning approaches has also steadily increased (Fig. 9c). The observed decline in the relative number of papers with terms relating to ‘numerical modelling’ since 2010 was unexpected. This is surprising given the continued increase in computational power and wide availability of numerical models capable of simulating fluid flow, solute and heat transport, including coupled surface–subsurface models as well as open access packages.

We also identify what we consider to be emerging research areas, noting that there is no formal definition for these areas (Fig. 9e). For example, contaminants, including PFAS, Trace Organic Compounds (TrOCs), and microplastics (among other chemical groups not highlighted in our analysis) are beginning to be featured in GW-SW interactions research. The dynamic conditions of GW-SW exchange may be particularly important in the fate and transport of PFAS given its sorption and speciation characteristics at multiple interfaces between groundwater, surface water, and soils (Divine et al., 2023). Open questions about how quickly PFAS are moving from aquifers to streams and the time needed to flush PFAS from aquifers remain unanswered (Briggs et al., 2020; Pétré et al., 2021). For example, fluctuations in GW-SW fluxes drive biogeochemical changes that can lead to seasonal variations in perfluoroalkyl acid concentrations and precursor loss (Tokranov et al., 2021). To date, there has been limited numerical simulation of GW-SW transport of PFAS, undoubtedly because modelling codes capable of simulating PFAS transport through variability saturated media have been only recently published (Brusseau et al., 2020; Guo et al., 2020; Silva et al., 2021).

The use of uncrewed vehicles, including aerial or aquatic vehicles (UAVs, or drones) is steadily advancing many research fields within the geosciences, including the study of GW-SW interactions (e.g., Mangel et al., 2022; Fig. 9e). Importantly, UAVs allow for precise, georeferenced surveys across landscapes or water bodies that are difficult to traverse by boat or by foot, including protected or contaminated water bodies, zones with unexploded ordinances, boggy or snowy terrain, and steep slopes. Although UAVs have been used for several years to construct digital elevation models using structure-from-motion photogrammetry for the study of GW-SW exchange (e.g., Pai et al., 2017), the number and type of sensors that can be installed on commercially available UAVs, along with UAV payloads, continues to expand. Given the ability of UAVs to capture high resolution data over large areas, we expect that this is an area of future research opportunities.

5.3. Future research opportunities

Using recent trends as presented above and our own insights, here we outline future opportunities in GW-SW interactions research. Although the fundamentals of hydrology and hydrogeology are largely unchanged since the 1970 s, for example, the vast majority of the *Groundwater* textbook by Freeze and Cherry (1979) remains current, the many disciplines that now participate in GW-SW interactions research have led to a proliferation of techniques, technologies and approaches. Indeed, over 20 years ago, Schwartz and Ibaraki (2001) suggested that in the field of hydrogeology “the number of truly impactful problems has dwindled to just a few.” However, through GW-SW interactions research, hydrogeology may still provide truly impactful research opportunities. The increased size of the GW-SW field since 2001 (Figs. 3, 5) could have led to issues of

citation dilution as discussed by Schwartz and Ibaraki (2001). Nonetheless, many knowledge gaps in GW-SW interaction research remain. Preferential flow pathways and heterogeneities in aquifers and bed materials continue to pose a challenge to quantifying or upscaling GW-SW interactions (e.g., Claude et al., 2019). The role that GW-SW interactions can play in removing contaminants is an area of active research (e.g., Ping et al. 2021). Climate change and groundwater extraction will continue to alter GW-SW systems (e.g., Benz et al., 2024; Petpongpan et al., 2022), potentially driving changes in stream drying that affect management (e.g., Walker et al., 2021). Similarly, sea-level rise will likely continue to pose problems in the coastal zone (e.g., Martínez-Pérez et al., 2022) by driving saltwater intrusion and potentially altering the geochemistry of coastal aquifers and receiving waters. This is clearly a partial list of challenges; and additional GW-SW interactions research challenges will likely develop in the future. Field-based research using existing and new techniques, sensing tools, and interpretive methods could be used to address these challenges. The discussion below outlines areas that we believe show promise as future directions in GW-SW interactions science.

5.3.1. Drones and uncrewed vehicles

Examples of recently developed commercial UAV sensors with applications in GW-SW interactions studies include thermal infrared, Light Detection and Ranging (LiDAR; e.g., Bandini et al., 2017), multi- and hyper-spectral imaging (e.g., Pai et al., 2017), electromagnetic, ground-penetrating radar, and remote sampling of surface water Straight et al. (2021). Many of the signals measured with UAV instruments could be used to parameterise models (e.g., topography for integrated surface–subsurface models) and detecting and characterising the distinctive features and conditions of groundwater discharge zones (e.g., temperature, biogeochemistry, wetted channel length). For example, Pai et al. (2017) used drone-based Normalised Difference Vegetation Index to explore river corridor vegetation health as related to subsurface flow through meanders in semi-arid terrain, while Dugdale et al. (2019) used UAV-based vegetation height metrics to parameterize stream-temperature models. Two reviews of UAV as applied to hydrologic systems broadly can be found in Vélez-Nicolás et al. (2021) and Acharya et al. (2021). High-resolution visual imagery from UAVs may also be used to detect zones of focused groundwater discharge in colder regions that manifest as ice-free areas in rivers and streams during winter months, a phenomenon that has been demonstrated at lower resolution from satellite imagery (O’Sullivan et al., 2019). This list of sensors, many of which can now be purchased commercially, include several geophysical instruments that enable hydrogeophysical studies of GW-SW interactions. We believe that the high-resolution, georeferenced data that can be collected with these UAVs sensors has the potential to be transformative for mapping and modelling GW-SW interactions.

5.3.2. Scaling with the aid of remote sensing

Stream reach-scale (a few km or shorter) GW-SW field investigations dominate the literature but are universally challenged in providing system-scale (catchment to basin) information of use to water-resource managers. Near-surface geophysical methods provide spatial continuity and hydrogeological context for point data (e.g., see Binley et al., 2015; McLachlan et al., 2017), but they are rarely collected at scales beyond reaches and transects, with some exceptions (e.g., Briggs et al., 2022; Lane et al., 2020). In contrast, data describing processes of GW-SW interactions collected by occupied aircraft and satellites are being paired with emerging AI-based analyses to push the spatial limitations of ground-based surveys. For example, high-resolution hyperspectral data acquired by occupied aircraft offer temporal snapshots of vegetation type and health (Chadwick et al., 2020), which is influenced by GW-SW exchange in more arid environments and can indicate controlling hydrologic processes (Meyers et al., 2021). Thermal infrared data are now routinely collected by aircraft to map zones of groundwater discharge and thermal heterogeneity across all surface water types (Steel et al.,

2017). Airborne electromagnetic imaging (AEM) data can be used to locate zones of surface water-groundwater connectivity at basin scales when either surface or groundwater is saline (Ball et al., 2015). However, occupied aircraft campaigns are costly and not often repeated at the timescales of seasonal change.

Satellite-based data, such as those from the Landsat satellite (U.S. Geological Survey, 2024b), provide global data every few weeks going back decades and are often freely available. Analysis products can indicate surface water inundation extent (Jones, 2019) and identify wetlands influenced by groundwater extraction (Heintzman et al., 2017). Landsat and AEM data have been combined to understand the impact of shallow permafrost thaw and arctic lake extent due to changes in lake-groundwater exchange (Jepsen et al., 2013; Rey et al., 2019). Groundwater discharge has been mapped using satellite thermal infrared data at regional scales in terrestrial (Sass et al., 2014) and ocean (Jou-Claus et al., 2021) environments. Although many surface water features are not currently resolvable due to the coarse spatial resolution (30–100 m pixels) and reduced accuracy of thermal infrared data due to edge effects, several higher-resolution commercial thermal infrared data sets are expected to become available in the coming years. New data types such as Surface Water and Ocean Topography (e.g., Altenau et al., 2021) are being made broadly available and have widespread applicability for estimating hydraulic gradients between surface and groundwater when paired with water-table elevations. New processing algorithms to infer large-scale algal blooms driven by the exchange of nutrient rich groundwater from satellite spectral imaging are allowing hindcasting of how such blooms have changed in timing and extent using publicly available time-series data (e.g., Cheng et al., 2020). Finally, remote sensing of river discharge from drones, occupied aircraft, and satellite platforms is becoming more accurate (Gleason and Durand, 2020), creating the potential for large-scale analysis of river gains and losses in remote areas across seasons. The citizens of many countries have made a collective investment in satellite-based remote sensing implemented by various governments, and the resultant publicly available data provide opportunities to inform GW-SW interactions science at management-relevant scales across the globe.

5.3.3. Real-time monitoring networks, low-cost sensors and application of newer tracers

The establishment and growth of real-time monitoring networks for groundwater and surface water is a recent development that can enable informed, real-time decision making for water-resources management (Zhou et al., 2018). Although commercial options for such real-time systems can be cost prohibitive for procurement and data management, the growth of “do-it-yourself” Internet-of-Things technologies and controllers (e.g., Arduino, Raspberry Pi) has led to the emergence of low-cost, often real-time monitoring of water levels, temperature and water quality that may enable more high-resolution or large-scale monitoring of GW-SW interactions because the low cost enables scalability (e.g., Calderwood et al., 2020; Drage and Kennedy, 2020; Espinoza Ortiz et al., 2023). The widespread use of real-time data can provide multiple benefits, including the potential for near-real-time numerical modelling. The availability of new wireless communications options, including Starlink, can also allow the use of real-time data to be measured in areas that previously did not have access to sufficient internet and phone-based services (e.g., Dinis et al., 2022).

The use of tracers has been a mainstay in the hydrological sciences. The wide array of existing approaches will likely continue to inform GW-SW interactions. The recently developed ability to measure noble gas concentrations in situ (e.g., Brennwald et al., 2016) is still in its relatively initial stages, though it is well suited for ‘Lagrangian’-type continuous surface-water parameter surveys collected at ambient streamflow velocity from watercraft (e.g., Hensley et al., 2020). The rise of DNA-based tracing techniques (e.g., Pang et al., 2020) also shows promise to inform GW-SW interactions. With the increasing array of chemicals in use, new environmental tracers may arise, as was the case

with chlorofluorocarbons (e.g., Hendlin, 2021). These approaches, amongst others, will expand the ability to inform various GW-SW interactions and associated processes. Future research could use these existing techniques and develop new tracing techniques. A continuing challenge will be to develop approaches that can be used to upscale the often point or local-scale investigations over which tracer studies are often applied.

5.3.4. Flow paths and contaminants

There are a wide range of contaminants that occur at the GW-SW interface and are receiving increased research attention (e.g., microplastics, PFAS, TrOCs, pharmaceutical products, and pathogens, amongst others). Although research examining the medical effects of the human exposure to some contaminants (e.g., PFAS, microplastics) there is in its preliminary stages, documenting the prevalence and extent of these contaminants in GW-SW systems is becoming increasingly important for informing management activities. Microplastics and PFAS are now ubiquitous in the environment; microplastics have been found in groundwater via karst/cave systems (Balestra and Bellopede, 2022; Panno et al., 2019), while PFAS chemicals have been found most everywhere (Johnson et al., 2022). Although the class of PFAS compounds is often referred to as ‘forever chemicals’, important reactions of precursors have recently been documented at the GW-SW interface (e.g., Tokranov et al., 2021). Challenges remain in understanding contaminant flow pathways in GW-SW systems due to heterogeneities in the subsurface, dominance of preferential flow processes, and the complex potential combination of biotic and abiotic chemical reactions that tend to occur in interface sediments. Understanding the surface or subsurface flow paths that contaminants take from source to receptacle and how those pathways are influenced by GW-SW exchange has long-term implications for human and ecosystem health given the ‘legacy’ effects of contaminants in groundwater systems (Basu et al., 2022b).

5.3.5. Knowledge transfer across disciplines

This review has highlighted the increasingly multi-disciplinary nature of GW-SW interactions research (e.g., Figs. 5 and 8). International collaborations on multi-disciplinary projects spur innovations that quickly advance the science of GW-SW interactions despite the challenges posed by the differences in language among researchers (Singha et al., 2024). Multidisciplinary collaborations provide opportunities for the translation of knowledge across disciplines and are often required to address complex research questions. This point has been made elsewhere for GW-SW systems, with suggestions for increased collaboration among hydrological scientists and ecologists (e.g., Humphreys, 2009), climate scientists (e.g., Hannah et al., 2007) and biogeochemists (e.g., Smith et al., 2008), and a wide range of other disciplines including non-scientific, disciplines. With increased pressure on water resources from contamination, overextraction, and climate change, we reiterate that ongoing collaborations can help promote existing knowledge transfer across disciplines. We suggest that GW-SW interactions is a natural topic for such collaborations because this topic spans the surface and subsurface aquatic environments that are often studied in disciplinary silos.

5.3.6. The future of numerical modelling and AI

Advances in computational power have outpaced our ability to measure field parameters at appropriate spatiotemporal resolution to characterise many processes relevant to GW-SW interactions. As a result, process-based models still use simplified representations of hydrogeological systems, particularly for subsurface properties (Thornton et al., 2022). Brunner and Simmons (2012) highlight the benefit of having models with a wide range of functionality, that can be used where the necessary data are available. Despite the challenges, large-scale (e.g., continental) models that include groundwater and surface-water processes in some manner are available (Cosgrove et al., 2024; Maxwell et al., 2015; Vaze et al., 2013) or are under development (Aquanty, 2024). Given the challenges in measuring model parameters,

Beven (2018) highlights that using models to test hypotheses remains a valuable pursuit. For example, studies exist where all models tested were ultimately rejected due to poor predictive abilities (Choi and Beven, 2007; Dean et al., 2009). This testing of multiple conceptual models or the use of heterogeneous realisations of hydraulic properties are approaches often used in hydrogeological investigations (e.g., Enemark et al., 2019). Although ultimately more computationally intensive, hypothesis testing can be informative about what is and is not, known about the region of interest.

Artificial Intelligence (AI) approaches (including a broad range of machine-learning techniques) are being used to assist the parameterisation of complex process-based numerical models. For example, Barclay et al. (2023) used guided deep-learning approaches to inform river-temperature models, Wunsch et al. (2022) conducted a national assessment of groundwater storage under different climate scenarios, and Yang et al. (2020) showed that model predictions of daily streamflow improved after using synthetic datasets and machine-learning approaches. Neural networks, deep learning, or unsupervised algorithms have also been used to assess groundwater quality; however, the majority of studies focus on a narrow band of chemical compounds such as nitrate, which accounts for nearly one half the studies (Haggerty et al., 2023). Unsupervised machine learning has also been used to cluster inverted electrical resistivity data to map biogeochemically important regions in a hyporheic system (Singley et al., 2022). These examples highlight data or analysis gaps in the types of field observations that could be filled to more fully understand GW-SW interactions. Where additional data can be collected, they should. Otherwise, the use of machine learning and AI approaches to address data gaps has already shown to be a possible approach to assist the understanding of GW-SW systems.

Just as some are raising the possibility that AI will be the end of computer programming as we know it (e.g., Welsh, 2023), we emphasise here that although numerical models will likely continue to be an important component of GW-SW interactions research, how numerical models will be implemented and applied in the future is uncertain in the face of rapidly developing AI approaches. Regardless of the exact approach to running numerical models, future modelling efforts that inform GW-SW interactions on appropriate scales could be used to inform water-resources management. Numerous important processes have been elucidated through decades of GW-SW interactions research, but predictive power and precision related to management-relevant themes, such as nutrient fate and transport above the reach scale, remains relatively low. This could mean that there is still a lack of process understanding and/or the existing models do not include the complexity of processes. Broad overviews of potential AI applications provide insights into AI opportunities in the hydrological sciences (e.g., Foroumadi et al., 2023; Halloran et al., 2023; Irvine et al., 2023). However, we are in the early stages of large-scale AI applications, and we hypothesise that AI will continue to have a wide-reaching influence on future research, across all disciplines.

5.3.7. Managing GW-SW systems

Ultimately, measured, modelled, and interpreted data are needed to understand hydrological systems and facilitate water-resources management. With the increased water demands to sustain human populations, agriculture, and energy production paired with the prevalence of large-scale contamination of many water resources, developing more effective management regimes for GW-SW systems is crucial (Gorelick and Zheng, 2015). The 2023 U.S. Supreme Court decision that the U.S. Clean Water Act only holds in continuously flowing surface water bodies has implications for interactions of groundwater and surface water systems (e.g., US EPA, 2023). Although the limitations and challenges in managing groundwater have been identified including in how to set effective trigger levels to minimise localised impacts (e.g., Rohde et al., 2024; Werner et al., 2011), broad management frameworks that acknowledge GW-SW as a single connected system are lagging despite

longstanding calls for conjunctive management (e.g., Winter et al., 1998). Acknowledging the dynamic relationship between groundwater and surface water when determining volume extraction limits could help inform management strategies that protect ecosystems and sustain groundwater and surface water systems. Science-based allocation limits with a two-way process between scientists and managers could help ensure that sufficient system knowledge is available to support and inform management activities.

With the potential long lag time between groundwater systems and discharges to inland and coastal waters, developing a more holistic approach that acknowledges groundwater as a resource and a conduit for transporting water and contaminants into terrestrial and marine environments could inform management decisions. The recently proposed 'Source-to-Sea' approach combines elements of the holistic management approaches of Integrated Water Resources Management and Integrated Coastal Management (Granit et al., 2017). These management approaches consider the interconnectivity of processes in hydrological and/or coastal systems. However, Source-to-Sea management approaches do not explicitly consider groundwater flow paths and the contaminants that they could transport (Granit et al., 2017; Michels-Brito et al., 2023). A new approach that views terrestrial, freshwater, and marine systems as a continuum (Fig. 1) rather than discrete segments could result in a more effective management framework.

Water-resources management is a complex process incorporating stakeholder views, and potential ecological disturbances and water extraction to support agriculture, industry, and town water supplies. There are multiple management approaches that provide opportunities to achieve this goal. For example, adaptive management, an approach that emphasises structured decision-making and uncertainty reduction (Williams et al., 2009), is increasingly cited in groundwater planning (Rohde et al., 2017). This approach involves the iterative integration of new knowledge through targeted monitoring to achieve desired outcomes (Thomann et al., 2022). Continuous reassessment and refinement of management actions via targeted monitoring improve the effectiveness of management actions over time (Thomann et al., 2020). Participatory management approaches (Ocampo-Melgar et al., 2022) can help involve stakeholders in solving complex problems. With many aggravated conflicts over water (Gleick and Shimabuku, 2023), shifting from top-down to participatory decision-making can help address the common issue of water-resources management decisions often being disconnected from their social context (Huggins et al., 2023).

6. Concluding remarks

Interactions between groundwater and surface water occur in a range of settings from headwater streams to coastlines and play a critical role in controlling many physical and biogeochemical processes. Given the broad range of processes that influence and are influenced by GW-SW interactions, the growth in research interest in GW-SW interactions in terms of both the number of research outputs and contributing disciplines is not at all surprising. This multidisciplinary in research should continue to inform our understanding of important parameters and processes that control GW-SW interactions. While there has been a notable volume of research into the complex interactions between surface water and groundwater, there remains much to do to integrate the learnings from the many disciplines that investigate these processes. The papers that we identified as having potential to have future impact on GW-SW interactions research (Table 3) focused on understanding processes using field measurements and/or numerical modelling. Much of the past research focus has been at relatively small scale, i.e., bedform to reach scale, with impediments in upscaling those findings in ways that could benefit resource managers of aquatic systems. Growing populations, continued economic development, and effects of climate change demonstrate the utility of approaches that upscale available local-scale knowledge. Effective management, mitigation, and adaptation strategies could help address these challenges to

provide water for communities and the environment. Thus, studying and understanding relevant processes in GW-SW systems could help generate knowledge that could be used to inform this conversation. The emergence of new mapping and monitoring technologies, tracer techniques and modelling capacity provide new opportunities to inform strategies to address the societal and technical challenge of monitoring, predicting, and managing GW-SW exchanges.

CRediT authorship contribution statement

Dylan J. Irvine: Writing – review & editing, Writing – original draft, Visualization, Software, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **Kamini Singha:** Writing – review & editing, Writing – original draft. **Barret L. Kurylyk:** Writing – review & editing, Writing – original draft. **Martin A. Briggs:** Writing – review & editing, Writing – original draft. **Yakub Sebastian:** Writing – review & editing, Writing – original draft, Visualization, Formal analysis. **Douglas R. Tait:** Writing – review & editing, Writing – original draft. **Ashley M. Helton:** Writing – review & editing, Writing – original draft.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Dylan Irvine reports financial support was provided by CRC For Developing Northern Australia. Martin Briggs reports financial support was provided by US Geological Survey. Dylan Irvine is an Associate Editor at the Journal of Hydrology. If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data available via FigShare

Acknowledgments

Generative AI was not used to produce text. During the preparation of this work the authors used ChatGPT for rephrasing and to facilitate elements of the data analysis. The authors reviewed and edited the content as needed and take full responsibility for the content of the publication. D.J. Irvine was supported by the CRC for Developing Northern Australia through the Water Security Program (AT.7.2223014). K. Singha was partially supported on this project by the ORNL Watershed Dynamics and Evolution (WaDE) Science Focus Area sponsored by the DOE Office of Science Office of Biological and Environmental Research Program. M.A. Briggs was supported on this project by the U.S. Geological Survey Next Generation Water Observing System Program. Additional resources and investigations are available in the [Supporting Information](#). Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. The authors would like to thank the anonymous peer reviewers and internal reviewer Julia Cantelon for their detailed reviews that helped to improve the work. The Scopus data and Python scripts to perform most of the analyses are available at Figshare, DOI: 10.6084/m9.figshare.26459395.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhydrol.2024.132061>.

References

- Abramo, G., D'Angelo, C.A., Mele, I., 2022. Impact of Covid-19 on research output by gender across countries. *Scientometrics* 127, 6811–6826. <https://doi.org/10.1007/s11192-021-04245-x>.
- Acharya, B.S., Bhandari, M., Bandini, F., Pizarro, A., Perks, M., Joshi, D.R., Wang, S., Dogwiler, T., Ray, R.L., Kharel, G., Sharma, S., 2021. Unmanned Aerial Vehicles in Hydrology and Water Management: Applications, Challenges, and Perspectives. *Water Resources Research* 57, e2021WR029925. <https://doi.org/10.1029/2021WR029925>.
- Acworth, R.L., Dasey, G.R., 2003. Mapping of the hyporheic zone around a tidal creek using a combination of borehole logging, borehole electrical tomography and cross-creek electrical imaging, New South Wales, Australia. *Hydrol. J.* 11, 368–377. <https://doi.org/10.1007/s10040-003-0258-4>.
- Alfaro, C., Wallace, M., 1994. Origin and classification of springs and historical review with current applications. *Geo* 24, 112–124. <https://doi.org/10.1007/BF00767884>.
- Ali, R., McFarlane, D., Varma, S., Dawes, W., Emelyanova, I., Hodgson, G., Charles, S., 2012. Potential climate change impacts on groundwater resources of south-western Australia. *J. Hydrol.* 475, 456–472. <https://doi.org/10.1016/j.jhydrol.2012.04.043>.
- Altenau, E.H., Pavelsky, T.M., Durand, M.T., Yang, X., Frasson, R.P. de M., Bendezu, L., 2021. The Surface Water and Ocean Topography (SWOT) Mission River Database (SWORD): A Global River Network for Satellite Data Products. *Water Resources Research* 57, e2021WR030054. <https://doi.org/10.1029/2021WR030054>.
- Anderson, M.P., 2005. Heat as a Ground Water Tracer. *Groundwater* 43, 951–968. <https://doi.org/10.1111/j.1745-6584.2005.00052.x>.
- Aquanty, 2015. HydroGeoSphere Reference Manual.
- Aquanty, 2024. Canada 1 Water [WWW Document]. Canada 1 Water. URL <https://www.canada1water.ca> (accessed 3.25.24).
- Arnold, J.G., Allen, P.M., 1999. Automated Methods for Estimating Baseflow and Ground Water Recharge from Streamflow Records. *JAWRA Journal of the American Water Resources Association* 35, 411–424. <https://doi.org/10.1111/j.1752-1688.1999.tb03599.x>.
- Ba, M.M., Heyman, J., Rivière, A., Soulayrol, M.O., Stubbe, V., Meric, F., Kergosien, B., Rolland, P., Petton, C., Lavenant, N., Kermarrec, J.J., Crave, A., 2023. A dataset on physico-chemical hyporheic variables in the Selune River: Towards understanding the impact of dam removal on riverbed clogging processes. *Data Brief* 46, 108837. <https://doi.org/10.1016/j.dib.2022.108837>.
- Bailey, R.T., Park, S., Bieger, K., Arnold, J.G., Allen, P.M., 2020. Enhancing SWAT+ simulation of groundwater flow and groundwater-surface water interactions using MODFLOW routines. *Environ. Model. Softw.* 126, 104660. <https://doi.org/10.1016/j.envsoft.2020.104660>.
- Balestra, V., Bellopede, R., 2022. Microplastic pollution in show cave sediments: First evidence and detection technique. *Environ. Pollut.* 292, 118261. <https://doi.org/10.1016/j.envpol.2021.118261>.
- Ball, L.B., Bloss, B.R., Bedrosian, P.A., Grauch, V.J.S., Smith, B.D., 2015. Airborne Electromagnetic and Magnetic Survey Data of the Paradox and San Luis Valleys, Colorado (No. 1015–1024). U.S. Geological Survey.
- Bandini, F., Jakobsen, J., Olesen, D., Reyna-Gutierrez, J.A., Bauer-Gottwein, P., 2017. Measuring water level in rivers and lakes from lightweight Unmanned Aerial Vehicles. *J. Hydrol.* 548, 237–250. <https://doi.org/10.1016/j.jhydrol.2017.02.038>.
- Banks, E.W., Shanfield, M.A., Noorduijn, S., McCallum, J., Lewandowski, J., Batelaan, O., 2018. Active heat pulse sensing of 3-D-flow fields in streambeds. *Hydrol. Earth Syst. Sci.* 22, 1917–1929. <https://doi.org/10.5194/hess-22-1917-2018>.
- Barclay, J.R., Briggs, M.A., Moore, E.M., Starn, J.J., Hanson, A.E.H., Helton, A.M., 2022. Where groundwater seeps: Evaluating modeled groundwater discharge patterns with thermal infrared surveys at the river-network scale. *Adv. Water Resour.* 160, 104108. <https://doi.org/10.1016/j.advwatres.2021.104108>.
- Barclay, J.R., Starn, J.J., Briggs, M.A., Helton, A.M., 2020. Improved Prediction of Management-Relevant Groundwater Discharge Characteristics Throughout River Networks. *Water Resources Research* 56, e2020WR028027. <https://doi.org/10.1029/2020WR028027>.
- Barclay, J.R., Topp, S.N., Koenig, L.E., Sleekman, M.J., Appling, A.P., 2023. Train, Inform, Borrow, or Combine? Approaches to Process-Guided Deep Learning for Groundwater-Influenced Stream Temperature Prediction. *Water Resources Research* 59, e2023WR035327. <https://doi.org/10.1029/2023WR035327>.
- Barlow, P.M., Hess, K.M., 1993. Simulated hydrologic responses of the Quashnet River stream-aquifer system to proposed ground-water withdrawals, Cape Cod, Massachusetts (No. 93–4064), Water-Resources Investigations Report. U.S. Geological Survey ; Earth Science Information Center, Open-File Reports Section [distributor]. <https://doi.org/10.3133/wri934064>.
- Basu, B., Morrissey, P., Gill, L.W., 2022a. Application of Nonlinear Time Series and Machine Learning Algorithms for Forecasting Groundwater Flooding in a Lowland Karst Area. *Water Resources Research* 58, e2021WR029576. <https://doi.org/10.1029/2021WR029576>.
- Basu, N.B., Van Meter, K.J., Byrnes, D.K., Van Cappellen, P., Brouwer, R., Jakobsen, B.H., Jarsjö, J., Rudolph, D.L., Cunha, M.C., Nelson, N., Bhattacharya, R., Destouni, G., Olsen, S.B., 2022b. Managing nitrogen legacies to accelerate water quality improvement. *Nat. Geosci.* 15, 97–105. <https://doi.org/10.1038/s41561-021-00889-9>.
- Befus, K.M., Cardenas, M.B., Erler, D.V., Santos, I.R., Eyre, B.D., 2013. Heat transport dynamics at a sandy intertidal zone. *Water Resour. Res.* 49, 3770–3786. <https://doi.org/10.1002/wrcr.20325>.
- Benz, S.A., Irvine, D.J., Rau, G.C., Bayer, P., Menberg, K., Blum, P., Jamieson, R.C., Griebler, C., Kurylyk, B.L., 2024. Global groundwater warming due to climate change. *Nat. Geosci.* 17, 545–551. <https://doi.org/10.1038/s41561-024-01453-x>.

- Beven, K.J., 2018. On hypothesis testing in hydrology: Why falsification of models is still a really good idea. *Wires Water* 5, e1278.
- Binley, A., Hubbard, S.S., Huisman, J.A., Revil, A., Robinson, D.A., Singha, K., Slater, L. D., 2015. The emergence of hydrogeophysics for improved understanding of subsurface processes over multiple scales. *Water Resour. Res.* 51, 3837–3866. <https://doi.org/10.1002/2015WR017016>.
- Blanc, T., Peel, M., Brennwald, M.S., Kipfer, R., Brunner, P., 2024. Efficient injection of gas tracers into rivers: A tool to study Surface water–Groundwater interactions. *Water Res.* 254, 121375. <https://doi.org/10.1016/j.watres.2024.121375>.
- Blaurock, K., Garthen, P., da Silva, M.P., Beudert, B., Gilfedder, B.S., Fleckenstein, J.H., Peiffer, S., Lechtenfeld, O.J., Hopp, L., 2022. Riparian Microtopography Affects Event-Driven Stream DOC Concentrations and DOM Quality in a Forested Headwater Catchment. *Journal of Geophysical Research: Biogeosciences* 127, e2022JG006831. <https://doi.org/10.1029/2022JG006831>.
- Blondel, V.D., Guillaume, J.-L., Lambiotte, R., Lefebvre, E., 2008. Fast unfolding of communities in large networks. *J. Stat. Mech.* 2008, P10008. <https://doi.org/10.1088/1742-5468/2008/10/P10008>.
- Blöschl, G., Ardoin-Bardin, S., Bonell, M., Dörninger, M., Goodrich, D., Gutknecht, D., Matamoros, D., Merz, B., Shand, P., Szolgay, J., 2007. At what scales do climate variability and land cover change impact on flooding and low flows? *Hydrol. Process.* 21, 1241–1247. <https://doi.org/10.1002/hyp.6669>.
- Boano, F., Camporeale, C., Revelli, R., Ridolfi, L., 2006. Sinuosity-driven hyporheic exchange in meandering rivers. *Geophys. Res. Lett.* 33. <https://doi.org/10.1029/2006GL027630>.
- Boano, F., Harvey, J.W., Marion, A., Packman, A.I., Revelli, R., Ridolfi, L., Wörman, A., 2014. Hyporheic flow and transport processes: Mechanisms, models, and biogeochemical implications. *Rev. Geophys.* 52, 603–679. <https://doi.org/10.1002/2012RG000417>.
- Boulton, A.J., Findlay, S., Marmonier, P., Stanley, E.H., Valett, H.M., 1998. The Functional Significance of the Hyporheic Zone in Streams and Rivers. *Annu. Rev. Ecol. Syst.* 29, 59–81. <https://doi.org/10.1146/annurev.ecolsys.29.1.59>.
- Boulton, A.J., Dattr, T., Kasahara, T., Mutz, M., Stanford, J.A., 2010. Ecology and management of the hyporheic zone: stream–groundwater interactions of running waters and their floodplains. *J. n. Am. Benthol. Soc.* 29, 26–40. <https://doi.org/10.1899/08-017.1>.
- Brennwald, M.S., Schmidt, M., Oser, J., Kipfer, R., 2016. A Portable and Autonomous Mass Spectrometric System for On-Site Environmental Gas Analysis. *Environ. Sci. Technol.* 50, 13455–13463. <https://doi.org/10.1021/acs.est.6b03669>.
- Brennwald, M.S., Peel, M., Blanc, T., Tomonaga, Y., Kipfer, R., Brunner, P., Hunkeler, D., 2022. New Experimental Tools to Use Noble Gases as Artificial Tracers for Groundwater Flow. *Front. Water* 4. <https://doi.org/10.3389/frwa.2022.925294>.
- Briggs, M.A., Goodling, P., Johnson, Z.C., Rogers, K.M., Hitt, N.P., Fair, J.B., Snyder, C. D., 2022. Bedrock depth influences spatial patterns of summer baseflow, temperature and flow disconnection for mountainous headwater streams. *Hydrol. Earth Syst. Sci.* 26, 3989–4011. <https://doi.org/10.5194/hess-26-3989-2022>.
- Briggs, M., Hare, D., 2018. Explicit consideration of preferential groundwater discharges as surface water ecosystem control points. *Hydrol. Process.* 32. <https://doi.org/10.1002/hyp.13178>.
- Briggs, M.A., Johnson, Z.C., Snyder, C.D., Hitt, N.P., Kurylyk, B.L., Lautz, L., Irvine, D.J., Hurley, S.T., Lane, J.W., 2018. Inferring watershed hydraulics and cold-water habitat persistence using multi-year air and stream temperature signals. *Sci. Total Environ.* 636, 1117–1127. <https://doi.org/10.1016/j.scitotenv.2018.04.344>.
- Briggs, M.A., Tokranov, A.K., Hull, R.B., LeBlanc, D.R., Haynes, A.B., Lane, J.W., 2020. Hillslope groundwater discharges provide localized stream ecosystem buffers from regional per- and polyfluoroalkyl substances contamination. *Hydrol. Process.* 34, 2281–2291. <https://doi.org/10.1002/hyp.13752>.
- Brunke, M., Gonser, T., 1997. The ecological significance of exchange processes between rivers and groundwater. *Freshw. Biol.* 37, 1–33. <https://doi.org/10.1046/j.1365-2427.1997.00143.x>.
- Brunner, P., Cook, P.G., Simmons, C.T., 2009. Hydrogeologic controls on disconnection between surface water and groundwater. *Water Resour. Res.* 45. <https://doi.org/10.1029/2008WR006953>.
- Brunner, P., Simmons, C.T., 2012. HydroGeoSphere: a fully integrated, physically based hydrological model. *Groundwater* 50, 170–176. <https://doi.org/10.1111/j.1745-6584.2011.00882.x>.
- Brunner, P., Therrien, R., Renard, P., Simmons, C.T., Franssen, H.-J.H., 2017. Advances in understanding river–groundwater interactions: River–Groundwater Interactions. *Rev. Geophys.* 55, 818–854. <https://doi.org/10.1002/2017RG000556>.
- Brunner, G.W., 2020. HEC-RAS Hydraulic Reference Manual.
- Brusseau, M.L., Anderson, R.H., Guo, B., 2020. PFAS concentrations in soils: Background levels versus contaminated sites. *Sci. Total Environ.* 740, 140017. <https://doi.org/10.1016/j.scitotenv.2020.140017>.
- Burkholder, B.K., Grant, G.E., Haggerty, R., Khangaonkar, T., Wampler, P.J., 2008. Influence of hyporheic flow and geomorphology on temperature of a large, gravel-bed river, Clackamas River, Oregon, USA. *Hydrol. Process.* 22, 941–953. <https://doi.org/10.1002/hyp.6984>.
- Burnett, W.C., Bokuniewicz, H., Huettel, M., Moore, W.S., Taniguchi, M., 2003. Groundwater and pore water inputs to the coastal zone. *Biogeochemistry* 66, 3–33. <https://doi.org/10.1023/B:BIOG.0000006066.21240.53>.
- Calderwood, A.J., Pauloo, R.A., Yoder, A.M., Fogg, G.E., 2020. Low-Cost, Open Source Wireless Sensor Network for Real-Time Scalable Groundwater Monitoring. *Water* 12, 1066. <https://doi.org/10.3390/w12041066>.
- Cantelon, J.A., Guimond, J.A., Robinson, C.E., Michael, H.A., Kurylyk, B.L., 2022. Vertical Saltwater Intrusion in Coastal Aquifers Driven by Episodic Flooding: A Review. *Water Resources Research* 58, e2022WR032614. <https://doi.org/10.1029/2022WR032614>.
- Cardenas, M.B., 2008. Surface water–groundwater interface geomorphology leads to scaling of residence times. *Geophys. Res. Lett.* 35. <https://doi.org/10.1029/2008GL033753>.
- Cardenas, M.B., 2009. Stream–aquifer interactions and hyporheic exchange in gaining and losing sinuous streams. *Water Resour. Res.* 45. <https://doi.org/10.1029/2008WR007651>.
- Cardenas, M.B., Markowski, M.S., 2011. Geoelectrical Imaging of Hyporheic Exchange and Mixing of River Water and Groundwater in a Large Regulated River. *Environ. Sci. Technol.* 45, 1407–1411. <https://doi.org/10.1021/es103438a>.
- Cardenas, M., Wilson, J.L., Haggerty, R., 2008. Residence time of bedform-driven hyporheic exchange. *Adv. Water Resour.* 31, 1382–1386. <https://doi.org/10.1016/j.advwatres.2008.07.006>.
- Cardenas, M.B., Zlotnik, V.A., 2003a. Three-dimensional model of modern channel bend deposits. *Water Resour. Res.* 39. <https://doi.org/10.1029/2002WR001383>.
- Cardenas, M.B., Zlotnik, V.A., 2003b. A simple constant-head injection test for streambed hydraulic conductivity estimation. *Groundwater* 41, 867–871. <https://doi.org/10.1111/j.1745-6584.2003.tb02428.x>.
- Cartwright, I., Irvine, D., 2020. The spatial extent and timescales of bank infiltration and return flows in an upland river system: Implications for water quality and volumes. *Sci. Total Environ.* 743, 140748. <https://doi.org/10.1016/j.scitotenv.2020.140748>.
- Chadwick, K.D., Brodrick, P.G., Grant, K., Goulden, T., Henderson, A., Falco, N., Wainwright, H., Williams, K.H., Bill, M., Breckheimer, I., Brodie, E.L., Steltzer, H., Williams, C.F.R., Blonder, B., Chen, J., Dafflon, B., Damerow, J., Hancher, M., Khurram, A., Lamb, J., Lawrence, C.R., McCormick, M., Musinsky, J., Pierce, S., Polussa, A., Hastings Porro, M., Scott, A., Singh, H.W., Sorensen, P.O., Varadharajan, C., Whitney, B., Maher, K., 2020. Integrating airborne remote sensing and field campaigns for ecology and Earth system science. *Methods Ecol. Evol.* 11, 1492–1508. <https://doi.org/10.1111/2041-210X.13463>.
- Chawanda, C.J., George, C., Thiery, W., van Griensven, A., Tech, J., Arnold, J., Srinivasan, R., 2020. User-friendly workflows for catchment modelling: Towards reproducible SWAT+ model studies. *Environ. Model. Softw.* 134, 104812. <https://doi.org/10.1016/j.envsoft.2020.104812>.
- Chen, X., 2000. Measurement of streambed hydraulic conductivity and its anisotropy. *Environ. Geol.* 39, 1317–1324. <https://doi.org/10.1007/s002540000172>.
- Chen, C., 2012. Predictive effects of structural variation on citation counts. *J. Am. Soc. Inf. Sci. Technol.* 63, 431–449. <https://doi.org/10.1002/asi.21694>.
- Chen, C., Chen, Y., Horowitz, M., Hou, H., Liu, Z., Pellegrino, D., 2009b. Towards an explanatory and computational theory of scientific discovery. *Journal of Informetrics, Science of Science: Conceptualizations and Models of Science* 3, 191–209. <https://doi.org/10.1016/j.joi.2009.03.004>.
- Chen, Z., Hartmann, A., Wägener, T., Goldscheider, N., 2018. Dynamics of water fluxes and storages in an Alpine karst catchment under current and potential future climate conditions. *Hydrol. Earth Syst. Sci.* 22, 3807–3823. <https://doi.org/10.5194/hess-22-3807-2018>.
- Chen, Y., Qiao, S., Zhang, G., Xu, Y.J., Chen, L., Wu, L., 2020. Investigating the potential use of Sentinel-1 data for monitoring wetland water level changes in China's Momoge National Nature Reserve. *PeerJ* 8, e8616.
- Chen, X., Song, J., Cheng, C., Wang, D., Lackey, S.O., 2009a. A new method for mapping variability in vertical seepage flux in streambeds. *Hydrogeol. J.* 17, 519–525. <https://doi.org/10.1007/s10040-008-0384-0>.
- Chen, C., Song, M., 2019. Visualizing a field of research: A methodology of systematic scientometric reviews. *PLoS One* 14, e0223994.
- Cheng, K.H., Luo, X., Jiao, J.J., 2020. Two-decade variations of fresh submarine groundwater discharge to Tolo Harbour and their ecological significance by coupled remote sensing and radon-222 model. *Water Res.* 178, 115866. <https://doi.org/10.1016/j.watres.2020.115866>.
- Choi, H.T., Beven, K., 2007. Multi-period and multi-criteria model conditioning to reduce prediction uncertainty in an application of TOPMODEL within the GLUE framework. *J. Hydrol.* 332, 316–336. <https://doi.org/10.1016/j.jhydrol.2006.07.012>.
- Claude, C., Cockenpot, S., Arfib, B., Meulé, S., Radakovitch, O., 2019. Accuracy and sensitivity of radium mass balances in assessing karstic submarine groundwater discharge in the stratified Calanque of Port-Miou (Mediterranean Sea). *J. Hydrol.* 578, 124034. <https://doi.org/10.1016/j.jhydrol.2019.124034>.
- Conant Jr., B., 2004. Delineating and quantifying ground water discharge zones using streambed temperatures. *Groundwater* 42, 243–257. <https://doi.org/10.1111/j.1745-6584.2004.tb02671.x>.
- Conant, B., Cherry, J.A., Gillham, R.W., 2004. A PCE groundwater plume discharging to a river: influence of the streambed and near-river zone on contaminant distributions. *J. Contam. Hydrol.* 73, 249–279. <https://doi.org/10.1016/j.jconhyd.2004.04.001>.
- Constantz, J., 2008. Heat as a tracer to determine streambed water exchanges. *Water Resour. Res.* 44. <https://doi.org/10.1029/2008WR006996>.
- Constantz, J., 2016. Streambeds merit recognition as a scientific discipline. *Wires Water* 3, 13–18. <https://doi.org/10.1002/wat2.1119>.
- Constantz, J., Thomas, C.L., Zellweger, G., 1994. Influence of diurnal variations in stream temperature on streamflow loss and groundwater recharge. *Water Resour. Res.* 30, 3253–3264. <https://doi.org/10.1029/94WR01968>.
- Cook, P.G., 2013. Estimating groundwater discharge to rivers from river chemistry surveys: GROUNDWATER DISCHARGE TO RIVERS. *Hydrol. Process.* 27, 3694–3707. <https://doi.org/10.1002/hyp.9493>.
- Cosgrove, B., Gochis, D., Flowers, T., Dugger, A., Ogden, F., Graziano, T., Clark, E., Cabell, R., Casiday, N., Cui, Z., Eicher, K., Fall, G., Feng, X., Fitzgerald, K., Frazier, N., George, C., Gibbs, R., Hernandez, L., Johnson, D., Jones, R., Karsten, L., Kefeleg, H., Kitzmiller, D., Lee, H., Liu, Y., Mashriqui, H., Mattern, D., McCluskey, A., McCreight, J.L., McDaniel, R., Midekisa, A., Newman, A., Pan, L., Pham, C., RafieeNasab, A., Rasmussen, R., Read, L., Rezaeianzadeh, M., Salas, F., Sang, D., Sampson, K., Schneider, T., Shi, Q., Sood, G., Wood, A., Wu, W., Yates, D.,

- Yu, W., Zhang, Y., 2024. NOAA's National Water Model: Advancing operational hydrology through continental-scale modeling. *JAWRA Journal of the American Water Resources Association* N/a. <https://doi.org/10.1111/1752-1688.13184>.
- Covino, T., 2017. Hydrologic connectivity as a framework for understanding biogeochemical flux through watersheds and along fluvial networks. *Geomorphology, Connectivity in Geomorphology from Binghamton 2016* (277), 133–144. <https://doi.org/10.1016/j.geomorph.2016.09.030>.
- Cranswick, R.H., Cook, P.G., Lamontagne, S., 2014. Hyporheic zone exchange fluxes and residence times inferred from riverbed temperature and radon data. *J. Hydrol.* 519, 1870–1881. <https://doi.org/10.1016/j.jhydrol.2014.09.059>.
- Crook, N., Binley, A., Knight, R., Robinson, D.A., Zarnetske, J., Haggerty, R., 2008. Electrical resistivity imaging of the architecture of streambed sediments. *Water Resour. Res.* 44. <https://doi.org/10.1029/2008WR006968>.
- CUASHI, 2024. Find, Analyze and Share Water Data | CUAHSI HydroShare [WWW Document]. HydroShare. URL <https://www.hydroshare.org/landingPage/> (accessed 8.6.24).
- Currell, M.J., Werner, A.D., McGrath, C., Webb, J.A., Berkman, M., 2017. Problems with the application of hydrogeological science to regulation of Australian mining projects: Carmichael Mine and Doongmabulla Springs. *J. Hydrol.* 548, 674–682. <https://doi.org/10.1016/j.jhydrol.2017.03.031>.
- Currell, M.J., Irvine, D.J., Werner, A.D., McGrath, C., 2020. Science sidelined in approval of Australia's largest coal mine. *Nat Sustain* 3, 644–649. <https://doi.org/10.1038/s41893-020-0527-4>.
- Dai, C., Chen, Q., Wan, T., Liu, F., Gong, Y., Wang, Q., 2021. Literary runaway: Increasingly more references cited per academic research article from 1980 to 2019. *PLoS One* 16, e0255849.
- Davis, J., Pavlova, A., Thompson, R., Sunnucks, P., 2013. Evolutionary refugia and ecological refuges: key concepts for conserving Australian arid zone freshwater biodiversity under climate change. *Glob. Chang. Biol.* 19, 1970–1984. <https://doi.org/10.1111/gcb.12203>.
- Dean, S., Freer, J., Beven, K., Wade, A.J., Butterfield, D., 2009. Uncertainty assessment of a process-based integrated catchment model of phosphorus. *Stoch Environ Res Risk Assess* 23, 991–1010. <https://doi.org/10.1007/s00477-008-0273-z>.
- Deerwester, S., Dumais, S.T., Furnas, G.W., Landauer, T.K., Harshman, R., 1990. Indexing by latent semantic analysis. *J. Am. Soc. Inf. Sci.* 41, 391–407. [https://doi.org/10.1002/\(SICI\)1097-4571\(199009\)41:6<391::AID-ASII>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1097-4571(199009)41:6<391::AID-ASII>3.0.CO;2-9).
- DelVecchia, A.G., Shanfield, M., Zimmer, M.A., Busch, M.H., Krabbenhoft, C.A., Stubbington, R., Kaiser, K.E., Burrows, R.M., Hosen, J., Detry, T., Kampf, S.K., Zipper, S.C., Fritz, K., Costigan, K., Allen, D.C., 2022. Reconceptualizing the hyporheic zone for nonperennial rivers and streams. *Freshwater Science* 41, 167–182. <https://doi.org/10.1086/720071>.
- DHI, 2023. MIKE SHE User Guide and Reference Manual.
- Dinis, H., Rocha, J., Matos, T., Gonçalves, L.M., Martins, M., 2022. The Challenge of Long-Distance Over-the-Air Wireless Links in the Ocean: A Survey on Water-to-Water and Water-to-Land M2M Communication. *Appl. Sci.* 12, 6439. <https://doi.org/10.3390/app12136439>.
- Divine, C., Wadhawan, A., Pulikkal, V., Khambhammettu, P., Erickson, J., 2023. Polyfluoroalkyl substances requiring a renewed focus on groundwater-surface water interactions. *Groundwater Monitoring & Remediation* 43, 14–31. <https://doi.org/10.1111/gwmr.12569>.
- Doble, R., Brunner, P., McCallum, J., Cook, P.G., 2012. An Analysis of River Bank Slope and Unsaturated Flow Effects on Bank Storage. *Groundwater* 50, 77–86. <https://doi.org/10.1111/j.1745-6584.2011.00821.x>.
- Doetsch, J., Linde, N., Pessognelli, M., Green, A.G., Günther, T., 2012. Constraining 3-D electrical resistance tomography with GPR reflection data for improved aquifer characterization. *Journal of Applied Geophysics, Developments in GPR and near-Surface Seismics - New Applications and Strategies for Data Integration, Inversion, and Modelling* 78, 68–76. <https://doi.org/10.1016/j.jappgeo.2011.04.008>.
- Doody, T.M., Bengel, S.N., Pritchard, J.L., Overton, I.C., 2014. Ecological response of *Eucalyptus camaldulensis* (river red gum) to extended drought and flooding along the River Murray, South Australia (1997–2011) and implications for environmental flow management. *Mar. Freshwater Res.* 65, 1082–1093. <https://doi.org/10.1071/MF13247>.
- Doody, T.M., Colloff, M.J., Davies, M., Koul, V., Benyon, R.G., Nagler, P.L., 2015. Quantifying water requirements of riparian river red gum (*Eucalyptus camaldulensis*) in the Murray-Darling Basin, Australia – implications for the management of environmental flows. *Ecology* 8, 1471–1487. <https://doi.org/10.1002/eco.1598>.
- Drage, J., Kennedy, G., 2020. Building a low-cost, internet-of-things, real-time groundwater level monitoring network. *Groundwater Monitoring & Remediation* 40, 67–73. <https://doi.org/10.1111/gwmr.12408>.
- Dugdale, S.J., Bergeron, N.E., St-Hilaire, A., 2013. Temporal variability of thermal refuges and water temperature patterns in an Atlantic salmon river. *Remote Sens. Environ.* 136, 358–373. <https://doi.org/10.1016/j.rse.2013.05.018>.
- Dugdale, S.J., Malcolm, I.A., Hannah, D.M., 2019. Drone-based Structure-from-Motion provides accurate forest canopy data to assess shading effects in river temperature models. *Sci. Total Environ.* 678, 326–340. <https://doi.org/10.1016/j.scitotenv.2019.04.229>.
- Duque, C., Knee, K.L., Russoniello, C.J., Sherif, M., Abu Risha, U.A., Sturchio, N.C., Michael, H.A., 2019. Submarine groundwater discharge data at meter scale (223Ra, 224Ra, 226Ra, 228Ra and 228Rn) in Indian River Bay (Delaware, US). *Data Brief* 27, 104728. <https://doi.org/10.1016/j.dib.2019.104728>.
- Duque, C., Russoniello, C.J., Rosenberry, D.O., 2020. History and evolution of seepage meters for quantifying flow between groundwater and surface water: Part 2 – Marine settings and submarine groundwater discharge. *Earth Sci. Rev.* 204, 103168. <https://doi.org/10.1016/j.earscirev.2020.103168>.
- Duvert, C., Priadi, C.R., Rose, A.M., Abdillah, A., Marthanty, D.R., Gibb, K.S., Kaestli, M., 2019. Sources and drivers of contamination along an urban tropical river (Ciliwung, Indonesia): Insights from microbial DNA, isotopes and water chemistry. *Sci. Total Environ.* 682, 382–393. <https://doi.org/10.1016/j.scitotenv.2019.05.189>.
- Dwivedi, D., Steefel, C.I., Arora, B., Banfield, J., Bargar, J., Boyanov, M.I., Brooks, S.C., Chen, X., Hubbard, S.S., Kaplan, D., Kemner, K.M., Nico, P.S., O'Loughlin, E.J., Pierce, E.M., Painter, S.L., Scheibe, T.D., Wainwright, H.M., Williams, K.H., Zavarin, M., 2022. From legacy contamination to watershed systems science: a review of scientific insights and technologies developed through DOE-supported research in water and energy security. *Environ. Res. Lett.* 17, 043004. <https://doi.org/10.1088/1748-9326/ac59a9>.
- Elsevier, 2024. Scopus.
- Enemark, T., Peeters, L.J.M., Mallants, D., Batelaan, O., 2019. Hydrogeological conceptual model building and testing: A review. *J. Hydrol.* 569, 310–329. <https://doi.org/10.1016/j.jhydrol.2018.12.007>.
- US EPA, 2023. To Conform with Recent Supreme Court Decision, EPA and Army Amend “Waters of the United States” Rule [WWW Document]. URL <https://www.epa.gov/newsreleases/conform-recent-supreme-court-decision-epa-and-army-amend-waters-united-states-rule> (accessed 8.6.24).
- Espinoza Ortiz, M., Apún Molina, J.P., Belmonte Jiménez, S.I., Herrera Barrientos, J., Peinado Guevara, H.J., Santamaria Miranda, A., 2023. Development of Low-Cost IoT System for Monitoring Piezometric Level and Temperature of Groundwater. *Sensors* 23, 9364. <https://doi.org/10.3390/s23239364>.
- Figshare, 2024. figshare - credit for all your research [WWW Document]. URL <https://figshare.com/> (accessed 8.6.24).
- Fleckenstein, J.H., Krause, S., Hannah, D.M., Boano, F., 2010. Groundwater-surface water interactions: New methods and models to improve understanding of processes and dynamics. *Advances in Water Resources, Special Issue on Ground Water-Surface Water Interactions* 33, 1291–1295. <https://doi.org/10.1016/j.advwatres.2010.09.011>.
- Foroumandi, E., Moradkhani, H., Sanchez-Vila, X., Singha, K., Castelletti, A., Destouni, G., 2023. ChatGPT in Hydrology and Earth Sciences: Opportunities, Prospects, and Concerns. *Water Resources Research* 59, e2023WR036288. <https://doi.org/10.1029/2023WR036288>.
- Fox, A., Boano, F., Arnon, S., 2014. Impact of losing and gaining streamflow conditions on hyporheic exchange fluxes induced by dune-shaped bed forms. *Water Resour. Res.* 50, 1895–1907. <https://doi.org/10.1002/2013WR014668>.
- Freeman, L.C., 1977. A Set of Measures of Centrality Based on Betweenness. *Sociometry* 40, 35–41. <https://doi.org/10.2307/3033543>.
- Freeze, R.A., Cherry, J.A., 1979. *Groundwater*. Prentice-Hall.
- García-Orellana, J., Rodellas, V., Tamborski, J., Diego-Feliu, M., van Beek, P., Weinstein, Y., Charette, M., Alorda-Kleinglass, A., Michael, H.A., Stieglitz, T., Scholten, J., 2021. Radium isotopes as submarine groundwater discharge (SGD) tracers: Review and recommendations. *Earth Sci. Rev.* 220, 103681. <https://doi.org/10.1016/j.earscirev.2021.103681>.
- U.S. Geological Survey, 2024a. ScienceBase Catalog Home [WWW Document]. URL <https://www.sciencebase.gov/catalog/> (accessed 8.6.24).
- U.S. Geological Survey, 2024b. Landsat 8 Scene LC08_L1TP_023030_20200305_20200310_01_T1.
- Gillfedder, B.S., Frei, S., Hofmann, H., Cartwright, I., 2015. Groundwater discharge to wetlands driven by storm and flood events: Quantification using continuous Radon-222 and electrical conductivity measurements and dynamic mass-balance modelling. *Geochim. Cosmochim. Acta* 165, 161–177. <https://doi.org/10.1016/j.gca.2015.05.037>.
- Gleason, C.J., Durand, M.T., 2020. Remote Sensing of River Discharge: A Review and a Framing for the Discipline. *Remote Sens. (basel)* 12, 1107. <https://doi.org/10.3390/rs12071107>.
- Gleick, P.H., Shimabuku, M., 2023. Water-related conflicts: definitions, data, and trends from the water conflict chronology. *Environ. Res. Lett.* 18, 034022. <https://doi.org/10.1088/1748-9326/acbb8f>.
- Gomez, M.L., Hoke, G., D'Ambrosio, S., Moreiras, S., Castro, A., 2022. Review: Hydrogeology of Northern Mendoza (Argentina), from the Andes to the eastern plains, in the context of climate change. *Hydrogeol J* 30, 725–750. <https://doi.org/10.1007/s10040-022-02462-z>.
- Gomez-Velez, J.D., Harvey, J.W., Cardenas, M.B., Kiel, B., 2015. Denitrification in the Mississippi River network controlled by flow through river bedforms. *Nature Geosci* 8, 941–945. <https://doi.org/10.1038/ngeo2567>.
- Gorelick, S.M., Zheng, C., 2015. Global change and the groundwater management challenge. *Water Resour. Res.* 51, 3031–3051. <https://doi.org/10.1002/2014WR016825>.
- Granit, J., Liss Lymer, B., Olsen, S., Tengberg, A., Nömmann, S., Clausen, T.J., 2017. A conceptual framework for governing and managing key flows in a source-to-sea continuum. *Water Policy* 19, 673–691. <https://doi.org/10.2166/wp.2017.126>.
- Guo, B., Zeng, J., Brusseau, M.L., 2020. A Mathematical Model for the Release, Transport, and Retention of Per- and Polyfluoroalkyl Substances (PFAS) in the Vadose Zone. *Water Resources Research* 56, e2019WR026667. <https://doi.org/10.1029/2019WR026667>.
- Gxokwe, S., Dube, T., Mazvimavi, D., 2020. Multispectral Remote Sensing of Wetlands in Semi-Arid and Arid Areas: A Review on Applications, Challenges and Possible Future Research Directions. *Remote Sens. (basel)* 12, 4190. <https://doi.org/10.3390/rs12244190>.
- Haggerty, R., Sun, J., Yu, H., Li, Y., 2023. Application of machine learning in groundwater quality modeling - A comprehensive review. *Water Res.* 233, 119745. <https://doi.org/10.1016/j.watres.2023.119745>.

- Hall, R., Jr, O., Baker, M.A., Arp, C.D., Koch, B.J., 2009. Hydrologic control of nitrogen removal, storage, and export in a mountain stream. *Limnol. Oceanogr.* 54, 2128–2142. <https://doi.org/10.4319/lo.2009.54.6.2128>.
- Halloran, L.J.S., Mhanna, S., Brunner, P., 2023. AI tools such as ChatGPT will disrupt hydrology, too. *Hydrol. Process.* 37, e14843.
- Hamdani, H., Eppheimer, D.E., Bogan, M.T., 2020. Release of treated effluent into streams: A global review of ecological impacts with a consideration of its potential use for environmental flows. *Freshw. Biol.* 65, 1657–1670. <https://doi.org/10.1111/fwb.13519>.
- Hammett, S., Day-Lewis, F.D., Trottier, B., Barlow, P.M., Briggs, M.A., Delin, G., Harvey, J.W., Johnson, C.D., Lane Jr, J.W., Rosenberry, D.O., Werkema, D.D., 2022. GW/SW-MST: A Groundwater/Surface-Water Method Selection Tool. *Groundwater* 60, 784–791. <https://doi.org/10.1111/gwat.13194>.
- Hannah, D.M., Brown, L.E., Milner, A.M., Gurnell, A.M., McGregor, G.R., Petts, G.E., Smith, B.P.G., Snook, D.L., 2007. Integrating climate–hydrology–ecology for alpine river systems. *Aquat. Conserv. Mar. Freshwat. Ecosyst.* 17, 636–656. <https://doi.org/10.1002/aqc.800>.
- Hare, D.K., Helton, A.M., Johnson, Z.C., Lane, J.W., Briggs, M.A., 2021. Continental-scale analysis of shallow and deep groundwater contributions to streams. *Nat Commun* 12, 1450. <https://doi.org/10.1038/s41467-021-21651-0>.
- Harvey, J.W., Bencala, K.E., 1993. The Effect of streambed topography on surface–subsurface water exchange in mountain catchments. *Water Resour. Res.* 29, 89–98. <https://doi.org/10.1029/92WR01960>.
- Harvey, J.W., Böhlke, J.K., Voytek, M.A., Scott, D., Tobias, C.R., 2013. Hyporheic zone denitrification: Controls on effective reaction depth and contribution to whole-stream mass balance. *Water Resour. Res.* 49, 6298–6316. <https://doi.org/10.1002/wrcr.20492>.
- Harvey, J., Gooseff, M., 2015. River corridor science: Hydrologic exchange and ecological consequences from bedforms to basins. *Water Resour. Res.* 51, 6893–6922. <https://doi.org/10.1002/2015WR017617>.
- Hatch, C.E., Fisher, A.T., Revenaugh, J.S., Constantz, J., Ruehl, C., 2006. Quantifying surface water–groundwater interactions using time series analysis of streambed thermal records: Method development. *Water Resour. Res.* 42. <https://doi.org/10.1029/2005WR004787>.
- Hatch, C.E., Fisher, A.T., Ruehl, C.R., Stemler, G., 2010. Spatial and temporal variations in streambed hydraulic conductivity quantified with time-series thermal methods. *J. Hydrol.* 389, 276–288. <https://doi.org/10.1016/j.jhydrol.2010.05.046>.
- Hayashi, M., Rosenberry, D.O., 2002. Effects of Ground Water Exchange on the Hydrology and Ecology of Surface Water. *Groundwater* 40, 309–316. <https://doi.org/10.1111/j.1745-6584.2002.tb02659.x>.
- Heeb, F., Singer, H., Pernet-Coudrier, B., Qi, W., Liu, H., Longrée, P., Müller, B., Berg, M., 2012. Organic Micropollutants in Rivers Downstream of the Megacity Beijing: Sources and Mass Fluxes in a Large-Scale Wastewater Irrigation System. *Environ. Sci. Technol.* 46, 8680–8688. <https://doi.org/10.1021/es301912q>.
- Heintzman, L.J., Starr, S.M., Mulligan, K.R., Barbato, L.S., McIntyre, N.E., 2017. Using Satellite Imagery to Examine the Relationship between Surface-Water Dynamics of the Salt Lakes of Western Texas and Ogallala Aquifer Depletion. *Wetlands* 37, 1055–1065. <https://doi.org/10.1007/s13157-017-0940-2>.
- Helton, A.M., Poole, G.C., Payn, R.A., Izurieta, C., Stanford, J.A., 2014. Relative influences of the river channel, floodplain surface, and alluvial aquifer on simulated hydrologic residence time in a montane river floodplain. *Geomorphology, Discontinuities in Fluvial Systems* 205, 17–26. <https://doi.org/10.1016/j.geomorph.2012.01.004>.
- Hendlin, Y.H., 2021. Surveying the Chemical Anthropocene. <https://doi.org/10.3167/ares.2021.120111>.
- Hensley, R.T., Spangler, M.J., DeVito, L.F., Decker, P.H., Cohen, M.J., Gooseff, M.N., 2020. Evaluating spatiotemporal variation in water chemistry of the upper Colorado River using longitudinal profiling. *Hydrol. Process.* 34, 1782–1793. <https://doi.org/10.1002/hyp.13690>.
- Herzog, S.P., Ward, A.S., Wondzell, S.M., 2019. Multiscale Feature-feature Interactions Control Patterns of Hyporheic Exchange in a Simulated Headwater Mountain Stream. *Water Resour. Res.* 55, 10976–10992. <https://doi.org/10.1029/2019WR025763>.
- Hester, E.T., Fox, G.A., 2020. Preferential Flow in Riparian Groundwater: Gateways for Watershed Solute Transport and Implications for Water Quality Management. *Water Resources Research* 56, e2020WR028186. <https://doi.org/10.1029/2020WR028186>.
- Hill, A.R., Lymburner, D.J., 1998. Hyporheic zone chemistry and stream–subsurface exchange in two groundwater-fed streams. *Can. J. Fish. Aquat. Sci.* 55, 495–506. <https://doi.org/10.1139/f97-250>.
- Hillebrand, O., Nödl, K., Sauter, M., Licha, T., 2015. Multitracer experiment to evaluate the attenuation of selected organic micropollutants in a karst aquifer. *Sci. Total Environ.* 506–507, 338–343. <https://doi.org/10.1016/j.scitotenv.2014.10.102>.
- Höhne, A., Müller, B.M., Schulz, H., Dara, R., Posselt, M., Lewandowski, J., McCallum, J. L., 2022. Fate of trace organic compounds in the hyporheic zone: Influence of microbial metabolism. *Water Res.* 224, 119056. <https://doi.org/10.1016/j.watres.2022.119056>.
- Howe, C.W., 2002. Policy issues and institutional impediments in the management of groundwater: lessons from case studies. *Environ. Dev. Econ.* 7, 625–641. <https://doi.org/10.1017/S1355770X02000384>.
- Huang, C.-S., Yang, T., Yeh, H.-D., 2018. Review of analytical models to stream depletion induced by pumping: Guide to model selection. *J. Hydrol.* 561, 277–285. <https://doi.org/10.1016/j.jhydrol.2018.04.015>.
- Huggins, X., Gleeson, T., Castilla-Rho, J., Holley, C., Re, V., Famiglietti, J.S., 2023. Groundwater Connections and Sustainability in Social-Ecological Systems. *Groundwater* 61, 463–478. <https://doi.org/10.1111/gwat.13305>.
- Humphreys, W.F., 2009. Hydrogeology and groundwater ecology: Does each inform the other? *Hydrogeol J* 17, 5–21. <https://doi.org/10.1007/s10040-008-0349-3>.
- Hunt, B., 2014. Review of Stream Depletion Solutions, Behavior, and Calculations. *J. Hydrol. Eng.* 19, 167–178. [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000768](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000768).
- Huntington, J.L., Niswonger, R.G., 2012. Role of surface-water and groundwater interactions on projected summertime streamflow in snow dominated regions: An integrated modeling approach. *Water Resour. Res.* 48. <https://doi.org/10.1029/2012WR012319>.
- Husic, A., Al-Aamery, N., Fox, J.F., 2022. Simulating hydrologic pathway contributions in fluvial and karst settings: An evaluation of conceptual, physically-based, and deep learning modeling approaches. *Journal of Hydrology X* 17, 100134. <https://doi.org/10.1016/j.jhydroa.2022.100134>.
- Irvine, D.J., Briggs, M.A., Lautz, L.K., Gordon, R.P., McKenzie, J.M., Cartwright, I., 2017. Using Diurnal Temperature Signals to Infer Vertical Groundwater–Surface Water Exchange. *Groundwater* 55, 10–26. <https://doi.org/10.1111/gwat.12459>.
- Irvine, D.J., Halloran, L.J.S., Brunner, P., 2023. Opportunities and limitations of the ChatGPT Advanced Data Analysis plugin for hydrological analyses. *Hydrol. Process.* 37, e15015.
- Jackson, J.L., Kuriyama, A., 2019. How Often Do Systematic Reviews Exclude Articles Not Published in English? *J Gen Intern Med* 34, 1388–1389. <https://doi.org/10.1007/s11606-019-04976-x>.
- Jägerbrand, A., Gasparovsky, D., Bouroussis, C., Schlangen, L., Lau, S., Donners, M., 2022. Correspondence: Obtrusive light, light pollution and sky glow: Areas for research, development and standardisation. *Light. Res. Technol.* 54, 191–194. <https://doi.org/10.1177/14771535211040973>.
- Jakeman, A.J., Barreteau, O., Hunt, R.J., Rinaudo, J.-D., Ross, A., Arshad, M., Hamilton, S., 2016. Integrated Groundwater Management: An Overview of Concepts and Challenges. In: Jakeman, A.J., Barreteau, O., Hunt, R.J., Rinaudo, J.-D., Ross, A. (Eds.), *Integrated Groundwater Management: Concepts, Approaches and Challenges*. Springer International Publishing, Cham, pp. 3–20. https://doi.org/10.1007/978-3-319-23576-9_1.
- Jepsen, S.M., Voss, C.I., Walvoord, M.A., Minsley, B.J., Rover, J., 2013. Linkages between lake shrinkage/expansion and sublacustrine permafrost distribution determined from remote sensing of interior Alaska, USA. *Geophys. Res. Lett.* 40, 882–887. <https://doi.org/10.1002/grl.50187>.
- Johnson, G.R., Brusseau, M.L., Carroll, K.C., Tick, G.R., Duncan, C.M., 2022. Global distributions, source-type dependencies, and concentration ranges of per- and polyfluoroalkyl substances in groundwater. *Sci. Total Environ.* 841, 156602. <https://doi.org/10.1016/j.scitotenv.2022.156602>.
- Johnson, A.G., Glenn, C.R., Burnett, W.C., Peterson, R.N., Lucey, P.G., 2008. Aerial infrared imaging reveals large nutrient-rich groundwater inputs to the ocean. *Geophys. Res. Lett.* 35. <https://doi.org/10.1029/2008GL034574>.
- Jones, J.W., 2019. Improved Automated Detection of Subpixel-Scale Inundation—Revised Dynamic Surface Water Extent (DSWE) Partial Surface Water Tests. *Remote Sens. (basel)* 11, 374. <https://doi.org/10.3390/rs11040374>.
- Jou-Claus, S., Folch, A., Garcia-Orellana, J., 2021. Applicability of Landsat 8 thermal infrared sensor for identifying submarine groundwater discharge springs in the Mediterranean Sea basin. *Hydrol. Earth Syst. Sci.* 25, 4789–4805. <https://doi.org/10.5194/hess-25-4789-2021>.
- Kahsay, K.D., Pingale, S.M., Hatiye, S.D., 2018. Impact of climate change on groundwater recharge and base flow in the sub-catchment of Tekeze basin, Ethiopia. *Groundw. Sustain. Dev.* 6, 121–133. <https://doi.org/10.1016/j.gsd.2017.12.002>.
- Kalbus, E., Reinstorf, F., Schirmer, M., 2006. Measuring methods for groundwater – surface water interactions: a review. *Hydrol. Earth Syst. Sci.* 10, 873–887. <https://doi.org/10.5194/hess-10-873-2006>.
- Kalvāns, A., Babre, A., Dēliņa, A., Popovs, K., 2020. Water stable isotope data set in temperate, lowland catchment, two years of monthly observations, River Salaca. *Latvia. Data in Brief* 30, 105607. <https://doi.org/10.1016/j.dib.2020.105607>.
- KarisAllen, J.J., Mohammed, A.A., Tamborski, J.J., Jamieson, R.C., Danieleacu, S., Kurylyk, B.L., 2022. Present and future thermal regimes of intertidal groundwater springs in a threatened coastal ecosystem. *Hydrol. Earth Syst. Sci.* 26, 4721–4740. <https://doi.org/10.5194/hess-26-4721-2022>.
- Karthe, D., Chalov, S., Borchardt, D., 2015. Water resources and their management in central Asia in the early twenty first century: status, challenges and future prospects. *Environ Earth Sci* 73, 487–499. <https://doi.org/10.1007/s12665-014-3789-1>.
- Kazmierczak, J., Nilsson, B., Postma, D., Sebok, E., Karan, S., Müller, S., Czekaj, J., Engesgaard, P., 2021. Transport of geogenic phosphorus to a groundwater-dominated eutrophic lake. *J. Hydrol.* 598, 126175. <https://doi.org/10.1016/j.jhydrol.2021.126175>.
- Keegan-Treloar, R., Irvine, D.J., Solórzano-Rivas, S.C., Werner, A.D., Banks, E.W., Currell, M.J., 2022. Fault-controlled springs: A review. *Earth Sci. Rev.* 230, 104058. <https://doi.org/10.1016/j.earscirev.2022.104058>.
- Kløve, B., Ala-aho, P., Bertrand, G., Boukalova, Z., Ertürk, A., Goldscheider, N., Ilmonen, J., Karakaya, N., Kupfersberger, H., Kværner, J., Lundberg, A., Mileusnić, M., Moszczynska, A., Muotka, T., Preda, E., Rossi, P., Sergieiev, D., Šimek, J., Wachniew, P., Angheluta, V., Widerlund, A., 2011. Groundwater dependent ecosystems. Part I: Hydroecological status and trends. *Environmental Science & Policy, Adapting to Climate Change: Reducing Water-related Risks in Europe* 14, 770–781. <https://doi.org/10.1016/j.envsci.2011.04.002>.
- Knapp, J.L.A., González-Pinzón, R., Drummond, J.D., Larsen, L.G., Cirkpa, O.A., Harvey, J.W., 2017. Tracer-based characterization of hyporheic exchange and benthic biolayers in streams. *Water Resour. Res.* 53, 1575–1594. <https://doi.org/10.1002/2016WR019393>.
- Koch, F.W., Voytek, E.B., Day-Lewis, F.D., Healy, R., Briggs, M.A., Lane Jr, J.W., Werkema, D., 2016. 1DTempPro V2: New Features for Inferring Groundwater/

- Surface-Water Exchange. *Groundwater* 54, 434–439. <https://doi.org/10.1111/gwat.12369>.
- Krause, S., Hannah, D.M., Fleckenstein, J.H., Heppell, C.M., Kaeser, D., Pickup, R., Pinay, G., Robertson, A.L., Wood, P.J., 2011. Inter-disciplinary perspectives on processes in the hyporheic zone. *Ecohydrology* 4, 481–499. <https://doi.org/10.1002/eco.176>.
- Kurylyk, B.L., MacQuarrie, K.T.B., Linnansaari, T., Cunjak, R.A., Curry, R.A., 2015. Preserving, augmenting, and creating cold-water thermal refugia in rivers: concepts derived from research on the Miramichi River, New Brunswick (Canada). *Ecohydrology* 8, 1095–1108. <https://doi.org/10.1002/eco.1566>.
- Ladson, A.R., Brown, R., Neal, B., Nathan, R., 2013. A Standard Approach to Baseflow Separation Using The Lyne and Hollick Filter. *Australasian Journal of Water Resources* 17, 25–34. <https://doi.org/10.7158/13241583.2013.11465417>.
- Lane, J.W., Briggs, M.A., Maurya, P.K., White, E.A., Pedersen, J.B., Auker, E., Terry, N., Minsley, B., Kress, W., LeBlond, D.R., Adams, R., Johnson, C.D., 2020. Characterizing the diverse hydrogeology underlying rivers and estuaries using new floating transient electromagnetic methodology. *Sci. Total Environ.* 740, 140074. <https://doi.org/10.1016/j.scitotenv.2020.140074>.
- Langevin, C.D., Hughes, J.D., Banta, E., Provost, A., Niswonger, R., Panday, S., 2017. MODFLOW 6, the U.S. Geological Survey Modular Hydrologic Model. <https://doi.org/10.5066/F76Q1VQV>.
- Lapworth, D.J., Baran, N., Stuart, M.E., Manamsa, K., Talbot, J., 2015. Persistent and emerging micro-organic contaminants in Chalk groundwater of England and France. *Environ. Pollut.* 203, 214–225. <https://doi.org/10.1016/j.envpol.2015.02.030>.
- Levy, Z.F., Fram, M.S., Faulkner, K.E., Alpers, C.N., Soltero, E.M., Taylor, K.A., 2020. Effects of montane watershed development on vulnerability of domestic groundwater supply during drought. *J. Hydrol.* 583, 124567. <https://doi.org/10.1016/j.jhydrol.2020.124567>.
- Lewandowski, J., Meinikmann, K., Nützmann, G., Rosenberry, D.O., 2015. Groundwater – the disregarded component in lake water and nutrient budgets. Part 2: effects of groundwater on nutrients. *Hydrol. Process.* 29, 2922–2955. <https://doi.org/10.1002/hyp.10384>.
- Li, Y., Tong, L., Zhang, J., Liu, H., Li, M., Wen, Z., 2023. Distribution and risk assessment of antibiotics under water level fluctuation in the riparian zone of the Hanjiang River. *Ecotoxicol. Environ. Saf.* 256, 114833. <https://doi.org/10.1016/j.ecoenv.2023.114833>.
- Lin, Y.-F., Chang, C.-H., Tsai, J.-P., 2022. Analytical solution for estimating transient vertical groundwater flux from temperature-depth profiles. *J. Hydrol.* 610, 127920. <https://doi.org/10.1016/j.jhydrol.2022.127920>.
- Luo, M., Zhang, Y., Li, H., Hu, W., Xiao, K., Yu, S., Zheng, C., Wang, X., 2022. Pollution assessment and sources of dissolved heavy metals in coastal water of a highly urbanized coastal area: The role of groundwater discharge. *Sci. Total Environ.* 807, 151070. <https://doi.org/10.1016/j.scitotenv.2021.151070>.
- Ma, R., Chen, K., Andrews, C.B., Loheide, S.P., Sawyer, A.H., Jiang, X., Briggs, M.A., Cook, P.G., Gorelick, S.M., Prommer, H., Scanlon, B.R., Guo, Z., Zheng, C., 2024. Methods for Quantifying Interactions Between Groundwater and Surface Water. <https://doi.org/10.1146/annurev-environ-111522-104534>.
- Ma, Q., Zhang, Y., 2020. Global Research Trends and Hotspots on Submarine Groundwater Discharge (SGD): A Bibliometric Analysis. *Int. J. Environ. Res. Public Health* 17, 830. <https://doi.org/10.3390/ijerph17030830>.
- Malenda, H.F., Sutfin, N.A., Guryan, G., Stauffer, S., Rowland, J.C., Williams, K.H., Singha, K., 2019. From Grain to Floodplain: Evaluating heterogeneity of floodplain hydrostratigraphy using sedimentology, geophysics, and remote sensing. *Earth Surf. Proc. Land.* 44, 1799–1815. <https://doi.org/10.1002/esp.4613>.
- Mangel, A.R., Dawson, C.B., Rey, D.M., Briggs, M.A., 2022. Drone applications in hydrogeophysics: Recent examples and a vision for the future. *Lead. Edge* 41, 540–547. <https://doi.org/10.1190/le41080540.1>.
- Markstrom, S.L., Niswonger, R.G., Regen, R.S., Prudic, D.E., Barlow, P.M., 2008. GSFLOW—Coupled Ground-Water and Surface-Water Flow Model Based on the Integration of the Precipitation-Runoff Modeling System (PRMS) and the Modular Ground-Water Flow Model (MODFLOW-2005).
- Martínez-Pérez, L., Luquot, L., Carrera, J., Marazuela, M.A., Goyetche, T., Pool, M., Palacios, A., Bellmunt, F., Lledo, J., Ferrer, N., del Val, L., Pezard, P.A., García-Orellana, J., Diego-Feliu, M., Rodellas, V., Saaltink, M.W., Vázquez-Suñé, E., Folch, A., 2022. A multidisciplinary approach to characterizing coastal alluvial aquifers to improve understanding of seawater intrusion and submarine groundwater discharge. *J. Hydrol.* 607, 127510. <https://doi.org/10.1016/j.jhydrol.2022.127510>.
- Maxwell, R.M., Condon, L.E., Kollet, S.J., 2015. A high-resolution simulation of groundwater and surface water over most of the continental US with the integrated hydrologic model ParFlow v3. *Geosci. Model Dev.* 8, 923–937. <https://doi.org/10.5194/gmd-8-923-2015>.
- Maxwell, R.M., Kollet, S.J., Condon, L.E., Smith, S.G., Woodward, C.S., Falgout, R.D., Ferguson, I.M., Engdahl, N., Hokkanen, J., Hector, B., Gilbert, J., Bearup, L., Jefferson, J., Baldwin, C., Bosl, W.J., Hornung, R., Ashby, S., Kulkarni, K.B., 2023. ParFlow Documentation Release 3.12.0 the ParFlow Developers.
- McCallum, A.M., Andersen, M.S., Rau, G.C., Acworth, R.I., 2012. A 1-D analytical method for estimating surface water-groundwater interactions and effective thermal diffusivity using temperature time series. *Water Resour. Res.* 48. <https://doi.org/10.1029/2012WR012007>.
- McCallum, A.M., Andersen, M.S., Rau, G.C., Larsen, J.R., Acworth, R.I., 2014. River-aquifer interactions in a semiarid environment investigated using point and reach measurements. *Water Resour. Res.* 50, 2815–2829. <https://doi.org/10.1002/2012WR012922>.
- McCallum, J.L., Cook, P.G., Brunner, P., Berhane, D., 2010. Solute dynamics during bank storage flows and implications for chemical base flow separation. *Water Resour. Res.* 46. <https://doi.org/10.1029/2009WR008539>.
- McLachlan, P.J., Chambers, J.E., Uhlemann, S.S., Binley, A., 2017. Geophysical characterisation of the groundwater-surface water interface. *Adv. Water Resour.* 109, 302–319. <https://doi.org/10.1016/j.advwatres.2017.09.016>.
- MDPI, 2023. Warning about a Suspicious Website Denouncing MDPI Journals [WWW Document]. Warning about a Suspicious Website Denouncing MDPI Journals. URL <https://www.mdpi.com/about/announcements/5482> (accessed 11.16.23).
- Meinikmann, K., Lewandowski, J., Nützmann, G., 2013. Lacustrine groundwater discharge: Combined determination of volumes and spatial patterns. *J. Hydrol.* 502, 202–211. <https://doi.org/10.1016/j.jhydrol.2013.08.021>.
- Meinikmann, K., Hupfer, M., Lewandowski, J., 2015. Phosphorus in groundwater discharge – A potential source for lake eutrophication. *J. Hydrol.* 524, 214–226. <https://doi.org/10.1016/j.jhydrol.2015.02.031>.
- Merill, L., Tonjes, D.J., 2014. A Review of the Hyporheic Zone, Stream Restoration, and Means to Enhance Denitrification. *Crit. Rev. Environ. Sci. Technol.* 44, 2337–2379. <https://doi.org/10.1080/10643389.2013.829769>.
- Meyers, Z.P., Frisbee, M.D., Rademacher, L.K., Stewart-Maddox, N.S., 2021. Old groundwater buffers the effects of a major drought in groundwater-dependent ecosystems of the eastern Sierra Nevada (CA). *Environ. Res. Lett.* 16, 044044. <https://doi.org/10.1088/1748-9326/abde5f>.
- Michels-Brito, A., Ferreira, J.C., Saito, C.H., 2023. Source-to-sea, integrated water resources management, and integrated coastal management approaches: integrative, complementary, or competing? *J. Coast Conserv.* 27, 66. <https://doi.org/10.1007/s11852-023-00999-z>.
- Miguez-Macho, G., Fan, Y., 2012. The role of groundwater in the Amazon water cycle: 1. Influence on seasonal streamflow, flooding and wetlands. *Journal of Geophysical Research: Atmospheres* 117. <https://doi.org/10.1029/2012JD017539>.
- Moore, W.S., 2010. The Effect of Submarine Groundwater Discharge on the Ocean. *Ann. Rev. Mar. Sci.* 2, 59–88. <https://doi.org/10.1146/annurev-marine-120308-081019>.
- Moore, E.M., Barclay, J.R., Haynes, A.B., Jackson, K.E., Bisson, A.M., Briggs, M.A., Helton, A.M., 2023. Where the past meets the present: connecting nitrogen from watersheds to streams through groundwater flowpaths. *Environ. Res. Lett.* 18, 124039. <https://doi.org/10.1088/1748-9326/ad0c86>.
- Murdoch, L.C., Kelly, S.E., 2003. Factors affecting the performance of conventional seepage meters. *Water Resour. Res.* 39. <https://doi.org/10.1029/2002WR001347>.
- Naranjo, R.C., Niswonger, R.G., Smith, D., Rosenberry, D., Chandra, S., 2019. Linkages between hydrology and seasonal variations of nutrients and periphyton in a large oligotrophic subalpine lake. *J. Hydrol.* 568, 877–890. <https://doi.org/10.1016/j.jhydrol.2018.11.033>.
- Neave, M., Luter, H., Padovan, A., Townsend, S., Schobben, X., Gibb, K., 2014. Multiple approaches to microbial source tracking in tropical northern Australia. *MicrobiologyOpen* 3, 860–874. <https://doi.org/10.1002/mbo3.209>.
- Nelson, A.R., Sawyer, A.H., Gabor, R.S., Saup, C.M., Bryant, S.R., Harris, K.D., Briggs, M.A., Williams, K.H., Wilkins, M.J., 2019. Heterogeneity in Hyporheic Flow, Pore Water Chemistry, and Microbial Community Composition in an Alpine Streambed. *J. Geophys. Res. Biogeosci.* 124, 3465–3478. <https://doi.org/10.1029/2019JG005226>.
- Newman, M.E.J., 2005. A measure of betweenness centrality based on random walks. *Soc. Networks* 27, 39–54. <https://doi.org/10.1016/j.socnet.2004.11.009>.
- Nogueira, G.E.H., Schmidt, C., Brunner, P., Graeber, D., Fleckenstein, J.H., 2021. Transit-Time and Temperature Control the Spatial Patterns of Aerobic Respiration and Denitrification in the Riparian Zone. *Water Resources Research* 57, e2021WR030117. <https://doi.org/10.1029/2021WR030117>.
- O'Sullivan, A.M., Kurylyk, B.L., 2022. Limiting External Absorptivity of UAV-Based Uncooled Thermal Infrared Sensors Increases Water Temperature Measurement Accuracy. *Remote Sens. (basel)* 14, 6356. <https://doi.org/10.3390/rs14246356>.
- O'Sullivan, A.M., Linnansaari, T., Curry, R.A., 2019. Ice cover exists: A quick method to delineate groundwater inputs in running waters for cold and temperate regions. *Hydrol. Process.* 33, 3297–3309. <https://doi.org/10.1002/hyp.13557>.
- Ocampo-Melgar, A., Barria, P., Chadwick, C., Rivas, C., 2022. Cooperation under conflict: participatory hydrological modeling for science policy dialogues for the Aculeo Lake. *Hydrol. Earth Syst. Sci.* 26, 5103–5118. <https://doi.org/10.5194/hess-26-5103-2022>.
- Padilla, F.M., Gallardo, M., Manzano-Agugliaro, F., 2018. Global trends in nitrate leaching research in the 1960–2017 period. *Sci. Total Environ.* 643, 400–413. <https://doi.org/10.1016/j.scitotenv.2018.06.215>.
- Pai, H., Malenda, H.F., Briggs, M.A., Singha, K., González-Pinzón, R., Goosseff, M.N., Tyler, S.W., Team, the A., 2017. Potential for Small Unmanned Aircraft Systems Applications for Identifying Groundwater-Surface Water Exchange in a Meandering River Reach. *Geophys. Res. Lett.* 44, 11868–11877. <https://doi.org/10.1002/2017GL075836>.
- Pan, Z., Ma, R., Sun, Z., Hu, Y., Chang, Q., Ge, M., Wang, S., Bu, J., Long, X., Pan, Y., Zhao, L., 2022. Integrated hydrogeological and hydrogeochemical dataset of an alpine catchment in the northern Qinghai-Tibet Plateau. *Earth Syst. Sci. Data* 14, 2147–2165. <https://doi.org/10.5194/essd-14-2147-2022>.
- Pang, L., Abeysekera, G., Hanning, K., Premaratne, A., Robson, B., Abraham, P., Sutton, R., Hanson, C., Hadfield, J., Heiligenthal, L., Stone, D., McBeth, K., Billington, C., 2020. Water tracking in surface water, groundwater and soils using free and alginate-chitosan encapsulated synthetic DNA tracers. *Water Res.* 184, 116192. <https://doi.org/10.1016/j.watres.2020.116192>.
- Panno, S.V., Kelly, W.R., Scott, J., Zheng, W., McNeish, R.E., Holm, N., Hoellein, T.J., Baranski, E.L., 2019. Microplastic Contamination in Karst Groundwater Systems. *Groundwater* 57, 189–196. <https://doi.org/10.1111/gwat.12862>.
- Payn, R.A., Goosseff, M.N., McGlynn, B.L., Bencala, K.E., Wondzell, S.M., 2009. Channel water balance and exchange with subsurface flow along a mountain headwater

- stream in Montana United States. *Water Resour. Res.* 45. <https://doi.org/10.1029/2008WR007644>.
- Petpongpan, C., Ekkawatpanit, C., Bailey, R.T., Kositgittiwong, D., Saraphirom, P., 2022. Evaluating Surface Water-groundwater Interactions in Consequence of Changes in Climate and Groundwater Extraction. *Water Resour Manage* 36, 5767–5783. <https://doi.org/10.1007/s11269-022-03334-7>.
- Pétre, M.-A., Genereux, D.P., Koropeckyj-Cox, L., Knappe, D.R.U., Duboscq, S., Gilmore, T.E., Hopkins, Z.R., 2021. Per- and Polyfluoroalkyl Substance (PFAS) Transport from Groundwater to Streams near a PFAS Manufacturing Facility in North Carolina. USA. *Environ. Sci. Technol.* 55, 5848–5856. <https://doi.org/10.1021/acs.est.0c07978>.
- Ping, X., Xian, Y., Jin, M., 2023. Effect of Particulate Organic Carbon Deposition on Nitrate Reduction in the Hyporheic Zone. *Water Resources Research* 59, e2022WR034253. <https://doi.org/10.1029/2022WR034253>.
- Ping, X., Wang, J., Jin, M., 2021. Experimental analysis of natural organic matter controls on nitrogen reduction during bank storage. *J. Contam. Hydrol.* 242, 103866. <https://doi.org/10.1016/j.jconhyd.2021.103866>.
- Poole, G.C., O'Daniel, S.J., Jones, K.L., Woessner, W.W., Bernhardt, E.S., Helton, A.M., Stanford, J.A., Boer, B.R., Beechie, T.J., 2008. Hydrologic spiralling: the role of multiple interactive flow paths in stream ecosystems. *River Res. Appl.* 24, 1018–1031. <https://doi.org/10.1002/rra.1099>.
- Post, V.E.A., Groen, J., Kooi, H., Person, M., Ge, S., Edmunds, W.M., 2013. Offshore fresh groundwater reserves as a global phenomenon. *Nature* 504, 71–78. <https://doi.org/10.1038/nature12858>.
- Predatory Reports, 2022. Is MDPI a predatory publisher? (*Updated) [WWW Document]. Predatory Reports. URL <https://predatoryreports.org/news/f/is-mdpi-a-predatory-publisher> (accessed 11.16.23).
- Pulido-Velazquez, M., Peña-Haro, S., García-Prats, A., Mocholi-Almudever, A.F., Henríquez-Dole, L., Macian-Sorribes, H., López-Nicolas, A., 2015. Integrated assessment of the impact of climate and land use changes on groundwater quantity and quality in the Mancha Oriental system (Spain). *Hydrol. Earth Syst. Sci.* 19, 1677–1693. <https://doi.org/10.5194/hess-19-1677-2015>.
- Python Software Foundation, 2024. *Python Language Reference. Version 3, 8*.
- Rau, G.C., Andersen, M.S., McCallum, A.M., Roshan, H., Acworth, R.I., 2014. Heat as a tracer to quantify water flow in near-surface sediments. *Earth Sci. Rev.* 129, 40–58. <https://doi.org/10.1016/j.earscirev.2013.10.015>.
- Rey, D.M., Walvoord, M., Minsley, B., Rover, J., Singha, K., 2019. Investigating lake-area dynamics across a permafrost-thaw spectrum using airborne electromagnetic surveys and remote sensing time-series data in Yukon Flats, Alaska. *Environ. Res. Lett.* 14, 025001. <https://doi.org/10.1088/1748-9326/aaf06f>.
- Rivett, M.O., Buss, S.R., Morgan, P., Smith, J.W.N., Bemment, C.D., 2008. Nitrate attenuation in groundwater: A review of biogeochemical controlling processes. *Water Res.* 42, 4215–4232. <https://doi.org/10.1016/j.watres.2008.07.020>.
- Rohde, M.M., Froend, R., Howard, J., 2017. A Global Synthesis of Managing Groundwater Dependent Ecosystems Under Sustainable Groundwater Policy. *Groundwater* 55, 293–301. <https://doi.org/10.1111/gwat.12511>.
- Rohde, M.M., Stella, J.C., Singer, M.B., Roberts, D., Caylor, K., Albano, C.M., 2024. Establishing ecological thresholds and targets for groundwater management. *Nature Water*.
- Rosenberry, D.O., Morin, R.H., 2004. Use of an Electromagnetic Seepage Meter to Investigate Temporal Variability in Lake Seepage. *Groundwater* 42, 68–77. <https://doi.org/10.1111/j.1745-6584.2004.tb02451.x>.
- Ruehl, C., Fisher, A.T., Hatch, C., Huertos, M.L., Stemler, G., Shennan, C., 2006. Differential gauging and tracer tests resolve seepage fluxes in a strongly-losing stream. *Journal of Hydrology, Hydro-Ecological Functioning of the Pang and Lambourn Catchments*, UK 330, 235–248. <https://doi.org/10.1016/j.jhydrol.2006.03.025>.
- Rumuri, R., 2020. Dataset on hydrogeochemical characteristics of groundwater and surface water in Kattumannarkoil Taluk. India. Data in Brief 32, 106058. <https://doi.org/10.1016/j.dib.2020.106058>.
- Sanford, W.E., Pope, J.P., 2013. Quantifying Groundwater's Role in Delaying Improvements to Chesapeake Bay Water Quality. *Environ. Sci. Technol.* 47, 13330–13338. <https://doi.org/10.1021/es401334k>.
- Santos, I.R., Eyre, B.D., Huettel, M., 2012. The driving forces of porewater and groundwater flow in permeable coastal sediments: A review. *Estuar. Coast. Shelf Sci.* 98, 1–15. <https://doi.org/10.1016/j.jecss.2011.10.024>.
- Santos, I.R., Chen, X., Lecher, A.L., Sawyer, A.H., Moosdorf, N., Rodellas, V., Tamborski, J., Cho, H.-M., Dimova, N., Sugimoto, R., Bonaglia, S., Li, H., Hajati, M.-C., Li, L., 2021. Submarine groundwater discharge impacts on coastal nutrient biogeochemistry. *Nat Rev Earth Environ* 2, 307–323. <https://doi.org/10.1038/s43017-021-00152-0>.
- Sass, G.Z., Creed, I.F., Riddell, J., Bayley, S.E., 2014. Regional-scale mapping of groundwater discharge zones using thermal satellite imagery. *Hydrol. Process.* 28, 5662–5673. <https://doi.org/10.1002/hyp.10068>.
- Sawyer, A.H., David, C.H., Famiglietti, J.S., 2016a. Continental patterns of submarine groundwater discharge reveal coastal vulnerabilities. *Science* 353, 705–707. <https://doi.org/10.1126/science.aag1058>.
- Sawyer, A.H., Michael, H.A., Schroth, A.W., 2016b. From soil to sea: the role of groundwater in coastal critical zone processes. *Wires Water* 3, 706–726. <https://doi.org/10.1002/wat2.1157>.
- Schilling, O.S., Irvine, D.J., Hendricks Franssen, H.-J., Brunner, P., 2017. Estimating the Spatial Extent of Unsaturated Zones in Heterogeneous River-Aquifer Systems. *Water Resour. Res.* 53, 10583–10602. <https://doi.org/10.1002/2017WR020409>.
- Schneidewind, U., van Berkel, M., Anibas, C., Vandersteen, G., Schmidt, C., Joris, I., Seuntjens, P., Batelaan, O., Zwart, H.J., 2016. LPMLE3: A novel 1-D approach to study water flow in streambeds using heat as a tracer. *Water Resour. Res.* 52, 6596–6610. <https://doi.org/10.1002/2015WR017453>.
- Schwartz, F.W., Ibaraki, M., 2001. Hydrogeological Research: Beginning of the End or End of the Beginning? *Groundwater* 39, 492–498. <https://doi.org/10.1111/j.1745-6584.2001.tb02337.x>.
- Sebastian, Y., Chen, C., 2021. The boundary-spanning mechanisms of Nobel Prize winning papers. *PLoS One* 16, e0254744.
- Sekar, S., Perumal, M., Roy, P.D., Ganapathy, M., Senapathi, V., Yong Chung, S., Elzain, H.E., Duraisamy, M., Kamaraj, J., 2022. A review on global status of fresh and saline groundwater discharge into the ocean. *Environ Monit Assess* 194, 915. <https://doi.org/10.1007/s10661-022-10566-y>.
- Shen, Z., Tian, Y., Yao, Y., Jiang, W., Dong, J., Huang, X., Wu, X., Farooq, T.H., Yan, W., 2023. Ecological restoration research progress and prospects: A bibliometric analysis. *Ecol. Ind.* 155, 110968. <https://doi.org/10.1016/j.ecolind.2023.110968>.
- Shi, W., Wang, Q., Klepikova, M., Irvine, D.J., Rashed, A., Liao, Z., 2024. BDFLUX: A new bounded domain model for quantifying vertical fluxes in streambeds using heat as a tracer. *J. Hydrol.* 633, 130981. <https://doi.org/10.1016/j.jhydrol.2024.130981>.
- Shishaye, H.A., Tait, D.R., Maher, D.T., Befus, K.M., Erler, D., Jeffrey, L., Reading, M.J., Morgenstern, U., Kaserzon, S., Mueller, J., De Verelle-Hill, W., 2021. The legacy and drivers of groundwater nutrients and pesticides in an agriculturally impacted Quaternary aquifer system. *Sci. Total Environ.* 753, 142010. <https://doi.org/10.1016/j.scitotenv.2020.142010>.
- Silva, J.A.K., Martin, W.A., McCray, J.E., 2021. Air-water interfacial adsorption coefficients for PFAS when present as a multi-component mixture. *J. Contam. Hydrol.* 236, 103731. <https://doi.org/10.1016/j.jconhyd.2020.103731>.
- Singh, T., Gupta, S., Chigna, G., Krause, S., Wohlmuth, B., 2022. Impacts of Peak-Flow Events on Hyporheic Denitrification Potential. *Water Resour. Res.* 58, e2021. <https://doi.org/10.1029/2021WR031407>.
- Singha, K., Navarre-Stichler, A., 2022. The Importance of Groundwater in Critical Zone Science. *Groundwater* 60, 27–34. <https://doi.org/10.1111/gwat.13143>.
- Singha, K., Pidlisecky, A., Day-Lewis, F.D., Gooseff, M.N., 2008. Electrical characterization of non-Fickian transport in groundwater and hyporheic systems. *Water Resour. Res.* 44. <https://doi.org/10.1029/2008WR007048>.
- Singha, K., Sullivan, P.L., Billings, S.A., Walls, L., Li, L., Jarecek, K.M., Barnard, H.R., Gasparini, N.M., Madoff, R.D., Dhital, S., Jones, C., Kastelic, E.C., Ma, L., Perilla-Castillo, P., Song, B., Zhu, T., 2024. Expanding the Spatial Reach and Human Impacts of Critical Zone Science. *Earth's Future* 12, e2023. <https://doi.org/10.1029/2023EF003971>.
- Singley, J.G., Singha, K., Gooseff, M.N., González-Pinzón, R., Covino, T.P., Ward, A.S., Dorley, J., Hinckley, E.-L.-S., 2022. Identification of hyporheic extent and functional zonation during seasonal streamflow recession by unsupervised clustering of time-lapse electrical resistivity models. *Hydrol. Process.* 36, e14713.
- Smith, J.W.N., Bonell, M., Gibert, J., McDowell, W.H., Sudicky, E.A., Turner, J.V., Harris, R.C., 2008. Groundwater-surface water interactions, nutrient fluxes and ecological response in river corridors: Translating science into effective environmental management. *Hydrol. Process.* 22, 151–157. <https://doi.org/10.1002/hyp.6902>.
- Sophocleous, M., 2002. Interactions between groundwater and surface water: the state of the science. *Hydrogeol. J.* 10, 52–67. <https://doi.org/10.1007/s10040-001-0170-8>.
- Springer, A.E., Stevens, L.E., 2009. Spheres of discharge of springs. *Hydrogeol J* 17, 83–93. <https://doi.org/10.1007/s10040-008-0341-y>.
- Steel, E.A., Beechie, T.J., Torgersen, C.E., Fullerton, A.H., 2017. Envisioning, Quantifying, and Managing Thermal Regimes on River Networks. *Bioscience* 67, 506–522. <https://doi.org/10.1093/biosci/bix047>.
- Sterte, E.J., Lidman, F., Sjöberg, Y., Ploum, S.W., Laudon, H., 2022. Groundwater travel times predict DOC in streams and riparian soils across a heterogeneous boreal landscape. *Sci. Total Environ.* 849, 157398. <https://doi.org/10.1016/j.scitotenv.2022.157398>.
- Stonedahl, S.H., Harvey, J.W., Wörman, A., Salehin, M., Packman, A.I., 2010. A multiscale model for integrating hyporheic exchange from ripples to meanders. *Water Resour. Res.* 46. <https://doi.org/10.1029/2009WR008865>.
- Straight, B.J., Castendyk, D.N., McKnight, D.M., Newman, C.P., Filiatreault, P., Pino, A., 2021. Using an unmanned aerial vehicle water sampler to gather data in a pit-lake mining environment to assess closure and monitoring. *Environ Monit Assess* 193, 572. <https://doi.org/10.1007/s10661-021-09316-3>.
- Stubbington, R., 2012. The hyporheic zone as an invertebrate refuge: a review of variability in space, time, taxa and behaviour. *Mar. Freshwater Res.* 63, 293–311. <https://doi.org/10.1071/MF11196>.
- Taft, L., Evers, M., 2016. A review of current and possible future human–water dynamics in Myanmar's river basins. *Hydrol. Earth Syst. Sci.* 20, 4913–4928. <https://doi.org/10.5194/hess-20-4913-2016>.
- Tague, C., Brandt, W.T., 2023. Critical zone science in the Western US—Too much information? *Frontiers in Water* 5.
- Tait, D.R., Erler, D.V., Santos, I.R., Cyronak, T.J., Morgenstern, U., Eyre, B.D., 2014. The influence of groundwater inputs and age on nutrient dynamics in a coral reef lagoon. *Mar. Chem.* 166, 36–47. <https://doi.org/10.1016/j.marchem.2014.08.004>.
- Tait, D.R., Santos, I.R., Lamontagne, S., Sippo, J.Z., McMahon, A., Jeffrey, L.C., Maher, D.T., 2023. Submarine Groundwater Discharge Exceeds River Inputs as a Source of Nutrients to the Great Barrier Reef. *Environ. Sci. Technol.* 57, 15627–15634. <https://doi.org/10.1021/acs.est.3c03725>.
- Tamborski, J.J., Rogers, A.D., Bokuniewicz, H.J., Cochran, J.K., Young, C.R., 2015. Identification and quantification of diffuse fresh submarine groundwater discharge via airborne thermal infrared remote sensing. *Remote Sens. Environ.* 171, 202–217. <https://doi.org/10.1016/j.rse.2015.10.010>.
- Taniguchi, M., Dulai, H., Burnett, K.M., Santos, I.R., Sugimoto, R., Stieglitz, T., Kim, G., Moosdorf, N., Burnett, W.C., 2019. Submarine Groundwater Discharge: Updates on

- Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects. *Frontiers in Environmental Science* 7.
- Tesoriero, A.J., Duff, J.H., Saad, D.A., Spahr, N.E., Wolock, D.M., 2013. Vulnerability of Streams to Legacy Nitrate Sources. *Environ. Sci. Technol.* 47, 3623–3629. <https://doi.org/10.1021/es305026x>.
- Tetzlaff, D., Smith, A., Kleine, L., Daempfling, H., Freymueller, J., Soulsby, C., 2023. Integrated ecohydrological hydrometric and stable water isotope data of a drought-sensitive mixed land use lowland catchment. *Earth Syst. Sci. Data* 15, 1543–1554. <https://doi.org/10.5194/essd-15-1543-2023>.
- Thomann, J.A., Werner, A.D., Irvine, D.J., Currell, M.J., 2020. Adaptive management in groundwater planning and development: A review of theory and applications. *J. Hydrol.* 586, 124871. <https://doi.org/10.1016/j.jhydrol.2020.124871>.
- Thomann, J.A., Werner, A.D., Irvine, D.J., 2022. Developing adaptive management guidance for groundwater planning and development. *J. Environ. Manage.* 322, 116052. <https://doi.org/10.1016/j.jenvman.2022.116052>.
- Thornton, J.M., Therrien, R., Mariéthoz, G., Linde, N., Brunner, P., 2022. Simulating Fully-Integrated Hydrological Dynamics in Complex Alpine Headwaters: Potential and Challenges. *Water Resour. Res.* 58, e2020. <https://doi.org/10.1029/2020WR029390>.
- Tokranov, A.K., LeBlanc, D.R., Pickard, H.M., Ruyle, B.J., Barber, L.B., Hull, R.B., Sunderland, E.M., Vecitis, C.D., 2021. Surface-water/groundwater boundaries affect seasonal PFAS concentrations and PFAS precursor transformations. *Environ. Sci.: Processes Impacts* 23, 1893–1905. <https://doi.org/10.1039/D1EM00329A>.
- Tonina, D., Buffington, J.M., 2009. Hyporheic Exchange in Mountain Rivers I: Mechanics and Environmental Effects. *Geogr. Compass* 3, 1063–1086. <https://doi.org/10.1111/j.1749-8198.2009.00226.x>.
- Torgersen, C.E., Faux, R.N., McIntosh, B.A., Poage, N.J., Norton, D.J., 2001. Airborne thermal remote sensing for water temperature assessment in rivers and streams. *Remote Sens. Environ.* 76, 386–398. [https://doi.org/10.1016/S0034-4257\(01\)00186-9](https://doi.org/10.1016/S0034-4257(01)00186-9).
- Tran, N.H., Gin, K.-Y.-H., Ngo, H.H., 2015. Fecal pollution source tracking toolbox for identification, evaluation and characterization of fecal contamination in receiving urban surface waters and groundwater. *Sci. Total Environ.* 538, 38–57. <https://doi.org/10.1016/j.scitotenv.2015.07.155>.
- Tripathi, M., Yadav, P.K., Chahar, B.R., Dietrich, P., 2021. A review on groundwater-surface water interaction highlighting the significance of streambed and aquifer properties on the exchanging flux. *Environ Earth Sci* 80, 604. <https://doi.org/10.1007/s12665-021-09897-9>.
- Uhl, A., Hahn, H.J., Jäger, A., Luftenstein, T., Siemensmeyer, T., Döll, P., Noack, M., Schwenk, K., Berkhoff, S., Weiler, M., Karwautz, C., Griebler, C., 2022. Making waves: Pulling the plug—Climate change effects will turn gaining into losing streams with detrimental effects on groundwater quality. *Water Res.* 220, 118649. <https://doi.org/10.1016/j.watres.2022.118649>.
- Ulrich, C., Hubbard, S.S., Florsheim, J., Rosenberry, D., Borglin, S., Trotta, M., Seymour, D., 2015. Riverbed Clogging Associated with a California Riverbank Filtration System: An Assessment of Mechanisms and Monitoring Approaches. *J. Hydrol.* 529, 1740–1753. <https://doi.org/10.1016/j.jhydrol.2015.08.012>.
- United Nations, 2022. World Population Prospects - Population Division - United Nations [WWW Document]. URL [https://population.un.org/wpp/Download/Standard/](https://population.un.org/wpp/Download/Standard/MostUsed/) (accessed 8.13.23).
- Vainu, M., Terasmaa, J., 2016. The consequences of increased groundwater abstraction for groundwater dependent closed-basin lakes in glacial terrain. *Environ. Earth Sci.* 75, 92. <https://doi.org/10.1007/s12665-015-4967-5>.
- Valet, H.M., Fisher, S.G., Grimm, N.B., Camill, P., 1994. Vertical Hydrologic Exchange and Ecological Stability of a Desert Stream Ecosystem. *Ecology* 75, 548–560. <https://doi.org/10.2307/1939557>.
- Van Meter, K.J., Van Cappellen, P., Basu, N.B., 2018. Legacy nitrogen may prevent achievement of water quality goals in the Gulf of Mexico. *Science* 360, 427–430. <https://doi.org/10.1126/science.aar4462>.
- Vaze, J., Viney, N., Stenson, M., Renzullo, L., van Dijk, A., Dutta, D., Crosbie, R., Lerat, J., Penton, D., Vleeshouwer, J., Peeters, L., Teng, J., Kim, S., Hughes, J., Dawes, W., Zhang, Y., Leighton, B., Perraud, J.-M., Joehnk, K., Daamen, C., 2013. The Australian Water Resource Assessment Modelling System (AWRA), MODSIM2013, 20th International Congress on Modelling and Simulation.
- Vélez-Nicolás, M., García-López, S., Barbero, L., Ruiz-Ortiz, V., Sánchez-Bellón, Á., 2021. Applications of Unmanned Aerial Systems (UASs) in Hydrology: A Review. *Remote Sens. (basel)* 13, 1359. <https://doi.org/10.3390/rs13071359>.
- Viola, M.R., de Mello, C.R., Chou, S.C., Yanagi, S.N., Gomes, J.L., 2015. Assessing climate change impacts on Upper Grande River Basin hydrology, Southeast Brazil. *Int. J. Climatol.* 35, 1054–1068. <https://doi.org/10.1002/joc.4038>.
- Voss, C.I., 2015. Editor's Message: University of Kansas (USA) plagiarism allegations regarding two Hydrogeology Journal articles by Dr. Marios Sophocleous. *Hydrogeol. J* 23, 213–214. <https://doi.org/10.1007/s10040-014-1215-0>.
- Walker, G.R., Crosbie, R.S., Chiew, F.H.S., Peeters, L., Evans, R., 2021. Groundwater Impacts and Management under a Drying Climate in Southern Australia. *Water* 13, 3588. <https://doi.org/10.3390/w13243588>.
- Wallace, C.D., Sawyer, A.H., Barnes, R.T., Soltanian, M.R., Gabor, R.S., Wilkins, M.J., Moore, M.T., 2020. A Model Analysis of the Tidal Engine That Drives Nitrogen Cycling in Coastal Riparian Aquifers. *Water Resources Research* 56, e2019WR025662. <https://doi.org/10.1029/2019WR025662>.
- Ward, A.S., 2016. The evolution and state of interdisciplinary hyporheic research. *Wires Water* 3, 83–103. <https://doi.org/10.1002/wat2.1120>.
- Ward, A.S., Gooseff, M.N., Singha, K., 2010. Imaging hyporheic zone solute transport using electrical resistivity. *Hydrol. Process.* 24, 948–953. <https://doi.org/10.1002/hyp.7672>.
- Ward, A.S., Gooseff, M.N., Fitzgerald, M., Voltz, T.J., Singha, K., 2014. Spatially distributed characterization of hyporheic solute transport during baseflow recession in a headwater mountain stream using electrical geophysical imaging. *J. Hydrol.* 517, 362–377. <https://doi.org/10.1016/j.jhydrol.2014.05.036>.
- Watson, A., Miller, J., Fink, M., Kralisch, S., Fleischer, M., de Clercq, W., 2019. Distributive rainfall-runoff modelling to understand runoff-to-baseflow proportioning and its impact on the determination of reserve requirements of the Verlorenvlei estuarine lake, west coast, South Africa. *Hydrol. Earth Syst. Sci.* 23, 2679–2697. <https://doi.org/10.5194/hess-23-2679-2019>.
- Watts, C.L., Hatch, C.E., Wicks, R., 2023. Mapping groundwater discharge seeps by thermal UAS imaging on a wetland restoration site. *Front. Environ. Sci.* 10.
- Weber, U.W., Cook, P.G., Brennwald, M.S., Kipfer, R., Stieglitz, T.C., 2019. A novel approach to quantify air-water gas exchange in shallow surface waters using high-resolution time series of dissolved atmospheric gases. *Environ. Sci. Technol.* 53, 1463–1470. <https://doi.org/10.1021/acs.est.8b05318>.
- Welch, E.M., Dulai, H., El-Kadi, A., Shuler, C.K., 2019. Submarine Groundwater Discharge and Stream Baseflow Sustain Pesticide and Nutrient Fluxes in Faga'alu Bay, American Samoa. *Frontiers in Environmental Science* 7.
- Welsh, M., 2023. The End of Programming. *Commun. ACM* 66, 34–35. <https://doi.org/10.1145/3570220>.
- Wen, C., Zhen, Z., Zhang, L., Yan, C., 2023. A bibliometric analysis of river health based on publications in the last three decades. *Environ. Sci. Pollut. Res.* 30, 15400–15413. <https://doi.org/10.1007/s11356-022-23267-3>.
- Werner, A.D., Alcoe, D.W., Ordens, C.M., Hutson, J.L., Ward, J.D., Simmons, C.T., 2011. Current practice and future challenges in coastal aquifer management: flux-based and trigger-level approaches with application to an Australian case study. *Water Resour. Manage.* 25, 1831–1853. <https://doi.org/10.1007/s11269-011-9777-2>.
- Wilhelmsen, K., Sawyer, A.H., Marshall, A., McFadden, S., Singha, K., Wohl, E., 2021. Laboratory Flume and Numerical Modeling Experiments Show Log Jams and Branching Channels Increase Hyporheic Exchange. *Water Resources Research* 57, e2021WR030299. <https://doi.org/10.1029/2021WR030299>.
- Williams, B.K., Szaro, R.C., Shapiro, C.D., 2009. Adaptive management: the US Department of the Interior technical guide. US Department of the Interior.
- Wilson, A.M., Morris, J.T., 2012. The influence of tidal forcing on groundwater flow and nutrient exchange in a salt marsh-dominated estuary. *Biogeochemistry* 108, 27–38. <https://doi.org/10.1007/s10533-010-9570-y>.
- Winter, T.C., 1999. Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrol. J.* 7, 28–45. <https://doi.org/10.1007/s100400050178>.
- Winter, T.C., Harvey, J.W., Franke, O.L., Alley, W.M., 1998. Ground water and surface water: A single resource (No. 1139). Circular. U.S. Geological Survey. <https://doi.org/10.3133/cir1139>.
- Woessner, W., 2000. Stream and fluvial plain ground water interactions: rescaling hydrogeologic thought. *Ground Water* 38, 423–429. <https://doi.org/10.1111/j.1745-6584.2000.tb00228.x>.
- Wohl, E., 2021. An Integrative Conceptualization of Floodplain Storage. *Reviews of Geophysics* 59, e2020RG000724. <https://doi.org/10.1029/2020RG000724>.
- Wolfe, W.W., Murgulet, D., Gyawali, B., Sterba-Boatwright, B., 2023. Long-term radon-222 (222Rn) and hydroclimatic dataset for a coastal estuary, Corpus Christi Bay, Texas. *Data in Brief* 51, 109651. <https://doi.org/10.1016/j.dib.2023.109651>.
- Wu, L., Gomez-Velez, J.D., Krause, S., Worman, A., Singh, T., Nützmann, G., Lewandowski, J., 2021. How daily groundwater table drawdown affects the diel rhythm of hyporheic exchange. *Hydrol. Earth Syst. Sci.* 25, 1905–1921. <https://doi.org/10.5194/hess-25-1905-2021>.
- Wunsch, A., Liesch, T., Broda, S., 2022. Deep learning shows declining groundwater levels in Germany until 2100 due to climate change. *Nat. Commun.* 13, 1221. <https://doi.org/10.1038/s41467-022-28770-2>.
- Xie, Y., Cook, P.G., Brunner, P., Irvine, D.J., Simmons, C.T., 2014. When Can Inverted Water Tables Occur Beneath Streams? *Groundwater* 52, 769–774. <https://doi.org/10.1111/gwat.12109>.
- Xiong, H., Wang, Y., Guo, X., Han, J., Ma, C., Zhang, X., 2022. Current status and future challenges of groundwater vulnerability assessment: A bibliometric analysis. *J. Hydrol.* 615, 128694. <https://doi.org/10.1016/j.jhydrol.2022.128694>.
- Yang, J., Griffiths, J., Zammitt, C., 2019. National classification of surface-groundwater interaction using random forest machine learning technique. *River Res. Appl.* 35, 932–943. <https://doi.org/10.1002/rra.3449>.
- Yang, S., Yang, D., Chen, J., Santisiriboon, J., Lu, W., Zhao, B., 2020. A physical process and machine learning combined hydrological model for daily streamflow simulations of large watersheds with limited observation data. *J. Hydrol.* 590, 125206. <https://doi.org/10.1016/j.jhydrol.2020.125206>.
- Yin, X., Feng, Q., Zheng, X., Wu, X., Zhu, M., Sun, F., Li, Y., 2021. Assessing the impacts of irrigated agriculture on hydrological regimes in an oasis-desert system. *J. Hydrol.* 594, 125976. <https://doi.org/10.1016/j.jhydrol.2021.125976>.
- Zarnetske, J.P., Haggerty, R., Wondzell, S.M., Baker, M.A., 2011. Dynamics of nitrate production and removal as a function of residence time in the hyporheic zone. *J. Geophys. Res.* 116. <https://doi.org/10.1029/2010JG001356>.
- Zhang, Y., He, H., Khandelwal, M., Du, K., Zhou, J., 2023b. Knowledge mapping of research progress in blast-induced ground vibration from 1990 to 2022 using CiteSpace-based scientometric analysis. *Environ. Sci. Pollut. Res.* 30, 103534–103555. <https://doi.org/10.1007/s11356-023-29712-1>.
- Zhang, S., Mao, G., Crittenden, J., Liu, X., Du, H., 2017. Groundwater remediation from the past to the future: A bibliometric analysis. *Water Res.* 119, 114–125. <https://doi.org/10.1016/j.watres.2017.01.029>.
- Zhang, X., Sawyer, A.H., Singha, K., 2023a. A numerical exploration of hyporheic zone solute transport behavior estimated from electrical resistivity inversions. *J. Hydrol.* 621, 129577. <https://doi.org/10.1016/j.jhydrol.2023.129577>.

- Zhao, S., Zhang, B., Sun, X., Yang, L., 2021. Hot spots and hot moments of nitrogen removal from hyporheic and riparian zones: A review. *Sci. Total Environ.* 762, 144168. <https://doi.org/10.1016/j.scitotenv.2020.144168>.
- Zhou, Z., Cartwright, I., 2021. Using geochemistry to identify and quantify the sources, distribution, and fluxes of baseflow to an intermittent river impacted by climate change: The upper Wimmera River, southeast Australia. *Sci. Total Environ.* 801, 149725. <https://doi.org/10.1016/j.scitotenv.2021.149725>.
- Zhou, Q., Chen, C., Zhang, G., Chen, H., Chen, D., Yan, Y., Shen, J., Zhou, R., 2018. Real-Time Management of Groundwater Resources Based on Wireless Sensors Networks. *J. Sens. Actuator Netw.* 7, 4. <https://doi.org/10.3390/jsan7010004>.
- Zhou, S., Yuan, X., Peng, S., Yue, J., Wang, X., Liu, H., Williams, D.D., 2014. Groundwater-surface water interactions in the hyporheic zone under climate change scenarios. *Environ. Sci. Pollut. Res.* 21, 13943–13955. <https://doi.org/10.1007/s11356-014-3255-3>.
- Zhu, R., Zheng, H., Croke, B.F.W., Jakeman, A.J., 2020. Quantifying climate contributions to changes in groundwater discharge for headwater catchments in a major Australian basin. *Sci. Total Environ.* 729, 138910. <https://doi.org/10.1016/j.scitotenv.2020.138910>.
- Ziliotto, F., Basilio Hazas, M., Rolle, M., Chiogna, G., 2021. Mixing Enhancement Mechanisms in Aquifers Affected by Hydropeaking: Insights From Flow-Through Laboratory Experiments. *Geophysical Research Letters* 48, e2021GL095336. <https://doi.org/10.1029/2021GL095336>.
- Zipper, S.C., Farmer, W.H., Brookfield, A., Ajami, H., Reeves, H.W., Wardropper, C., Hammond, J.C., Gleeson, T., Deines, J.M., 2022. Quantifying streamflow depletion from groundwater pumping: a practical review of past and emerging approaches for water management. *J. Am. Water Resour. Assoc.* 58, 289–312. <https://doi.org/10.1111/1752-1688.12998>.

A review of modelling of groundwater-surface water interactions in arid/semi-arid floodplains

Jolly, I.D.^{1,3} and D.W. Rassam^{2,3}

¹CSIRO Water for a Healthy Country National Research Flagship, South Australia
 Email: ian.jolly@csiro.au

²CSIRO Water for a Healthy Country National Research Flagship, Queensland
 Email: david.rassam@csiro.au

³eWater Cooperative Research Centre, Australian Capital Territory

Abstract: Floodplains and their wetlands are extremely important centres of biodiversity due to their ability to support higher amounts of biomass than surrounding upland areas. This is as a result of flooding which provides greater availability of water and more dynamic nutrient cycling. In arid and semi-arid regions, where evapotranspiration is often much greater than rainfall, this additional source of water (and nutrients) creates ecological oases for many biota in what would otherwise be a harsh environment. Floodplains also have a very important hydrogeological role in that they are the interface zone between streams and groundwater systems where exchange of water and solutes occur between these systems. The groundwater systems of this interface zone are complex and dynamic with processes such as bank storage exchange, hyporheic zone exchange, overbank inundation and recharge, evapotranspiration and groundwater pumping needing to be accounted for in water balances. While modelling of groundwater-surface water interactions in floodplains, from either water resource availability or ecological sustainability perspectives, is far from a mature area of science, there have been significant advances over the last 15 years, particularly in the area of spatial modelling. In this paper, we summarise the advances that have been made in saturated zone, unsaturated zone and combined modelling methods.

Modelling has progressed from relatively simple 1D and 2D analytical and empirical approaches to highly sophisticated 3D spatially distributed integrated modelling systems, with the advances mirroring the vast increases in desktop computing power over that time. In our view, future advances in this discipline are not so much limited by either computing power or solution methods, but rather by the availability of suitable data needed to parameterize, calibrate and validate the current-day models. As pointed out by Silberstein (2003), any model is only as good as the data upon which it is based and has been tested. Whilst calibration of very detailed process models has been possible for small (sub-reach scale) study areas, success has been limited at the reach, catchment and regional scales. However, new large-scale satellite/airborne data acquisition techniques are showing great promise, with some of the recent studies illustrating their value in model parameterization, calibration and validation. Because of severe data limitations in many areas, simple analytical models are still an attractive option in many cases as they generally only have a small number of parameters that require calibration. They can also be easily incorporated into publically available (i.e. inexpensive non-proprietary) GIS and other software frameworks. In our view, use of more sophisticated numerical models are only valid in data-rich situations or when used as research tools to study fundamental processes and parameter dependencies. At best, most detailed models will be only be usable for management as forecasting/what-if tools, and should not be considered as scenario prediction models unless they have been thoroughly calibrated and validated.

Keywords: Groundwater, surface water, interaction, floodplain, modelling

1. INTRODUCTION

Approximately one third of the world's land area is comprised of arid/semi-arid regions (Rogers, 1981). Extreme climatic variability and subsequent hydrological fluctuations are typical in these regions. The climatic variability occurs seasonally, inter-annually and on longer time frames. Consequently, arid/semi-arid areas are subjected to frequent and severe droughts and infrequent but significant floods. Climate variability and subsequent fluctuating hydrology are key drivers of ecology in arid/semi-arid environments.

To overcome the highly variable stream flow in arid/semi-arid areas, rivers have been regulated to provide year-round water supplies, transportation, waste assimilation, recreation and electricity generation. River regulation has often had profound hydrological and ecological side-effects on the rivers themselves and their adjacent floodplains (Jolly, 1996). Floodplains generally support higher amounts of biomass, have greater primary production and more dynamic nutrient cycling than surrounding upland areas. This is particularly the case in arid/semi-arid areas where potential evapotranspiration is greater than rainfall for most months and so flooding provides an additional source of water not available to non-floodplain areas (Hollis, 1990). Floodplains often also have an important hydrogeological role as the interface zone between streams and groundwater systems where exchange of water and solutes occur. Groundwater systems of this zone are complex and dynamic, with processes such as hyporheic exchange, bank storage exchange, overbank inundation, evapotranspiration and groundwater pumping needing to be accounted for in water balances (see Figure 1 for examples of some of these processes).

There have been significant advances over the last 15 years in the modelling of groundwater-surface water interactions in floodplains (water and solutes), both from a water resource availability perspective and from an ecological sustainability viewpoint. It is timely to take stock of what has been achieved to predict future directions. This short paper summarises the advances made and discusses some of the data issues which are limiting further progress in this area. The paper builds on the findings of a recent review of groundwater-surface water interactions in arid/semi-arid wetlands (Jolly et al., 2008) by expanding the scope to floodplains in general and by focusing specifically on modelling approaches. We describe saturated zone, unsaturated zone and combined methods.




Figure 1. Important floodplain hydrology processes and their interactions (from Refsgaard et al., 1998).

2. SATURATED ZONE METHODS

2.1. Bank storage

By the early 1990's considerable literature had been developed on analytical solutions to bank storage interactions (see extensive review by Rassam and Werner, 2008). The first published application of these solutions to a arid/semi-arid floodplain situation was the study of Jolly et al. (1994) which developed a 2D analytical solution for the prediction of groundwater heads beneath two 2 km long cross-sections of a floodplain which was controlled by head fluctuations of streams on either side of the floodplain. This relatively simple model provided good transient predictions of groundwater head during a flood event on the Chowilla floodplain in south-eastern Australia. The main shortcomings of the approach were (i) that it assumed that Dupuit-Forchheimer conditions hold (i.e. horizontal groundwater flow); (ii) it only predicted groundwater heads and not fluxes; and (iii) it did not apply to situations where there was vertical recharge or discharge within the floodplain (i.e. only the effects of bank storage were modelled).

Whiting and Pomeroy (1997) also developed a general 2D model of bank storage (WaTab2D). This model used a finite element approach that enabled representation of the seepage faces that form on the sides of stream channels, non-symmetrical valleys, non-symmetrical channel banks, non-uniform hydraulic geometry (i.e. Dupuit-Forchheimer conditions do not need to hold), and non-zero boundary fluxes. The un-calibrated model was used to show that the total volume of water released from bank storage is nearly proportional to the width of the floodplain, the height of the bank, and the specific yield of the aquifer. It was also used to show that the duration over which water is released from bank storage (i) increases with floodplain width and decreases with hydraulic conductivity; (ii) the rate of release decreases in an exponential-like manner; and (iii) can occur over time periods ranging from days (in gravels) to years (in clays).

The most recent work on 2D floodplain bank storage methods is the analytical model of Knight and Rassam (2007). This is a significant improvement on previous analytical methods in that the stream head fluctuations can be a random time series (described using basis splines) rather than having to be represented in functional form or as a convolution interval which must be evaluated numerically. The value of using analytical approaches such as this is that they can be easily incorporated into geographical information system (GIS) models. For example, Rassam et al. (2008) used this new method within a GIS framework to study the nitrate attenuation potential of riparian zones in the large (60,000 ha) Maroochy catchment in eastern Australia (note that this case study was in a sub-tropical area, however the method has application to arid/semi-arid areas). This approach is also being employed in a new analytical/empirical floodplain model presently being developed by the eWater Cooperative Research Centre (see paper by Rassam et al., 2009).

2.2. Overbank inundation

The most notable modelling of overbank inundation in a system which has remained relatively natural is the recent studies of the very large (1,200,000 ha) Okavango Delta in Botswana. Bauer et al. (2004) showed that transpiration from arid/semi-arid floodplains such as the Delta can be significant in situations where water tables are shallow (< 5 m below ground) as floodplain vegetation can use groundwater as a supplementary water supply in dry periods between floods. This, along with evaporation from bare soils, can lead to accumulation of salt in plant root zones and in the underlying groundwater. Overbank flooding can mitigate these accumulations of salt. Wolski et al. (2006) developed a GIS-based hydrological model of the Delta which was able to establish an overall water budget and was used to map areas that are inundated from floods. Bauer et al. (2006a) carried out coupled 3D groundwater flow and salinity transport modelling of the 10,800 ha Shashe River Valley on the fringe of the Delta using a modified version of the density-dependent SEAWAT model (Guo and Langevin, 2002). In this valley, a fresh groundwater lens had developed naturally over time due to annual flooding, but was being heavily exploited for domestic water supply and by riparian vegetation, leading to declines in groundwater levels and deterioration of water quality due to ingress of saline groundwater from adjacent aquifers. The modelling was used to gain an understanding of the complex interactions, including the interdependence of groundwater salinity and riparian vegetation transpiration. The study successfully reproduced the development of the groundwater lens and recent decline in piezometric heads, as well as explaining the old age of the saline groundwater surrounding the lens. Zimmermann et al. (2006) investigated groundwater flow and salt transport dynamics below an evaporation-dominated island in the Delta with 2D density-dependent simulations using three different numerical models. The simulation results supported geophysical observations reported in Bauer et al. (2004) of density fingering (caused by downward convection of saline water) in the groundwater beneath the island. They also showed that these density effects may be entirely overridden by lateral flow if islands are imbedded in a sufficiently transmissive aquifer within a high regional hydraulic gradient. Wolski and Savenije (2006) also studied the dynamics of floodplain-island groundwater flow in the Delta, but employed more traditional MODFLOW modelling techniques. They found that floodplain-island groundwater flow was in general very dynamic, was driven by island evaporation and transpiration, and that regular flooding was required to replenish groundwater stores. They concluded that a prolonged reduction in the duration or level of a flood could result in depletion of the groundwater storage with negative effects on the riparian vegetation.

In many floodplain systems, river regulation by weirs/locks and upstream storages has led to raised water tables beneath floodplains and reduced frequency and duration of overbank flooding. Water tables beneath floodplains can also be raised by increased groundwater inflows into floodplains from adjacent areas due to changes in land management such as the development of irrigation areas. The increased evapotranspiration and decreased recharge that can result can lead to net accumulation of salt in plant root zones and consequent floodplain vegetation dieback (Jolly, 1996). Doble et al. (2006) used relationships for evapotranspiration response to groundwater depth and salt concentration, within the MODFLOW-2000 framework, to successfully model spatial patterns of groundwater flux and salt accumulation within a small (500 ha) floodplain in south-eastern Australia (Clarks Floodplain). To more accurately model the recharge and

evapotranspiration processes in this floodplain environment, where the boundaries between recharging and nonrecharging cells change with time, Doble et al. (2009) modified the MODFLOW-2000 recharge (RCH) and segmented evapotranspiration (ETS) packages to produce a recharge-discharge function (CRD) that allows groundwater flux to be represented as a continuous process, dependent on water table depth.

Hyporheic exchange is the rapid movement of water from the stream channel into and out of the surface depths of the alluvial aquifer over short (sub-reach) distances. The importance of hyporheic exchange is that it keeps surface water in close contact with chemically reactive constituents and microbial colonies in the stream bed, thus enhancing biogeochemical reactions that influence downstream water quality and ecosystems (Harvey and Wagner, 2000). Over the last decade, there has been some progress in modeling these fine scale processes. The modeling has either been very site specific and based on finely-gridded groundwater models such as MODFLOW (e.g. Wroblicky et al., 1998; Poole et al., 2008) or has involved the development of analytical, numerical and/or statistical approaches to help understand some of the fundamental water and solute exchange processes (e.g. Qian et al., 2008; Boano et al., 2008). Research on hyporheic processes in general is a very active area of current research.

3. UNSATURATED ZONE METHODS

As described above, evapotranspiration from floodplains can result in accumulation of salt in plant root zones which can be leached back into the groundwater system by overbank floods. Slavich et al. (1999a) modelled soil salt accumulation/leaching processes at 2 sites in the Chowilla floodplain using a 1D soil-vegetation-atmosphere-transfer model (WAVES) to assess possible management options such as increasing flooding frequency by release of water from upstream storages and groundwater pumping to lower the water table. Whilst the modelling was successful, the authors recommended that the management options needed to be evaluated further at the floodplain scale to be useful for management. Slavich (1999b) then developed a salinity index (WINDS-Index), related to vegetation health, which could potentially be used in a GIS framework to evaluate management options at the floodplain scale. The approach utilised a simple 1D analytical root zone salt and water balance model which represented the salinisation process as a moving salt front. The dominant features of the more detailed WAVES simulations were adequately reproduced, and the WINDS-Index was found to be strongly dependent on the relative flood inundation time (the ratio of the duration of inundation to the duration between floods) of successive flood events. Overton et al. (2006) then implemented the WINDS-Index approach spatially in a GIS framework, tested its predictions against vegetation health mapping, and successfully used it to assess a wide range of potential management options aimed at improving the vegetation health of the large (17,400 ha) Chowilla floodplain.

4. COMBINED SATURATED AND UNSATURATED ZONE METHODS

To account for variably saturated conditions within bank storage, and the effects of density due to large salinity contrasts between stream water and groundwater, Jolly et al. (1998) applied a variable density flow and solute transport model (SUTRA; Voss, 1984) to a 4 km long cross-section of the Chowilla floodplain. The model was used to assess the importance of overbank floods in the transport of salt from the floodplain groundwater system to the lower River Murray system. The simulations showed that the mixing of flood water and saline groundwater within bank storage could explain short-term (<12 months) salt load recessions, but the observed long term (12-24 months) salt load recessions were due to the groundwater impacts of localised recharge from overbank floods at some locations on the floodplain.

To study the environmental impacts of a hydropower scheme in the 130 km long Danubian Lowland in eastern Europe, Refsgaard et al. (1998) developed an integrated modelling system comprised of catchment (MIKE SHE; Refsgaard and Storm, 1995), river (MIKE 11 and MIKE 21; Havno et al., 1995 and DHI, 1995) and unsaturated zone (DAISY; Hansen et al., 1991) models. This coupling allowed detailed modelling of the dynamic hydrological regime of this area which captured the crucial links and feedback mechanisms between the various parts of the surface and sub-surface water regimes, in particular those of the floodplain. One of the issues with this study was that while it was possible to model a complex area such as this in great detail, and the individual models could generally be validated, only a few tests on the integrated model were possible. This highlights one of the key problems with very detailed process modelling; there are often insufficient data available to both parameterise the models and then calibrate and validate them.

Hammersmark et al. (2008) have also used MIKE SHE and MIKE 11 to model surface water-groundwater interactions, in this case to quantify the hydrological effects of stream restoration in a small (230 ha) montane meadow system in northern California. The system was small enough that sufficient data was collected to enable a very good calibration and validation of the models. As a result they were able to successfully

evaluate the pre- and post- hydrological responses in the meadow to the ‘pond and plug’ restoration methods (alluvium excavated, forming ponds; excavated material used to plug incised channels; and smaller dimension channels restored to the floodplain surface level) commonly proposed for these systems.

Bauer et al. (2006b) developed a large-scale (1 km² grid) coupled 2D surface water-groundwater model to study the water balance of the entire Okavango Delta under a range of scenarios (increased water abstraction, development of upstream hydropower and irrigation schemes, channel dredging, local climatic change, local tectonic events). The model was comprised of components which simulate overland flow (based on a diffusive wave approximation of the 2D Saint-Venant equations), groundwater flow (based on a modified version of MODFLOW-96), and recharge and evapotranspiration through the unsaturated zone (using mass balance and an empirical relationship between evapotranspiration and water table depth). Due to a lack of site data, the model was not calibrated with point hydrographs but with pixel by pixel comparisons with flooding pattern time series derived from satellite imagery (151 images from 1972-2000). The authors cautioned that given the high uncertainty of both the calibrated and input parameters the model outputs could not be considered as predictions, rather they gave indications of the potential changes in flooding patterns.

There have also been extensive studies of a large (19,800 ha) floodplain of the Lower Havel River in northern Germany that have involved the development of the Integrated Modelling of Water Balance and Nutrient Dynamics (IWAN) modelling system (Krause and Bronstert, 2005, 2007; Krause et al., 2007a, 2007b). IWAN is comprised of the two way coupling (i.e. feedbacks in both directions) of the WASIM-ETH-I spatially distributed runoff generation and vertical soil water dynamics model (Schulla and Jasper, 1999) with MODFLOW-96. The most recent version of IWAN includes the SWIM (Krysanova et al., 2000) and MT3D (Zheng and Wang, 1999) models to simulate nitrate dynamics in both the unsaturated zone and groundwater (Krause et al., 2008). Use of IWAN to simulate the water balance of this floodplain indicated that interactions between groundwater and surface water controlled recharge dynamics in the floodplain and outweighed the influence of vertical percolation and root water uptake. While groundwater contributions represented only 1% of the annual total discharge of the river, in summer ~30% of the river flow was derived from groundwater discharge from the floodplain. Simulations of proposed land use and land management changes on the floodplain using IWAN showed that rates of nitrate leaching from the root zone into the groundwater and then the river could be reduced substantially, and this would have a large impact on river nitrate loads during low flow conditions.

5. CONCLUDING REMARKS

There have been very significant advances over the last 15 years in the modelling of groundwater-surface water-floodplain interactions. In addition to the studies described here there are at least a further 20 published studies focusing on temperate and tropical regions. Modelling has progressed from relatively simple 1D and 2D analytical and empirical approaches to highly sophisticated 3D spatially distributed integrated modelling systems, with advances mirroring the vast increases in desktop computing power over that time.

In our view future advances in this discipline are not so much limited by either computing power or solution methods, but rather by the availability of suitable data needed to parameterize, calibrate and validate the current-day models. As always, data collection is expensive, and therefore generally very difficult to fund. The scaling down of routine hydrological and hydrogeological monitoring in many jurisdictions has further exacerbated this problem. Many natural resource management policy makers and managers tend to consider process models as “silver bullets” without fully understanding that any model is only as good as the data upon which it is based and has been tested (Silberstein, 2003). Whilst calibration of very detailed process models has been possible for small (sub-reach scale) study areas (i.e. Hammersmark et al., 2008; Doble et al., 2006), success has been limited at the reach, catchment and regional scales. However, new large-scale satellite/airborne data acquisition techniques are showing great promise; the studies of Overton et al. (2006) and Bauer et al. (2006a) illustrate their value in model parameterization, calibration and validation.

In data-poor situations, simple analytical models are still an attractive option as they generally only have a small number of parameters that require calibration. They can also be easily incorporated into publically available (i.e. inexpensive non-proprietary) GIS and other software frameworks (e.g. Overton et al., 2006; Rassam et al., 2008, 2009). In our view, use of more sophisticated numerical models are only valid in data-rich situations or when used as research tools to study fundamental processes and parameter dependencies. At best, most detailed models will only be usable for management as forecasting/what-if tools, and should not be considered as scenario prediction models unless they have been thoroughly calibrated and validated.

ACKNOWLEDGMENTS

This review was funded by the eWater Cooperative Research Centre (and its partner agencies) and the National Water Commission Raising National Water Standards Program, under the Project “Australian Hydrological Modelling Initiative - Groundwater Surface Water Interaction Tool”.

REFERENCES

- Bauer, P., G. Thabeng, F. Stauffer, and W. Kinzelbach, (2004), Estimation of the evapotranspiration rate from diurnal groundwater level fluctuations in the Okavango Delta, Botswana. *Journal of Hydrology*, 288, 344-355.
- Bauer, P., R.J. Held, S. Zimmermann, F. Linn, and W. Kinzelbach, (2006a), Coupled flow and transport modelling in semi-arid environments: The Shashe River Valley, Botswana. *Journal of Hydrology*, 316, 163-183.
- Bauer, P., T. Gumbricht, and W. Kinzelbach, (2006b), A regional coupled surface water/groundwater model of the Okavango Delta, Botswana. *Water Resources Research*, 42, W04403, doi:10.1029/2005WR004234.
- Boano, F., R. Revelli, and L. Ridolfi, (2008), Reduction of the hyporheic zone volume due to the stream-aquifer interaction. *Geophysical Research Letters*, 35, L09401, doi:10.1029/2008GL033554.
- DHI (1995), MIKE 21 Short Description. Danish Hydraulic Institute, Horsholm.
- Doble, R.C., C.T. Simmons, I.D. Jolly, and G.R. Walker, (2006), Spatial relationships between vegetation cover and irrigation-induced groundwater discharge on a semi-arid floodplain, Australia. *Journal of Hydrology*, 329, 75-97.
- Doble, R.C. C.T Simmons, and G.R. Walker, (2009), using MODFLOW 2000 to model ET and recharge for shallow ground water problems. *Groundwater*, 47, 129-135.
- Guo, W. and C.D. Langevin, (2002), User's Guide to SEAWAT: A Computer Program For Simulation of Three-Dimensional Variable Density Groundwater Flow. Techniques in Water Resources Investigations 06-A7, U.S. Geological Survey, Tallahassee.
- Hammersmark, C.T., M.C.Rains, and J.F. Mount, (2008), Quantifying the hydrological effects of stream restoration in a montane meadow, northern California, USA. *River Research and Applications*, 24, 735-753.
- Hansen, S., H.E. Jensen, N.E., Neilsen, and H. Svendsen, (1991), Simulation of nitrogen dynamics and biomass production in winter wheat using the Danish model DAISY. *Fertilizer Research*, 27, 245-259.
- Harvey, J.W. and Wagner, B.J., (2000), Quantifying hydrologic interactions between streams and their subsurface hyporheic zones. In *Streams and Ground Waters* (Eds J.B. Jones and P.J. Mulholland), Academic Press, San Diego, pp. 3-44.
- Havno, K., M.N. Madsen and J. Dorge, (1995), MIKE 11 – A Generalized River Modelling Package. In V.P. Singh (Ed), *Computer Models of Watershed Hydrology*, Water Resources Publications, pp. 733-782.
- Hollis, G.E. (1990), Environmental impacts of development on wetlands in arid and semi-arid lands. *Hydrological Science*, 35, 411-428.
- Jolly, I.D. (1996), The effects of river management on the hydrology and hydroecology of arid and semi-arid floodplains. In *Floodplain Processes* (Eds M.G. Anderson, D.E. walling and P.D. Bates), John Wiley and Sons, Chichester, pp. 577-609.
- Jolly, I.D., G.R. Walker, and K.A. Narayan, (1994), Floodwater recharge processes in the Chowilla anabranch system, South-Australia. *Australian Journal of Soil Research*, 32, 417-435.
- Jolly, I.D., K.A. Narayan, D. Armstrong, and G.R. Walker, G.R., (1998), The impact of flooding on modelling salt transport process to streams. *Environmental Modelling and Software*, 13, 87-104.
- Jolly, I.D., K.L., McEwan, and K.L. Holland, (2008), A review of groundwater-surface water interactions in arid/semi-arid wetlands and the consequences of salinity for wetland ecology. *Ecohydrology*, 1, 43-58.
- Knight, J.H. and D.W. Rassam, (2007), Groundwater head responses due to random stream stage fluctuations using basis splines. *Water Resources Research*, 43, W06501, doi:10.1029/2006WR005155.
- Krause, S and A. Bronstert, (2005), An advanced approach for catchment delineation and water balance modelling within wetlands and floodplains. *Advances in Geosciences*, 5, 1-5.
- Krause, S. and A. Bronstert, (2007), The impact of groundwater-surface water interactions on the water balance of a mesoscale lowland river catchment in northeastern Germany. *Hydrological Processes*, 21, 169-184.
- Krause, S.; A., Bronstert, and E. Zehe, (2007a), Groundwater-surface water interactions in a north German lowland floodplain - Implications for the river discharge dynamics and riparian water balance. *Journal of Hydrology*, 347, 404-417.
- Krause, S.; J. Jacobs, and A. Bronstert, (2007b), Modelling the impacts of land-use and drainage density on the water balance of a lowland-floodplain landscape in northeast Germany. *Ecological Modelling*, 200, 475-492.
- Krause, S.; J. Jacobs, A. Voss, A. Bronstert, and E. Zehe, E., (2008), Assessing the impact of changes in landuse and management practices on the diffuse pollution and retention of nitrate in a riparian floodplain. *Science of the Total Environment*, 389, 149-164.
- Krysanova, V., F. Wechsung, J. Arnold, R. Srinivasan, and J. Williams, (2000), SWIM (Soil and Water Integrated Model) User Manual. PIK Report No. 69. Potsdam Institute for Climate Impact Research, Potsdam.

- Overton I.C., I.D. Jolly, P.G. Slavich, M.M. Lewis, and G.R. Walker, (2006), Modelling vegetation health from the interaction of saline groundwater and flooding on the Chowilla floodplain, South Australia. *Australian Journal of Botany*, 54, 207-220.
- Poole, G.C., S.J. O'Daniel, K.L. Jones, W.W. Woessner, E.S. Bernhardt, A.M. Helton, J.A. Stanford, B.R. Boer, and T.J. Beechie, (2008), Hydrologic spiralling: The role of multiple interactive flow paths in stream ecosystems. *River Research and Applications*, 24, 1018-1031.
- Qian, Q., V.R. Voller and H.G. Stefan, (2008), A vertical dispersion model for solute exchange induced by underflow and periodic hypoheic flow in a stream gravel bed. *Water Resources Research*, 44, W07422, doi:10.1029/2007WR006366.
- Rassam, D.W. and A.D. Werner, (2008), Review of Groundwater-Surfacewater Interaction Modelling Approaches and Their Suitability for Australian Conditions. eWater Technical Report. eWater Cooperative Research Centre, Canberra.
- Rassam, D.W., D.E. Pagendam, and H.M. Hunter, (2008), Conceptualisation and application of models for groundwater-surface water interactions and nitrate attenuation potential in riparian zones. *Environmental Modelling and Software*, 23, 859-875.
- Rassam, D., T. Pickett, and J. Knight, (2009), The Floodplain Model. In Proceedings of 18th World IMACS/MODSIM Congress, Cairns, Australia, 13-17 July 2009.
- Refsgaard, J.C. and B. Storm, (1995), MIKE SHE. In V.P. Singh (Ed), *Computer Models of Watershed Hydrology*, Water Resources Publications, pp. 809-846.
- Refsgaard, J.C., H.R. Sorensen, I. Mucha, D. Rodak, Z. Hlavaty, L. Banský, J. Klucovská, J. Topolská, J. Takáč, V. Kosc, H.G. Enggrob, P. Engesgaard, J.K. Jensen, J. Fiselier, J., Griffioen, and S. Hansen, (1998), An integrated model for the Danubian Lowland - Methodology and applications. *Water Resources Management*, 12, 433-465.
- Rogers, B.R. (1981), Fools rush in, Part 3: selected dryland areas of the world. *Arid Lands Newsletter*, 14, 24-24.
- Schulla, J. and K. Jasper, (1999), Modellbeschreibung WASIM-ETH. Swiss Federal Institute of Technology (ETH), Zurich.
- Silberstein, R. (2003), Why collect data when we can model it? In Proceedings of MODSIM 2003, Townsville, Australia, 14-17 July 2003, 915-920.
- Slavich, P.G., G.R. Walker, I.D. Jolly, T.J. Hatton, and W.R. Dawes, (1999a), Dynamics of *Eucalyptus largiflorens* growth and water use in response to modified watertable and flooding regimes on a saline floodplain. *Agricultural Water Management*, 39, 245-264.
- Slavich, P.G., G.R. Walker, and I.D. Jolly, (1999b), A flood history weighted index of average root-zone salinity for assessing flood impacts on health of vegetation on a saline floodplain. *Agricultural Water Management*, 39, 135-151.
- Voss, C.I. (1984), SUTRA: A Finite-Element Simulation Model For Saturated-Unsaturated Fluid-Density-Dependent Groundwater Flow With Energy Transport or Chemically Reactive Species Solute Transport. Water Resources Investigations Report 84-4369, U.S. Geological Survey, Reston.
- Werner A.D., M.R. Gallagher, and S.W. Weeks, (2006), Regional-scale fully coupled modelling of stream-aquifer interaction in a tropical catchment. *Journal of Hydrology*, 328, 497-510.
- Whiting, P.J. and M.A. Pomeroy, (1997), Numerical study of bank storage and its contribution to streamflow. *Journal of Hydrology*, 202, 121-136.
- Wolski, P. and H.H.G. Savenije, (2006), Dynamics of floodplain-island groundwater flow in the Okavango Delta, Botswana. *Journal of Hydrology*, 320, 283-301.
- Wolski, P., H.H.G. Savenije, M. Murray-Hudson, and T. Gumbrecht, (2006), Modelling of the flooding in the Okavango Delta, Botswana, using a hybrid reservoir-GIS model. *Journal of Hydrology*, 331, 58-72.
- Wroblicky, G.J., M.E. Campana, H.M. Valett, and C.N. Dahm, (1998), Seasonal variation in surface-subsurface water exchange and lateral hyporheic area of two stream-aquifer systems. *Water Resources Research*, 34, 317-328.
- Zheng, C and P.P. Wang, (1999), MT3DMS: A Modular Three-Dimensional Multi-Species Transport Model For Simulation of Advection, Dispersion, and Chemical Reactions of Contaminants in Groundwater Systems; Documentation and User's Guide. Contract report SERDP-99-1, US Army Corps of Engineers Engineer Research and Development Center, Vicksburg.
- Zimmermann, S., P. Bauer, R. Held, W. Kinzelbach, and J.H. Walther, (2006), Salt transport on islands in the Okavango Delta: Numerical investigations. *Advances in Water Resources*, 29, 11-29.




**US Army Corps
of Engineers®**

Hydrologic Engineering Center

HEC-RAS

River Analysis System



HEC-RAS Hydraulic Reference Manual

Version 6.5

Exported - March 2024

4 THEORETICAL BASIS FOR ONE-DIMENSIONAL AND TWO-DIMENSIONAL HYDRODYNAMIC CALCULATIONS

This chapter describes the methodologies used in performing the one-dimensional (1D) steady flow and unsteady flow calculations, as well as the two-dimensional (2D) unsteady flow calculations within HEC-RAS. The basic equations are presented along with discussions of the various terms. Solution schemes for the various equations are described. Discussions are provided as to how the equations should be applied, as well as applicable limitations.

1D Steady Flow Water Surface Profiles

HEC-RAS is currently capable of performing 1D water surface profile calculations for steady gradually varied flow in natural or constructed channels. Subcritical, supercritical, and mixed flow regime water surface profiles can be calculated. Topics discussed in this section include: equations for basic profile calculations; cross section subdivision for conveyance calculations; composite Manning's n for the main channel; evaluation of the mean kinetic energy head (velocity weighting coefficient α); friction loss evaluation; contraction and expansion losses; computational procedure; critical depth determination; applications of the momentum equation; air entrainment in high velocity streams; and limitations of the steady flow model.

Equations for Basic Profile Calculations

Water surface profiles are computed from one cross section to the next by solving the Energy equation with an iterative procedure called the standard step method. The Energy equation is written as follows:

$$1) \quad Z_2 + Y_2 + \frac{\alpha_2 V_2^2}{2g} = Z_1 + Y_1 + \frac{\alpha_1 V_1^2}{2g} + h_e$$

Symbol	Description	Units
Z_1, Z_2	elevation of the main channel inverts	
Y_1, Y_2	depth of water at cross sections	
V_1, V_2	average velocities (total discharge/ total flow area)	
α_1, α_2	velocity weighting coefficients	
g	gravitational acceleration	
h_e	energy head loss	

A diagram showing the terms of the energy equation is shown in the figure below.