

**BEFORE THE HON'BLE NATIONAL GREEN TRIBUNAL PRINCIPAL
BENCH, NEW DELHI**

IN

ORIGINAL APPLICATION NO. 100 /2025

IN THE MATTER OF:

AMIT KUMAR YADAV

.... APPLICANT.

VS.

STATE OF U.P. & ORS.

.... RESPONDENT.

INDEX

<u>S. NO.</u>	<u>PARTICULARS</u>	<u>PAGE NO.</u>
1.	Supplementary Affidavit of District Magistrate, Hamirpur in furtherance of hearing dated 22.05.2026	
2.	<u>Annexure No. SA -1: -</u> A copy of the Letter dated 25.05.2026 issued by the Deponent	
3.	<u>Annexure No. SA-2: -</u> A copy of the Letter dated 25.05.2026 issued by the Central Ground Water Board	
4.	<u>Annexure No. SA – 3 (Collectively): -</u> A copy of the table containing the incorrectly mentioned years i.e. 2022–2023, along with the corresponding corrected rainfall data pertaining to the years 2023–2024	

DATED: - 26.05.2026

THROUGH

Ankit Verma

(ANKIT VERMA)

STANDING COUNSEL STATE OF UP

A-15 FF, NIZAMUDDIN EAST, NEW DELHI- 110013

MOB:- 0999080440

Email-ankit.scngtup@gmail.com

**BEFORE THE HON'BLE NATIONAL GREEN TRIBUNAL PRINCIPAL
BENCH, NEW DELHI**

IN

ORIGINAL APPLICATION NO. 100 /2025

IN THE MATTER OF:

AMIT KUMAR YADAV

VS.

STATE OF U.P. & ORS.

.... RESPONDENT.

**SUPPLEMENTARY AFFIDAVIT OF DISTRICT MAGISTRATE,
HAMIRPUR IN FURTHERENCE OF HEARING DATED 22.05.2026**

The Respondent No. 3 herein states as under:

MOST RESPECTFULLY SHOWETH:


I, Abhishek Goyal, aged about 33 years, S/o Rajesh Kumar Goyal R/o H.N. 5A Lokmanya Tilak Nagar Ward no-5, P.S.- Chandauli Tahshil Chandauli District- Chandauli Occupation-Government Servant, Qualification Graduation, presently posted as District Magistrate – Hamirpur, the deponent, do hereby solemnly state and affirm as under:-

1. That I am the above-mentioned answering Respondent No. 3 and is duly competent to file the present Supplementary Affidavit. That the Deponent is well conversant with the facts and the circumstances of the instant case and is competent to swear this Supplementary Affidavit.
2. That the Deponent is posted as District Magistrate-Hamirpur, since 21.04.2026 and is swearing this Supplementary Affidavit in his official capacity.




3. That the present Supplementary Affidavit is being filed pursuant to the query raised by this Hon'ble Tribunal during the course of hearing held on 22.05.2026.
4. That the query so raised by this Hon'ble Tribunal pertains to the rainfall data downloaded by the Office of the Sub-Divisional Committee from the official website of the Central Ground Water Board. It is most respectfully submitted that, owing to an inadvertent clerical and typographical error in Table 2.0 of the said report, the years "2022" and "2023" came to be mentioned in place of "2023" and "2024" respectively. It is further submitted that the figures reflected under the headings "Actual Rainfall 2022" and "Actual Rainfall 2023" in fact correspond to the actual rainfall data pertaining to the years 2023 and 2024 respectively. The rainfall figures contained therein are otherwise accurate and have been duly extracted from the "Ground Water Bulletin, Uttar Pradesh, May 2024" published by the Central Ground Water Board.
5. That in respectful compliance of the observations and queries raised by this Hon'ble Tribunal, the Deponent herein addressed a communication dated 25.05.2026 to the Regional Director, Central Ground Water Board, requesting therein verification of the aforesaid report and issuance of the authenticated clarification at the earliest, so as to enable the same to be placed before this Hon'ble Tribunal for its kind consideration. A copy of the Letter dated 25.05.2026 issued by the Deponent is being annexed as **Annexure No. SA-1** to the present Supplementary Affidavit.

6. That in response thereto, the Central Ground Water Board, vide its Letter dated 25.05.2026 addressed to the Deponent herein, categorically clarified



A handwritten signature in blue ink, consisting of stylized initials.

that the "Ground Water Level Bulletin, Uttar Pradesh, May 2024" is a duly verified report of the Board and is also available in the public domain on its official website. It was further clarified therein that, upon examination of the report, a minor correction was noticed in Table 2 (Page No. 8), wherein the years "2022" and "2023" had inadvertently been mentioned in place of "2023" and "2024" respectively. However, the Board unequivocally confirmed that the rainfall data reflected in the said table is correct and is fully consistent with the data supplied by the India Meteorological Department. A copy of the Letter dated 25.05.2026 issued by the Central Ground Water Board is being annexed as **Annexure No. SA-2** to the present Supplementary Affidavit.

7. That the aforesaid inadvertent discrepancy in mentioning the years corresponding to the rainfall data has, in any event, no material or substantive bearing upon the ultimate findings and quantities reflected in the Replenishment Studies, and as such, no prejudice whatsoever is caused nor does anything substantial turn upon the said clerical error. For the kind and convenient perusal of this Hon'ble Tribunal, the table containing the incorrectly mentioned years i.e. 2022-2023, along with the corresponding corrected rainfall data pertaining to the years 2023-2024, is collectively being annexed as **Annexure No. SA-3** to the present Supplementary Affidavit.

8. That the Deponent most respectfully submits that he remains duty bound to ensure faithful compliance of all orders and directions passed by this Hon'ble Tribunal and accordingly prays that the present Supplementary




Affidavit be graciously taken on record for the just, proper, and effective adjudication of the present Original Application, in the interest of justice.


DEPONENT


VERIFICATION


Verified at H.M.R......, on 25/5/26... that the contents of the paras 1 to of this affidavit are true and correct to the best of my knowledge. No part of it is false and nothing material has been concealed therefrom.


DEPONENT




Execution of this...
And Signed Before Me By Smt
.....
In The Court Premises My Office
At H.M.R. Dated 25/5/26




Kalicharan Nisha
Advocate & Notary
Distt.-Hamirpur (U.P.)

भारत निर्वाचन आयोग
ELECTION COMMISSION OF INDIA
मतदाता फोटो पहचान पत्र - ELECTOR PHOTO IDENTITY CARD

IUA4227245



नाम : अभिषेक गोयल
Name : ABHISHEK GOYAL
पिता का नाम : राजेश कुमार गोयल
Father's Name : RAJESH KUMAR GOYAL

लिंग / Gender : पुरुष / Male
जन्म तिथि/आयु : 15-07-1993
Date of Birth/ Age :
पता : म.क्र.5 ए, लोकमान्य तिलक नगर वार्ड नं० ५, वार्ड नं०-5 लोकमान्य
तिलक नगर, थाना-चन्दाली, तहसील-चन्दाली, जिला-चन्दाली-232104
Address : H.N.5 A, LOKMANYA TILAK NAGAR WARD NO-
5, WARD NO-5 LOKMANYA TILAK NAGAR, P.S.-
CHANDAULI, Tahsil-Chandauli, Dist-Chandauli, 232104

दिनांक / Date : 28-04-2019 निर्वाचक रजिस्ट्रीकरण अधिकारी
Electoral Registration Officer

विधान सभा क्षेत्र की संख्या व नाम : 380-मुगतसराय
Assembly Constituency No. and Name : 380-Mughalsarai

भाग संख्या व नाम : 396-महेन्द्र टेक्निकल इण्टर कालेज चन्दाली कंस०
Part No. and Name : 396-MAHENDRA TECKNIKAL
COLLEGE CHANDAULI ROOM N05

नोट / Note :

1. इस कार्ड को धारण करने मात्र से यह कोई गारंटी नहीं है कि आप वर्तमान निर्वाचक नामावली में निर्वाचक हैं। कृपया अपना नाम प्रत्येक चुनाव से पहले वर्तमान नामावली में जांच लें।
Mere possession of this card is no guarantee that you are elector in the current electoral roll. Please check your name in the current electoral roll before every election.
2. इस कार्ड में उल्लिखित जन्मतिथि को निर्वाचक नामावली में पंजीकरण के अलावा अन्य किसी भी स्थिति में आयु के प्रमाण के रूप में नहीं माना जाएगा।
Date of Birth mentioned in this card shall not be treated as a proof of age / D.O.B. for any purpose other than registration in electoral roll. B13P87 380 396 0035201922150219




प्रेषक,

जिलाधिकारी
हमीरपुर।

सेवा में,

क्षेत्रीय निदेशक,
केन्द्रीय भूजल आयोग,
उत्तरी क्षेत्र, लखनऊ।

संख्या- 340 / खनिज-एम0एम0सी-तीस-विविध (2026-27)

दिनांक- 25.05.2026

विषय- केन्द्रीय भूजल आयोग द्वारा पब्लिक डोमेन प्रसारित रिपोर्ट-2024 के सत्यापन के सम्बन्ध में।

महोदय,

उपर्युक्त विषयक अवगत कराना है कि जनपद हमीरपुर के जिला सर्वेक्षण रिपोर्ट निर्मित किये जाने के दौरान आपके वेबसाइट से प्रसारित रिपोर्ट के आंकड़े वर्ष-2024 को जिला सर्वेक्षण रिपोर्ट में शामिल किया गया था। वर्तमान में जिला सर्वेक्षण रिपोर्ट-2024 जनपद हमीरपुर के विरुद्ध याचिका मा0 न्यायालय हरित अधिकरण नई दिल्ली में ओ0ए0 संख्या-09/2025 तथा 100/2025 योजित किया गया है जिसमें उक्त रिपोर्ट के भी सत्यापन की प्रति प्रस्तुत की जानी है जबकि वर्तमान में उक्त रिपोर्ट (C.G.W.B. Northern Region) आपके वेबसाइट पर प्रदर्शित नहीं हो रहा है इसके कारण उक्त रिपोर्ट की सत्यापन में असुविधा हो रही है।

अतः C.G.W.B. Northern Region की वर्ष-2024 की रिपोर्ट को संलग्न कर इस अनुरोध के साथ प्रेषित है कि उक्त रिपोर्ट का सत्यापन कराते हुए अविलम्ब उपलब्ध कराने का कष्ट करें जिससे मा0 न्यायालय हरित अधिकरण नई दिल्ली को प्रति ससमय उपलब्ध कराया जा सके।

(राकेश कुमार)

अपर जिलाधिकारी (वि0/रा0)
हमीरपुर।

भारत सरकार
जल शक्ति मंत्रालय
जल संसाधन, नदी विकास और गंगा संरक्षण विभाग
केन्द्रीय भूमि जल बोर्ड
उत्तरी क्षेत्र, भूजल भवन,
सेक्टर-बी, सीतापुर रोड योजना,
अलीगंज, लखनऊ - 226021
दूरभाष : (0522)-2363812
ई-मेल: rdnr-cgwb@nic.in



GOVERNMENT OF INDIA
MINISTRY OF JAL SHAKTI
DEPARTMENT OF WATER RESOURCES, RIVER DEVELOPMENT
AND GANGA REJUVENATION
CENTRAL GROUND WATER BOARD
NORTHERN REGION, BHUJAL BHAWAN
SECTOR-B, SITAPUR ROAD YOJANA
ALIGANJ, LUCKNOW-226 021
PHONE : (0522)-2363812
E-mail: rdnr-cgwb@nic.in

न. -4(232)/के.भू.ज.बो./उ.क्षे./एस&आइ/2026

580

दिनांक :

25 MAY 2026

सेवा में,

जिलाधिकारी महोदय,
हमीरपुर, उत्तर प्रदेश।

विषय: केन्द्रीय भूमि जल बोर्ड द्वारा पब्लिक डोमेन में प्रसारित रिपोर्ट-2024 के सत्यापन के संबंध में।
महोदय,

पत्रांक संख्या-340/खनिज-एम0एम0सी0-तीस-विधि(2026-27) दिनांक 25.05.2026 के संदर्भ में, आपको अवगत कराना है की Ground Water Level Bulletin, Uttar Pradesh, May 2024 की रिपोर्ट केन्द्रीय भूमि जल बोर्ड की सत्यापित रिपोर्ट है जो की इसके official website (<https://cgwb.gov.in/>) पर भी पब्लिक डोमेन में उपलब्ध है।

हालाँकि, जांच करने पर यह पाया गया कि रिपोर्ट की सारणी संख्या-2 (पृष्ठ सं.-8) में एक मामूली संशोधन है, जिसमें वर्ष 2022 एवं 2023 अंकित किए गए हैं, जबकि इनके स्थान पर क्रमशः 2023 एवं 2024 होना चाहिए। तथापि, सारणी में प्रदर्शित वर्षा संबंधी आँकड़े सही हैं तथा आईएमडी द्वारा उपलब्ध कराए गए आँकड़ों के अनुरूप हैं।

यह आपके आवश्यक जानकारी हेतु प्रेषित है।


भवदीय,

(एस. के. स्वरूप)

क्षेत्रीय निदेशक के तकनीकी सचिव

प्रतिलिपि:-

1. क्षेत्रीय निदेशक, केन्द्रीय भूमि जल बोर्ड, उत्तरी क्षेत्र, लखनऊ को जानकारी हेतु।



GROUND WATER LEVEL BULLETIN
UTTAR PRADESH
 MAY 2024

ABSTRACT

Ground water level Scenario during May – 2024 highlighting the finding, status of ground water level in different aquifer and its seasonal, annual and decadal comparison.

CGWB, NORTHERN REGION LUCKNOW

1.0. INTRODUCTION

Groundwater bulletin is prepared by CGWB depicting changes in groundwater regime of the country through different seasons. It is an effort to obtain information on groundwater levels through representative monitoring wells. The important attributes of groundwater regime monitoring are groundwater level. The natural conditions affecting the groundwater regime involve climatic parameters like rainfall, vapor transpiration etc. whereas anthropogenic in fluences include pumpage from the aquifer, recharge due to irrigation systems and other practices like waste disposal etc.

Groundwater levels are being measured by Central Ground Water Board four times a year during January, March, April, May, August and November. The regime monitoring started in the year 1969 by Central Groundwater Board. A network of 1464 observation wells called National Hydrograph Network Stations (NHNS), as on 30.05.2024, located all over the State is being monitored.

2.0. STUDY AREA

The State of Uttar Pradesh forms a part of vast Gangetic Alluvial Plain covering an area of 2,40,928 Sq. Km. and lies between North latitude 23°52'12" & 30°24'30" and East longitude 77°05'38" & 84°38'30". It is bounded by Uttarakhnad on the NW, Nepal on the NE, Bihar on the

East, Madhya Pradesh in the South, and Haryana, Delhi & Rajasthan in the West as shown in figure 1.




Figure-1: Map showing major aquifers and administrative divisions of UP

The state is covered with rich fertile soil and underlain by a large thickness of alluvium making it one of the richest grounds waterrepositories of the world. Ground water is a major source of fresh water on earth. It is the most dependable source of water, comparatively

free from the vagaries of nature, easily accessible, available at the point of use and economical. Hence it is being developed indiscriminately and the ground water reservoir is stressed. The State being the most populous in the country, with a population density of 829 persons per sq. km and a high rate of population growth (20%), its demand for water is soaring. Also due to industrialization, urbanization and modern farming practices, its quality is also at stake.

demand for water is soaring. Also due to industrialization, urbanization and modern farming practices its quality is also at stake.

The food production in Uttar Pradesh is commensurate with the self-sufficiency of the country. One of the major contributors for this sufficiency is irrigation. To meet this high irrigational requirement, water resources are being increasingly developed. Ground water contributes to about 71 % of the irrigation needs of the State. The indiscriminate development of ground water has resulted in depletion of groundwater storage and lowering of water level in certain areas on one hand. On other side the surface water development in are as having shallow water level has resulted in water logging and soil salinization. The geology and structure of the formations existing in an area control by occurrence and movement of ground water. The geomorphic conditions also have a great impact on ground water scenario. The larger part of the State is underlain by fluvial sediments laid down in

the fore deep between Plateau region in south and Himalayas in north during the Quaternary period by the Indus-Ganga system of drainage over the Precambrian topography existing during geological past. These deposits owe their origin to riverine activity. The southern part of the State has entirely different geological conditions being underlain by Precambrian formations under a thin alluvial cover. Broadly, the State can be divided into two hydrogeological units.

1. Unconsolidated zone.
2. Consolidated, hardrock zone.

The hydrogeological conditions of the above two units widely differ and are discussed subsequently in brief.

Unconsolidated Zone:

This unit covers nearly 85% of the State area. The unconsolidated formations comprising the area have been deposited through mighty rivers originating from the great Himalayan Mountains. These sediments are an admixture of pebble, gravel, sand, silt, clay and kankar. The sediments are generally coarser in the north and gradually become finer southeastward along downstream of the drainage which is a typical feature of fluvial deposits. This zone consists of mainly two parts, the Terai and the Alluvial Plain. However, the foot hill zone is very small part of Bhabar belt and lies in the northern parts of Bijanore

and Saharanpur districts. The Terai is a narrow-disconnected belt along the northwestern fringe of the State. The Alluvial Plain occupies the area south of Terai and can further be divided into two sub units - Younger Alluvium and Older Alluvium.

The younger alluvium occurs mostly along the present-day flood plain area. The continuous shifting of the drainage network with time caused reworking of their earlier deposits giving rise to the younger alluvium. The older alluvium occupying comparatively high area covers major part of the Plain. A typical characteristic of older alluvium is formation of kankar within itself due to leaching of calcium carbonate under favourable climatic conditions. The kankar occasionally forms pans restricting downward movement of water.

The thickness of alluvial sediments is variable and generally goes upto 500m. below which occur the semi-consolidated Upper Siwalik formations. The Shallower basement occurs in isolated areas which are known as "Basement highs." This unconsolidated zone is porous and permeable with primary intergranular porosity and has good ground water potential. The sub-surface correlation of formations in the state has shown presence of several aquifers down to a depth of 750 m below the ground. These aquifers mainly encountered in Central Ganga Plain have been grouped on the basis of lithological characters as well as based on interpretation of electrical logs of Boreholes drilled and are as

follows:

1. First aquifer	0.0 – 150.00 mbgl
2. Second aquifer	160.00 – 210.00 mbgl
3. Third aquifer	250.00 – 360.00 mbgl
4. Forth deep aquifer	380.00 – 600.00 mbgl

The upper part of the first aquifer down to 50 mbgl is the main source of drinking water through hand pumps and dug wells and is unconfined in nature. The first aquifer as a whole which is under unconfined to semi-confined conditions, it is the most potential aquifer group which is the main source of groundwater in the State extensively exploited through private as well as Government tube wells to meet the drinking water and irrigation needs. The deeper aquifers are confined in nature being exploited to a very limited extent. The yield of the second aquifer is limited while the third aquifer is potential. The shallow and phreatic aquifers are under heavy stress.

Consolidated Zone:

The Bundelkhand Vindhyan plateau region is underlain by a variety of Precambrian formations, mostly granite and granite gneisses, Vindhyan sandstone, limestone & shale, under a thin alluvial cover or without alluvial cover. As such these formations are hard and compact and devoid of any primary porosity. The ground water in these formations

occurs in the secondary porosity of these formations. The secondary porosity has developed due to cracks and fractures which are open at the surface and tighten at depth. The ground water occurs under unconfined or water level conditions in these formations.

The alluvial sediments of moderate depth along the river course sand in valleys form potential ground water repositories. The weathered mantle over the entire unit so forms potential aquifers. These aquifers are being monitored mostly through open wells over the area.

3.0. GROUNDWATER LEVEL MONITORING

Central Ground Water Board, Northern Region, is monitoring changes in groundwater regime in Uttar Pradesh State on quarterly basis continuously. This is facilitated by a network of monitoring stations in the State located in diverse hydrogeological and geomorphic units. The number of operational wells till May 2024 was 1464 which include 1001 dug wells and 463 piezometers shown in figure 2. The district-wise breakup of the water level monitoring stations is given in Table-1.

Table-1: District-wise distribution of water level monitoring station

S.No	District	Number Of Water Level Monitoring Stations		
		Dw	Pz	Total
1	Agra	6	11	17

S.No	District	Number Of Water Level Monitoring Stations		
		Dw	Pz	Total
2	Aligarh	9	4	13
3	Ambekarnagar	9	16	25
4	Amethi	36	2	38
5	Amroha		9	9
6	Auraiya	9	2	11
7	Ayodhya	13	5	18
8	Azamgarh	22	10	32
9	Baghpat	2	9	11
10	Bahraich	20	15	35
11	Ballia	21	2	23
12	Balrampur	17	9	26
13	Banda	10	18	28
14	Barabanki	37	4	41
15	Barilly	11	4	15
16	Basti	14	3	17
17	Bhadohi	7		7
18	Bijnor	6	14	20
19	Budaun		10	10

S.No	District	Number Of Water Level Monitoring Stations 2024		
		Dw	Pz	Total
20	Bulandshahr	2	14	16
21	Chandauli	15	4	19
22	Chitrakoot	10	14	24
23	Deoria	28	1	29
24	Etah	2	4	6
25	Etawah	11	2	13
26	Farrukhabad	1	4	5
27	Fatehpur	13	14	27
28	Firozabad		6	6
29	Gautam Buddha Nagar		8	8
30	Ghaziabad		3	3
31	Ghaziipur	22	11	33
32	Gonda	25	4	29
33	Gorakhpur	15	16	31
34	Hamirpur	12	11	33
35	Hapur		4	4
36	Hardoi	16	7	23
37	Hathras	5	4	9

S.No	District	Number Of Water Level Monitoring Stations 2024		
		Dw	Pz	Total
38	Jalaun	32	6	38
39	Jaunpur	30	11	41
40	Jhansi	20	2	22
41	Kannauj	11	2	13
42	Kanpur Dehat	12	1	13
43	Kanpur Nagar	16	1	17
44	Kasganj	4	11	15
45	Kaushambi	9	2	11
46	Kheri	26	5	31
47	Kushinagar	28		28
48	Lalitpur	19	4	23
49	Lucknow	9	15	24
50	Mahoba	10	6	16
51	Mahrajganj	13	1	14
52	Mainpuri	4	5	9
53	Mathura	17	3	20
54	Mau	13	3	16
55	Meerut	1	12	13

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
56	Mirzapur	25		25
57	Moradabad	5	6	11
58	Muzaffarnagar	2	10	12
59	Pilibhit	8	3	11
60	Pratapgarh	29	2	31
61	Prayagraj	38		38
62	Rae Bareli	30	5	35
63	Rampur	4	6	10
64	Saharanpur	7	11	18
65	Sambhal		11	11
66	Sant Kabir Nagar	9	1	10
67	Shahjahanpur	3		3
68	Shamli		4	4
69	Shravasti	13	6	19
70	Siddharthnagar	14	10	24
71	Sitapur	26	7	33
72	Sonhadra	22		22
73	Sultanpur	31	6	37

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
74	Unnao	24	5	29
75	Varanasi	11	2	13
	Grand total	1001	463	1464




Figure-2: Map showing monitoring wells during May 2024

4.0. RAINFALL

The district wise monthly grided rainfall data collected from Indian Meteorological Department; India WRIS were used to analyzed the

rainfall pattern. Table-2 gives the district wise normal rainfall and actual rainfall of Pre-monsoon 2023 and Pre-monsoon 2024 with the departure from normal rainfall.




Figure-3: Actual Rainfall of the district during the period of March–May, 2024

Table-2: District wise variability of rainfall in Uttar Pradesh (2024)

S.No	District	Normal rainfall (mm)	Actual rainfall (mm) 2022	Deviation%	Actual rainfall (mm) 2023	Deviation%	% Deviation 2023 To 2022
1	Agra	24.5	65.69	168.12	8.23	-66.41	-698.18
2	Aligarh	29.9	87.68	193.24	19.18	-35.85	-357.14
3	Ambikapur	35.5	49.37	39.07	7.92	-77.69	-523.36
4	Aunriya	23.4	34.96	49.40	8.96	-61.71	-290.18
5	Ayodhya	29.8	49.18	65.03	8.58	-71.21	-473.19
6	Azamgarh	29.6	39.85	34.63	17.04	-42.43	-133.86
7	Baghpat	32.7	108.1	230.58	12.67	-61.25	-753.20
8	Balranch	47.6	90.18	89.45	25.7	-46.01	-250.89
9	Ballia	29.9	47.03	57.29	20.5	-31.44	-129.41
10	Balrampur	60	37.98	-36.70	24.4	-59.33	-55.66
11	Banda	18.5	54.78	196.11	16.87	-8.81	-224.72
12	Barabanki	26.9	86.88	222.97	24.7	-8.18	-251.74
13	Barilly	35.9	186.4	419.22	42.18	17.49	-341.92
14	Basni	42.9	36.43	-15.08	18.66	-56.50	-95.23
15	Bijnor	45.6	161	253.07	48.79	7.00	-229.99
16	Budhan	29.2	83.23	185.03	27.88	-4.52	-198.53
17	Bulandshahr	28.8	109.11	278.85	17.06	-40.76	-539.57
18	Chandauli		78.94		12.48		-532.53
19	Chitrakoot		73.82		14.89		-395.77
20	Deoria	51.4	59.97	16.67	29.3	-43.00	-104.68
21	Etah	27.7	56.31	103.29	14.08	-49.17	-299.93
22	Etawah	24.9	41.13	65.18	17.46	-29.88	-155.57
23	Farrukhbad	26.8	57.43	114.29	12.07	-54.96	-375.81
24	Fatehpur	19.9	28.6	43.72	5.55	-72.11	-115.32
25	Firozabad	21.2	66.57	214.01	13.73	-35.24	-384.85
26	Gautam Buddha Nagar		77.37		16.09		-380.86

27	Ghazalbad	36.3	96.55	165.98	12.67	-65.10	-662.04	57	Prantighath	68.65	10.2	-573.04
28	Ghanpur	25.4	58.55	130.51	21.44	-15.59	-173.09	58	Prangpur	18.5	8.93	-568.53
29	Gonda	37.6	63.01	67.58	15.5	-58.78	-306.52	59	Rae Bareilly	18.6	12.18	-293.60
30	Gorakhpur	60.3	43.84	-27.30	12.99	-78.46	-237.49	60	Rampur	53.1	53.45	-263.95
31	Hannpur	13.6	53.35	292.28	24.73	81.84	-115.73	61	Saharanpur	41.4	34.72	-373.07
32	Hardoi	30.7	63.34	106.32	16.39	-46.61	-286.46	62	Samt Kabir Nagar	35.36	13.06	-170.75
33	Jaloun	17.1	41.01	139.82	30.47	78.19	-34.59	63	Samt Ravi Das Nagar	108.75	14.97	-40.36
34	Jaunpur	23.1	73.56	218.44	9.28	-59.83	-692.67	64	Shahjahanpur	35.7	29.75	-165.88
35	Jhansi	17	22.93	34.88	27.65	62.65	17.07	65	Shrawasti	47.6	7.33	-363.77
36	Jyotiba Phule Nagar	34.4	134.36	348.72	21.31	-38.05	-624.35	66	Siddharth Nagar	51.9	54.01	40.90
37	Kannauj		72.06		11.31		-537.14	67	Sitapur	33	18.92	-358.46
38	Kanpurdehat	19.3	55.99	190.10	12.24	-36.58	-357.43	68	Sonbhadra	46.55	26.65	-74.67
39	Kanpur Nagar	17.1	67.27	293.39	14.74	-13.80	-356.38	69	Sulampur	70.46	10.91	-60.18
40	Kaushambi		53.34		16.5		-223.27	70	Utranchi	20.4	12.05	-437.10
41	Kaushambi		54.72		10.72		-410.45	71	Varanasi	85.49	17.47	-389.35
42	Kheri	49.8	98.48	97.75	20.57	-58.69	-378.76	Average	24.7	246.11	-29.27	
43	Kushinagar	68.8	45.3	-34.16	66.15	-3.85	31.52	Average	74.97	158.07	-329.33	
44	Lalitpur	17.8	31.14	74.94	49.27	176.80	36.80					
45	Lucknow	25.3	87.98	247.75	23.31	-7.87	-277.43					
46	Mahmud Nagar	22.7	70.16	209.07	17.17	-24.36	-308.62					
47	Maharajganj	74.9	51.7	-30.97	56.5	-24.57	8.50					
48	Makoba		52.95		24.65		-114.81					
49	Meerut	23.4	60.66	159.23	18.27	-21.92	-232.02					
50	Mehaura	17.5	82.22	369.83	11.36	-35.09	-623.77					
51	Mi	33.6	31.24	-7.02	11.36	-66.19	-175.00					
52	Mirzapur	39.4	165.06	318.93	17.66	-55.18	-834.65					
53	Muzaffarpur	17.6	62.22	253.52	17.51	-0.51	-255.34					
54	Muzaffarpur	35.8	141.44	295.08	19.95	-44.27	-608.97					
55	Muzaffarpur	30.7	164.88	437.07	29.38	-4.30	-461.20					
56	Pilibhit	38.8	144.29	271.88	26.08	-32.78	-453.26					




Figure-4: District wise deviation % in Rainfall during the period of March-May, 2024

5.0. GROUNDWATER LEVEL SCENARIO (May 2024)

5.1. SHALLOW AQUIFER (UNCONFINED)

5.1.1. DEPTH TO WATER LEVEL

Depth To Water Level in Unconfined Aquifer (May2024)

The depth to water level of 971 wells is used for the analysis. It shows that water levels vary between 0.01mbgl (Aligarh district) to 44.56 mbgl (Gautam Buddh Nagar district). Water level of less than 2 mbgl is recorded in 1.59% of wells, between 2 to 5 mbgl in 31.18% of wells, between 5 to 10 mbgl in 42.50% of wells, between 10 to 20 mbgl in 20.36% of wells, between 20-30 mbgl in 3.57% of well and water level between 30-40 mbgl is registered in 0.79% of wells. Percentage of wells shown in Figure No. 6 for unconfined aquifers and Depth to Water level of unconfined aquifers is shown in Figure – 5. Shallow water level of less than 2 mbgl is seen in isolated patches in parts of Mathura, Aligarh, Kannauj, Mau, Moradabad, districts covering only an area of 1.5% of the State. Water level of 2 to 5 mbgl is majorly observed in the parts of Moradabad, Rampur, Bareilly, Pilibhit, Shahjahanpur, Lakhimpur Kheri, Sitapur, Shravasti, Bahraich, Balrampur, Gonda, Sidhdharth Nagar, Basti, Sant Kabir Nagar, Gorakhpur, Maharajganj, Kushinagar, Deoria, Mau, Ballia and isolated patches are seen in the parts of Ghazipur, Chandauli, Sonbhadra, Mirzapur, Varanasi, Prayagraj, Jaunpur, Sultanpur, Amethi, Raebareli,

Barabanki, Unnao, Kanpur Nagar, Kannauj, Jalaun, Auraiya, Etawah, Mainpuri, Hardoi, Farrukhabad, Lalitpur, Mahoba districts of Uttar Pradesh covering the area of 31% the State.




Figure-5: Depth to water level of unconfined aquifer during May 2024

The 42.5% area has depth to water level of 5 to 10 mbgl which is observed in the parts of Saharanpur, Bijnaur, Muzaffarnagar, Moradabad, Rampur, Bareilly, Shahjahanpur, Lakhimpur Kheri, Hardoi, Sitapur, Lucknow, Barabanki, Raebareli, Amethi, Ayodhya, Pratapgarrh, Sultanpur, Ambedkar Nagar, Prayagraj, Jaunpur,

Azamgarh, Gorakhpur, Mau, Ballia, Ghazipur, Chandauli, Mirzapur, Sonbhadra, Unnao, Kanpur Nagar, Kanpur Dehat, Banda, Mahoba, Hamirpur, Jhansi, Lalitpur, Jalaun, Aurraiya, Kannauj, Etawah, Mainpuri, Kasganj, Etah, Aligarh, Hathras, Mathura.




Figure-6 Percentage of wells in different water level ranges in Unconfined aquifer

20% of the area falls under Water level of 10 to 20 mbgl and are observed in parts of Saharanpur, Shamli, Baghpat, Ghaziabad, Muzaffarnagar, Bijnor, Meerut, Hapur, Amroha, Sambhal, Gautam Budh Nagar, Bulandshahar, Budaun, Shahjahanpur, Farrukhabad, Etah, Hathras, Mathura, Agra, Firozabad, Mainpuri, Etawah, Aurraiya, Kanpur Dehat and Nagar, Jalaun, Hamirpur, Mahoba, Banda, Fatehpur,

Raebareli, Lucknow, Kaushambi, Chitrakoot, Pratapgarh, Prayagraj, Mirzapur, Varanasi, Ghazipur, Sonbhadra. Deeper water levels of more than 20mbgl cover 4% area mainly in the parts of Baghpat, Shamli, Muzaffarnagar, Ghaziabad, G.B. Nagar, Sambhal, Agra, Firozabad, Etawah, Lucknow, Hamirpur, Fatehpur and Banda.

5.1.2 ANNUAL FLUCTUATION IN WATER LEVEL

Rise in Water Levels: Out of 874 wells analyzed, it is observed that, the rise in water level of less than 2m is recorded in 20.71% wells, 2 to 4 in 4% wells and more than 4 m in 11.1% of the wells. Water level rise of less than 2m is seen in parts of Bijnore, Sambhal, Moradabad, Rampur, Bareilly, Firozabad, Mainpuri, Etawah, Aurraiya, Kannauj, Farrukhabad, Kanpur Dehat and Nagar, Jhansi, Hamirpur, Mahoba, Lalitpur, Lakhimpur Kheri, Shahjahanpur, Hardoi, Lucknow, Barabanki, Gonda, Sidhdharth Nagar, Balrampur, Basti, Maharajganj, Kushinagar, Ballia, Ghazipur, Chandauli, Sonbhadra, Chitrakoot districts. Water level rise of 2 to 4 m is observed mainly in Baghpat, Meerut, Ghaziabad, Agra, Firozabad, Mainpuri, Etawah, Aurraiya, Kanpur Nagar, Lucknow, Chitrakoot, Fatehpur, Balrampur, Varanasi, Ghazipur, Chandauli, Sonbhadra etc. Rise of more than 4m is significantly observed in Baghpat, Meerut, Ghaziabad, Firozabad, Lucknow, Chitrakoot, Fatehpur, Balrampur and Varanasi districts.




Figure-7: Percentage of wells showing rise and fall in WL in unconfined aquifer (May 2023 to May 2024)

Falling Water Levels:

Out of 874 wells analyzed, 58.58% of the area shows fall in water levels and recorded water level of less than 2m fall, while 4.12% of wells are in the range of 2 to 4m and remaining 1.48% shows fall of more than 4m. Fall of less than 2m is mainly observed in Bareilly, Budaun, Aligarh, Kasganj, Hardoi, Sitapur, Pilibhit, Lakhimpur kheri, Lucknow, Unnao, Fatehpur, Banda, Hamirpur, Pratapgarrh, Prayagraj, Jaunpur, Azamgarh, Mau, Deoria, Mirzapur and Sonhadra etc.

Fall of 2 to 4 m is observed mainly in isolated patches of G.B. Nagar, Mathura, Etah, Kasganj, Agra, Jalaun, Kanpur Nagar, Jhansi, Banda, Pratapgarrh and Mirzapur etc. Fall of more than 4m is observed in isolated patches of G.B. Nagar, Mathura, Kasganj, Jalaun, Jhansi, Banda and Pratapgarrh districts. Annual water level fluctuation in

unconfined aquifer (May 2023 -May2024) is shown in Figure- 8 and Percentage of wells showing rise and fall in WL for unconfined aquifer (May2023 to May2024) in Figure – 7.




Figure-8: Annual water level fluctuation in unconfined aquifer (May 2023 -24)




Figure-9: Decadal water level fluctuation in unconfined aquifer (May 2014 -May2023 with respect to 2024)

Decadal Fluctuation of Water Level in Unconfined Aquifer (Decadal Mean May (2014-2023) to May 2024)

Rise in Water Levels:

Out of 554 analyzed wells, the rise in water level of less than 2m is recorded in 28.52%wells, 2 to 4 m in 3.24 % wells and more than 4m in 0.9% of the wells. Water level rise of less than 2m is seen in Bijnor, Aligarh, Firozabad, Kanpur Dehat, Jalaun, Hamirpur, Jhansi, Mahoba,

Banda, Basif, Siddharth Nagar, Balrampur, Shravasti, Bahraich, Gonda and Chandali regions. Water level rise of 2 to 4 m is observed mainly in isolated patches of Bijnor, Agra, Lalitpur, Hamirpur, Sonbhadra and Lucknow districts etc and rise of more than 4m is significantly observed in isolated patches of Agra, Amroha, Lalitpur, Lucknow and Varanasi districts.

Fall in Water Levels:

Out of the 554 analyzed wells, 55.77% of the area shows, fall in water levels of less than 2m while 8.3% in the range of 2 to 4m and remaining 3.24% wells registered water level fall of more than 4m. Fall of less than 2 m is observed in major parts of Eastern Uttar Pradesh such as Sonbhadra, Ghazipur, Ballia, Mau, Jaunpur, Deoria, Ayodhya and Central parts of Uttar Pradesh such as Kannauj, Hardoi, Siapur, Farrukhabad and Kasganj districts etc. Fall of 2 to 4m is observed in isolated patches of Shamli, Ghaziabad, Gautam Budh Nagar, Bulandshahar, Sambhal Budaan, Hardoi. Chitrakoot, Pratapgarh, Mirzapur and Sonbhadra districts etc. Fall more than 4m is observed in isolated patches of Shamli, Gautam Budh Nagar, Bulandshahar, Hardoi, Raebareli, Chitrakoot, Pratapgarh, Mirzapur and Sonbhadra districts etc. Decadal Water level fluctuation form (May 2014- 2023) with respect to May 2024 is shown in the Figure – 9 and percentage of wells showing

rise and fall in WL for unconfined Aquifer (Decadal fluctuation (2014-2023) w.r.t. May 2024) is shown in Figure-10.




Figure-10: Percentage of wells showing rise and fall in WL in unconfined Aquifer (Decadal Fluctuation (2014-2023) w.r.t. May 2024)


5.2 DEEPER AQUIFER (CONFINED/SEMI-CONFINED)

5.2.1 DEPTH TO WATER LEVEL

Depth To Piezometric Level in Confined/Semi-Confined Aquifer (May 2024)

Analysis of piezometric level data of 205 wells shows piezometric levels vary between 0.57 mbgl (Mau) to 39.11 mbgl (Agra district). Piezometric level of less than 2 mbgl is recorded in 0.88% of wells, between 2 to 5 mbgl (17.70%) of wells, between 5 to 10 mbgl in

31.42% of wells, between 10 to 20 mbgl in 38.50% of wells, between 20-30 mbgl in 9.29% of wells, between 30 – 45 mbgl in 2.21% and none of the wells have registered piezometric level more than 40 mbgl. Percentage of wells in different water level ranges for Confined aquifer is shown in Figure-11. Shallow piezometer level of less than 2 mbgl is noticed in isolated patches of Azamgarh and Mau districts of the State. Piezometric level of 2 to 5 mbgl mainly observed in parts of Bahraich, Barabanki, Gonda, Shravasti, Balrampur, Siddharth Nagar, Maharajganj, Kushinagar, Gorakhpur etc. districts of the State.



0 - 2 02 to 5 5 to 10 10 to 20 20 to 30 30 to 45 >45

Figure-11: Percentage of wells in different water level ranges in Confined aquifer

Eastern to North-Eastern part of the State falls under piezometric level of 5 to 10 mbgl with significant area of Ambedkar Nagar, Azamgarh, Bahraich, Sitapur, Kheri, Meerut districts etc. Piezometric level of 10 to 20 mbgl is observed mostly in North western parts of the state that is in Saharanpur, Shamli, Muzaffarnagar, Baghpat, Meerut, Ghaziabad, Bulandshahar, Bijnor, Moradabad, Sambhal, Bareilly, Budann, Pilibhit, Shahjahanpur, Unnao, Pratapgarh, Varanasi etc. districts. The peizometric level of 20-30 mbgl is observed in the isolated patches of Baghpat, Banda, Fatehpur, Hamirpur, Hathras, Moradabad, Muzaffarnagar, Sambhal, Shamli, Varanasi etc and peizometric level of 30 – 45 mbgl observed mainly in areas of Agra, Hamirpur, Mathura, Muzaffarnagar district.

52.3 ANNUAL FLUCTUATION IN PIEZOMETRIC LEVEL

Annual Fluctuation of Piezometric Level in Confined / Semi-confined Aquifer (May 2023 to May 2024)

Rise in piezometric levels:

Out of 96 wells, the rise of less than 2m piezometric level is recorded in 7.3% wells and more than 2m – 4m in 1.04% of the wells. Piezometric level rise more than 4m in 7.3% wells. Piezometric level rise of less than 2m is seen in the districts, of Chitrakoot, Fatehpur, Rampur, Sambhal, districts. Peizometric level rise from 2 – 4m is seen

significantly in Unnao district. Piezometric level rise greater than 4m is seen in parts of Banda, Chitrakoot, Fatehpur, Hamirpur, Mahoba, Rampur and Sambhal districts.

Fall in Piezometric Levels: Out of 96 analyzed wells 75% of wells shows fall in piezometric levels of less than 2m while 4.16% wells registered piezometric level for fall of more than 2m. Fall of less than 2m is mainly observed in parts of Ambedkar Nagar, Amroha, Fatehpur, Ballia, Banda, Baghpat, Chitrakoot, Gorakhpur, Mau, Meerut, Sidharth Nagar, Sitapur, Unnao etc. districts. Piezometric level fall of 2 – 4m in minor parts of Ambedkar Nagar, Bulandshahar, Banda and Moradabad districts. Piezometric level fall of >4m is seen in Banda, Fatehpur and Mahoba districts. Percentage of wells showing rise and fall in piezometer level of confined aquifer (May 2023 to May2024) is shown in the Figure – 12.




Figure 12: Percentage of wells showing rise and fall in piezometric level in unconfined (May 2023 to May 2024)

Decadal Fluctuation of Piezometric Level in Confined/Semi-confined Aquifer (Decadal Mean May (2014-2023) to May 2024)

Rise in piezometric levels:

Out of 6 wells, that have registered rise in piezometric levels, 16.67 % have recorded less than 2m and remaining 16.67% wells registered piezometric level rise of more than 4m. Fall of less than 2m is observed 50% and fall of 2 – 4m range is seen in 16.67% and none of the districts is observed in fall of greater than 4m decadal fluctuation. Out of 6 wells rise of water level fluctuation less 2m is seen in Rampur district, whereas rise of more than 4m is typically seen in Amroha district. Fall of less than 2m is observed in all districts mainly in parts of Ghaziabad, Meerut districts, which covered 50% of the area

shows fall of more than 4 m. Percentage of wells showing rise and fall in piezometric level in confined/semi- confined Aquifer (Decadal Mean May (2014-2023 to May2024) in Figure – 13.




Figure-13: Percentage of wells showing rise and fall in piezometric level in confined/semi- confined Aquifer (Decadal Mean May (2014-2023 to May 2024).




6.0. SUMMARY

As a component of the National Ground Water Monitoring Programme, the CGWB, NR, Lucknow conducts monitoring of the ground water conditions on a quarterly basis: in January, pre-monsoon May, August, and post-monsoon November. Additionally, a yearly assessment of ground water quality is performed in May. As of May 31, 2024, the Northern Region-Central Ground Water Board supervises

1001 dug wells and 463 piezometers. This comprehensive effort aims to portray the variations in the states ground water conditions across different aquifers. In May 2024, around 75% of the state's area exhibited a depth to waterlevel within 10 meters below ground level for unconfined aquifers. And around 49% for confined. Deeper water levels of more than 20m cover 4% area of the State covering mainly Agra, Gautam Buddha Nagar, Lucknow, Baghpat, Hamirpur, Jhansi districts.

The ground water level in Uttar Pradesh during May 2024 has been significantly influenced by rainfall patterns from March 2024 to May 2024. This period witnessed a departure of -28.19% from the normal, classifying most of the region in the category of deficit rainfall. This has result in fall in the water level in many districts of UP. Annual water level comparison with previous year May-2023 to May-2024 has shown fall in about 63% area of the state in unconfined aquifer and 84% in confined aquifer because of the rainfall deficit in 2024. Around 32% of the area experienced rise of water level in decadal mean water level fluctuation of May, 2013-2024, with respect to May, 2024, in unconfined aquifer whereas 49% of the area experienced rise in decadal mean water level of May 2014-2023 with respect to May, 2024 in confined aquifer. The monsoon in 2024 witnessed significant fluctuations in rainfall pattern across the state. The evident fall in

annual ground water level during May 2024 in Uttar Pradesh can be attributed to a substantial deficit in rainfall when comparing 2024 to 2023. Rainfall distribution varied with major region of Uttar Pradesh receiving deficit in rainfall while region of Lalitpur experienced comparatively excess rainfall.



GROUND WATER LEVEL BULLETIN
UTTAR PRADESH
 MAY 2024

ABSTRACT

Ground water level Scenario during May – 2024 highlighting the finding, status of ground water level in different aquifer and its seasonal, annual and decadal comparison.

CGWB, NORTHERN REGION LUCKNOW

1.0. INTRODUCTION

Groundwater bulletin is prepared by CGWB depicting changes in groundwater regime of the country through different seasons. It is an effort to obtain information on groundwater levels through representative monitoring wells. The important attributes of groundwater regime monitoring are groundwater level. The natural conditions affecting the groundwater regime involve climatic parameters like rainfall, vapor transpiration etc. whereas anthropogenic in fluences include pumpage from the aquifer, recharge due to irrigation systems and other practices like waste disposal etc.

Groundwater levels are being measured by Central Ground Water Board four times a year during January, March, April, May, August and November. The regime monitoring started in the year 1969 by Central Groundwater Board. A network of 1464 observation wells called National Hydrograph Network Stations (NHNS), as on 30.05.2024, located all over the State is being monitored.

2.0. STUDY AREA

The State of Uttar Pradesh forms a part of vast Gangetic Alluvial Plain covering an area of 2,40,928 Sq. Km. and lies between North latitude $23^{\circ}52'12''$ & $30^{\circ}24'30''$ and East longitude $77^{\circ}05'38''$ & $84^{\circ}38'30''$. It is bounded by Uttarakhnad on the NW, Nepal on the NE, Bihar on the

East, Madhya Pradesh in the South, and Haryana, Delhi & Rajasthan in the West as shown in figure 1.




Figure-1: Map showing major aquifers and administrative divisions of UP

The state is covered with rich fertile soil and underlain by a large thickness of alluvium making it one of the richest grounds water repositories of the world. Ground water is a major source of fresh water on earth. It is the most dependable source of water, comparatively free

from the vagaries of nature, easily accessible, available at the point of use and economical. Hence it is being developed indiscriminately and the ground water reservoir is stressed. The State being the most populous in the country, with a population density of 829 persons per sq. km and a high rate of population growth (20%), its demand for water is soaring. Also due to industrialization, urbanization and modern farming practices, its quality is also at stake.

demand for water is soaring. Also due to industrialization, urbanization and modern farming practices its quality is also at stake.

The food production in Uttar Pradesh is commensurate with the self-sufficiency of the country. One of the major contributors for this sufficiency is irrigation. To meet this high irrigational requirement, water resources are being increasingly developed. Ground water contributes to about 71 % of the irrigation needs of the State. The indiscriminate development of ground water has resulted in depletion of groundwater storage and lowering of water level in certain areas on one hand. On other side the surface water development in are as having shallow water level has resulted in water logging and soil salinization. The geology and structure of the formations existing in an area control by occurrence and movement of ground water. The geomorphic conditions also have a great impact on ground water scenario. The larger part of the State is under lain by fluvial sediments laid down in

the fore deep between Plateau region in south and Himalayas in north during the Quaternary period by the Indus-Ganga system of drainage over the Precambrian topography existing during geological past. These deposits owe their origin to riverine activity. The southern part of the State has entirely different geological conditions being underlain by Precambrian formations under a thin alluvial cover. Broadly, the State can be divided into two hydrogeological units.

1. Unconsolidated zone.
2. Consolidated, hardrock zone.

The hydrogeological conditions of the above two units widely differ and are discussed subsequently in brief.

Unconsolidated Zone:

This unit covers nearly 85% of the State area. The unconsolidated formations comprising the area have been deposited through mighty rivers originating from the great Himalayan Mountains. These sediments are an admixture of pebble, gravel, sand, silt, clay and kankar. The sediments are generally coarser in the north and gradually become finer southeast ward along downstream of the drainage which is a typical feature of fluvial deposits. This zone consists of mainly two parts, the Terai and the Alluvial Plain. However, the foot hill zone is very small part of Bhaber belt and lies in the northern parts of Bijnore

and Saharanpur districts. The Terai is a narrow-disconnected belt along the northwestern fringe of the State. The Alluvial Plain occupies the area south of Terai and can further be divided into two sub units - Younger Alluvium and Older Alluvium.

The younger alluvium occurs mostly along the present-day flood plain area. The continuous shifting of the drainage network with time caused reworking of their earlier deposits giving rise to the younger alluvium. The older alluvium occupying comparatively high area covers major part of the Plain. A typical characteristic of older alluvium is formation of kankar within itself due to leaching of calcium carbonate under favourable climatic conditions. The kankar occasionally forms pans restricting downward movement of water.

The thickness of alluvial sediments is variable and generally goes upto 500m. below which occur the semi-consolidated Upper Siwalik formations. The Shallower basement occurs in isolated areas which are known as "Basement highs." This unconsolidated zone is porous and permeable with primary intergranular porosity and has good ground water potential. The sub-surface correlation of formations in the state has shown presence of several aquifers down to a depth of 750 m below the ground. These aquifers mainly encountered in Central Ganga Plain have been grouped on the basis of lithological characters as well as based on interpretation of electrical logs of Boreholes drilled and are as

follows:

1. First aquifer	0.0 – 150.00 mbgl
2. Second aquifer	160.00 – 210.00 mbgl
3. Third aquifer	250.00 – 360.00 mbgl
4. Forth deep aquifer	380.00 – 600.00 mbgl

The upper part of the first aquifer down to 50 mbgl is the main source of drinking water through hand pumps and dug wells and is unconfined in nature. The first aquifer as a whole which is under unconfined to semi-confined conditions, it is the most potential aquifer group which is the main source of groundwater in the State extensively exploited through private as well as Government tube wells to meet the drinking water and irrigation needs. The deeper aquifers are confined in nature being exploited to a very limited extent. The yield of the second aquifer is limited while the third aquifer is potential. The shallow and phreatic aquifers are under heavy stress.

Consolidated Zone:

The Bundelkhand Vindhyan plateau region is underlain by a variety of Precambrian formations, mostly granite and granite gneisses, Vindhyan sandstone, limestone & shale, under a thin alluvial cover or without alluvial cover. As such these formations are hard and compact and devoid of any primary porosity. The ground water in these formations

occurs in the secondary porosity of these formations. The secondary porosity has developed due to cracks and fractures which are open at the surface and tighten at depth. The ground water occurs under unconfined or water level conditions in these formations.

The alluvial sediments of moderate depth along the river course sand in valleys form potential ground water repositories. The weathered mantle over the entire until so forms potential aquifers. These aquifers are being monitored mostly through open wells over the area.

3.0. GROUNDWATER LEVEL MONITORING

Central Ground Water Board, Northern Region, is monitoring changes in groundwater regime in Uttar Pradesh State on quarterly basis continuously. This is facilitated by a network of monitoring stations in the State located in diverse hydrogeological and geomorphic units. The number of operational wells till May 2024 was 1464 which include 1001 dug wells and 463 piezometers shown in figure 2. The district-wise breakup of the water level monitoring stations is given in Table-1.

Table-1: District-wise distribution of water level monitoring station

S.No	District	Number Of Water Level Monitoring Stations 2024		
		Dw	Pz	Total
1	Agra	6	11	17

S.No	District	Number Of Water Level Monitoring Stations 2024		
		Dw	Pz	Total
2	Aligarh	9	4	13
3	Ambekkamagar	9	16	25
4	Amethi	36	2	38
5	Amroha		9	9
6	Auraiya	9	2	11
7	Ayodhya	13	5	18
8	Azamgarh	22	10	32
9	Baghpat	2	9	11
10	Bahraich	20	15	35
11	Ballia	21	2	23
12	Balrampur	17	9	26
13	Banda	10	18	28
14	Barabanki	37	4	41
15	Barilly	11	4	15
16	Basti	14	3	17
17	Bhadohi	7		7
18	Bijnor	6	14	20
19	Budann		10	10

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
20	Bulandshahr	2	14	16
21	Chandauli	15	4	19
22	Chitrakoot	10	14	24
23	Deoria	28	1	29
24	Etah	2	4	6
25	Etawah	11	2	13
26	Farrukhabad	1	4	5
27	Fatehpur	13	14	27
28	Firozabad		6	6
29	Gautam Buddha Nagar		8	8
30	Ghaziabad		3	3
31	Ghaziipur	22	11	33
32	Gonda	25	4	29
33	Gorakhpur	15	16	31
34	Hamirpur	12	11	33
35	Hapur		4	4
36	Hardoi	16	7	23
37	Hathras	5	4	9

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
38	Jalaun	32	6	38
39	Jaunpur	30	11	41
40	Jhansi	20	2	22
41	Kannauj	11	2	13
42	Kanpur Dehat	12	1	13
43	Kanpur Nagar	16	1	17
44	Kasganj	4	11	15
45	Kaushambi	9	2	11
46	Kheri	26	5	31
47	Kushinagar	28		28
48	Lalitpur	19	4	23
49	Lucknow	9	15	24
50	Mahoba	10	6	16
51	Mahrajganj	13	1	14
52	Mainpuri	4	5	9
53	Mathura	17	3	20
54	Mau	13	3	16
55	Meerut	1	12	13

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
56	Mirzapur	25		25
57	Moradabad	5	6	11
58	Muzaffarnagar	2	10	12
59	Pilibhit	8	3	11
60	Pratapgarh	29	2	31
61	Prayagraj	38		38
62	Rae Bareli	30	5	35
63	Rampur	4	6	10
64	Saharampur	7	11	18
65	Sambhal		11	11
66	Sant Kabir Nagar	9	1	10
67	Shahjahanpur	3		3
68	Shamli		4	4
69	Shrawasiti	13	6	19
70	Siddharthnagar	14	10	24
71	Sitapur	26	7	33
72	Sonbhadra	22		22
73	Sultanpur	31	6	37

S.No	District	Number Of Water Level Monitoring Stations		
		2024		
		Dw	Pz	Total
74	Unnao	24	5	29
75	Varanasi	11	2	13
	Grand total	1001	463	1464




Figure-2: Map showing monitoring wells during May 2024

4.0. RAINFALL

The district wise monthly grided rainfall data collected from Indian Meteorological Department; India WRIS were used to analyzed the

rainfall pattern. Table-2 gives the district wise normal rainfall and actual rainfall of Pre-monsoon 2023 and Pre-monsoon 2024 with the departure from normal rainfall.




Figure-3: Actual Rainfall of the district during the period of March-May, 2024

Table-2: District wise variability of rainfall in Uttar Pradesh (2024)

S.No	District	Normal rainfall (mm)	Actual rainfall 2023	Deviation%	Actual rainfall (mm) 2024	Deviation%	% Deviation 2024 To 2023
1	Agra	24.5	65.69	168.12	8.23	-66.41	-698.18
2	Aligarh	29.9	87.68	193.24	19.18	-35.85	-357.14
3	Ambekankaragar	35.5	49.37	39.07	7.92	-77.69	-523.36
4	Auraya	23.4	34.96	49.40	8.96	-61.71	-250.18
5	Ayodhya	29.8	49.18	65.03	8.58	-71.21	-473.19
6	Azamgarh	29.6	39.85	34.63	17.04	-42.43	-133.86
7	Baghpat	32.7	108.1	230.58	12.67	-61.25	-753.20
8	Bahraich	47.6	90.18	89.45	25.7	-6.01	-250.89
9	Ballia	29.9	47.03	57.29	20.5	-31.44	-129.41
10	Baranpur	60	37.98	-36.70	24.4	-59.33	-55.66
11	Banda	18.5	54.78	196.11	16.87	-8.81	-224.72
12	Barabanki	26.9	86.88	222.97	24.7	-8.18	-251.74
13	Bareilly	35.9	186.4	419.22	42.18	17.49	-341.92
14	Basti	42.9	36.43	-15.08	18.66	-56.50	-95.23
15	Bijnor	45.6	161	253.07	48.79	7.00	-229.99
16	Budhan	29.2	83.23	185.03	27.88	-4.52	-198.53
17	Bulanahar	28.8	109.11	278.85	17.06	-40.76	-539.57
18	Chandauli		78.94		12.48	-532.53	
19	Chitkoot		73.82		14.89	-395.77	
20	Deoria	51.4	59.97	16.67	29.3	-43.00	-104.68
21	Etah	27.7	56.31	103.29	14.08	-49.17	-299.93
22	Etaawah	24.9	41.13	65.18	17.46	-29.88	-135.57
23	Farrukhabad	26.8	57.43	114.29	12.07	-54.96	-375.81
24	Fatehpur	19.9	28.6	43.72	5.55	-72.11	-415.32
25	Firozabad	21.2	66.57	214.01	13.73	-53.24	-354.85
26	Gautam Buddha Nagar		77.37		16.09	-380.86	

27	Ghazipur	36.3	96.55	165.98	12.67	-65.10	-662.04
28	Ghazipur	25.4	58.55	130.51	21.44	-15.59	-173.09
29	Gonda	37.6	63.01	67.58	15.5	-58.78	-306.52
30	Gorakhpur	60.3	43.84	-27.30	12.99	-78.46	-237.49
31	Hampiur	13.6	53.35	292.28	24.73	81.84	-115.73
32	Hardoi	30.7	63.34	106.32	16.39	-66.61	-286.46
33	Jajau	17.1	41.01	139.82	30.47	78.19	-34.59
34	Jaunpur	23.1	73.56	218.44	9.28	-59.83	-692.67
35	Jhansi	17	22.93	34.88	27.65	62.63	17.07
36	Jyotiba Pratinagar	34.4	154.36	348.72	21.31	-38.05	-624.35
37	Kannauj		72.06		11.31		-537.14
38	Kanpurdehat	19.3	55.99	190.10	12.24	-36.58	-357.43
39	Kanpur Nagar	17.1	67.27	293.39	14.74	-13.80	-356.38
40	Kanshanpur		53.34		16.5		-233.27
41	Kaushambi		54.72		10.72		-110.45
42	Kheri	49.8	98.48	97.75	20.57	-58.69	-378.76
43	Kushinagar	68.8	45.3	-34.16	66.15	-3.85	31.52
44	Lalitpur	17.8	31.14	74.94	49.27	176.80	36.80
45	Lucknow	25.3	87.98	247.75	23.31	-7.87	-277.43
46	Maharaja Nagar	22.7	70.16	209.07	17.17	-24.36	-308.62
47	Maharajganj	74.9	51.7	-30.97	56.5	-24.57	8.50
48	Malerba		52.95		24.65		-114.81
49	Meerut	23.4	60.66	159.23	18.27	-21.92	-232.02
50	Methua	17.5	82.22	369.83	11.36	-33.09	-623.77
51	Mirzapur	33.6	31.24	-7.02	11.36	-66.19	-175.00
52	Muzaffarpur	39.4	165.06	318.93	17.66	-55.18	-834.65
53	Muzaffarpur	17.6	62.22	253.52	17.51	-0.51	-255.34
54	Muzaffarpur	35.8	141.44	295.08	19.95	-44.27	-608.97
55	Muzaffarpur	30.7	164.88	437.07	29.38	-4.30	-161.20
56	Prabhat	38.8	144.29	271.88	26.08	-32.78	-453.26

57	Pratapnagar		68.65		10.2		-573.04
58	Prayagraj	18.5	59.7	222.70	8.93	-51.73	-568.53
59	Rae Bareilly	18.6	47.94	157.74	12.18	-34.52	-293.60
60	Rampur	53.1	190.89	259.49	52.45	-1.22	-263.95
61	Saharanpur	41.4	164.25	296.74	34.72	-16.14	-373.07
62	Sant Kabir Nagar		35.36		13.06		-170.75
63	Sant Ravi Das Nagar	25.1	108.75	333.27	14.97	-10.36	-626.45
64	Shahjahanpur	35.7	79.1	121.57	29.75	-16.67	-165.88
65	Shrawasti	47.6	51.09	7.33	11.04	-76.81	-362.77
66	Siddharth Nagar	51.9	31.92	-38.50	54.01	4.07	40.90
67	Sitapur	33	86.74	162.85	18.92	-12.67	-358.46
68	Sonbhadra	32	46.55	45.47	26.65	-16.72	-74.67
69	Sultanpur	27.4	70.46	157.15	10.91	-60.18	-545.83
70	Unnao	20.4	64.72	217.25	12.05	-10.93	-437.10
71	Varanasi	24.7	85.49	246.11	17.47	-29.27	-389.35
	Average		32.26	74.97	158.07	20.84	-329.33




Figure-4: District wise deviation % in Rainfall during the period of March-May, 2024

5.0. GROUNDWATER LEVEL SCENARIO (May 2024)

5.1. SHALLOW AQUIFER (UNCONFINED)

5.1.1. DEPTH TO WATER LEVEL

Depth To Water Level in Unconfined Aquifer (May2024)

The depth to water level of 971 wells is used for the analysis. It shows that water levels vary between 0.01mbgl (Aligarh district) to 44.56 mbgl (Gautam Buddha Nagar district). Water level of less than 2 mbgl is recorded in 1.59% of wells, between 2 to 5 mbgl in 31.18% of wells, between 5 to 10 mbgl in 42.50% of wells, between 10 to 20 mbgl in 20.36% of wells, between 20-30 mbgl in 3.57% of well and water level between 30-40 mbgl is registered in 0.79% of wells. Percentage of wells shown in Figure No. 6 for unconfined aquifers and Depth to Water level of unconfined aquifers is shown in Figure – 5. Shallow water level of less than 2 mbgl is seen in isolated patches in parts of Mathura, Aligarh, Kannauj, Mau, Moradabad, districts covering only an area of 1.5% of the State. Water level of 2 to 5 mbgl is majorly observed in the parts of Moradabad, Rampur, Bareilly, Pilibhit, Shahjahanpur, Lakhimpur Kheri, Sitapur, Shravasti, Bahraich, Balrampur, Gonda, Sidhharth Nagar, Basti, Sant Kabir Nagar, Gorakhpur, Maharajganj, Kushinagar, Deoria, Mau, Ballia and isolated patches are seen in the parts of Ghazipur, Chandauli, Sonbhadra, Mirzapur, Varanasi, Prayagraj, Jaunpur, Sultanpur, Amethi, Raebareli,

Barabanki, Unnao, Kanpur Nagar, Kannauj, Jalaun, Auraiya, Etawah, Mainpuri, Hardoi, Farrukhabad, Lalitpur, Mahoba districts of Uttar Pradesh covering the area of 31% the State.




Figure-5: Depth to water level of unconfined aquifer during May2024

The 42.5% area has depth to water level of 5 to 10 mbgl which is observed in the parts of Saharanpur, Bijnaur, Muzaffarnagar, Moradabad, Rampur, Bareilly, Shahjahanpur, Lakhimpur Kheri, Hardoi, Sitapur, Lucknow, Barabanki, Raebareli, Amethi, Ayodhya, Pratapgarh, Sultanpur, Ambedkar Nagar, Prayagraj, Jaunpur,

Azamgarh, Gorakhpur, Mau, Ballia, Ghazipur, Chandauli, Mirzapur, Sonbhadra, Unnao, Kanpur Nagar, Kanpur Dehat, Banda, Mahoba, Hamirpur, Jhansi, Lalitpur, Jalaun, Aurraiya, Kannauj, Etawah, Mainpuri, Kasganj, Etah, Aligarh, Hathras, Mathura.




Figure-6 Percentage of wells in different water level ranges in Unconfined aquifer

20% of the area falls under Water level of 10 to 20 mbgl and are observed in parts of Saharanpur, Shamli, Baghpat, Ghaziabad, Muzaffarnagar, Bijnor, Meerut, Hapur, Amroha, Sambhal, Gautam Budh Nagar, Bulandshahar, Budaun, Shahjahanpur, Farrukhabad, Etah, Hathras, Mathura, Agra, Firozabad, Mainpuri, Etawah, Aurraiya, Kanpur Dehat and Nagar, Jalaun, Hamirpur, Mahoba, Banda, Fatehpur,

Raebareli, Lucknow, Kaushambhi, Chitrakoot, Pratapgarh, Prayagraj, Mirzapur, Varanasi, Ghazipur, Sonbhadra. Deeper water levels of more than 20mbgl cover 4% area mainly in the parts of Baghpat, Shamli, Muzaffarnagar, Ghaziabad, G.B. Nagar, Sambhal, Agra, Firozabad, Etawah, Lucknow, Hamirpur, Fatehpur and Banda.

5.1.2 ANNUAL FLUCTUATION IN WATER LEVEL

Rise in Water Levels: Out of 874 wells analyzed, it is observed that, the rise in water level of less than 2m is recorded in 20.71% wells; 2 to 4 in 4% wells and more than 4 m in 11.1% of the wells. Water level rise of less than 2m is seen in parts of Bijnore, Sambhal, Moradabad, Rampur, Bareilly Firozabad, Mainpuri, Etawah, Aurraiya, Kannauj, Farrukhabad, Kanpur Dehat and Nagar, Jhansi, Hamirpur, Mahoba, Lalitpur, Lakhimpur Kheri, Shahjahanpur, Hardoi, Lucknow, Barabanki, Gonda, Sidhharth Nagar, Balrampur, Basti, Maharajganj, Kushinagar, Ballia, Ghazipur, Chandauli, Sonbhadra, Chitrakoot districts. Water level rise of 2 to 4 m is observed mainly in Baghpat, Meerut, Ghaziabad, Agra, Firozabad, Mainpuri, Etawah, Aurraiya, Kanpur Nagar, Lucknow, Chitrakoot, Fatehpur, Balrampur, Varanasi, Ghazipur, Chandauli, Sonbhadra etc. Rise of more than 4m is significantly observed in Baghpat, Meerut, Ghaziabad, Firozabad, Lucknow, Chitrakoot, Fatehpur, Balrampur and Varanasi districts.




Figure-7: Percentage of wells showing rise and fall in WL in unconfined aquifer (May 2023 to May 2024)
Falling Water Levels:

Out of 874 wells analyzed, 58.58% of the area shows fall in water levels and recorded water level of less than 2m fall, while 4.12% of wells are in the range of 2 to 4m and remaining 1.48% shows fall of more than 4m. Fall of less than 2m is mainly observed in Bareilly, Buddann, Aligarh, Kasganj, Hardoi, Sitapur, Pilibhit, Lakhimpur kheri, Lucknow, Unnao, Fatehpur, Banda, Hamirpur, Pratapgarh, Prayagraj, Jaunpur, Azamgarh, Mau, Deoria, Mirzapur and Sonhadra etc.

Fall of 2 to 4 m is observed mainly in isolated patches of G.B. Nagar, Mathura, Etah, Kasganj, Agra, Jalaun, Kanpur Nagar, Jhansi, Banda, Pratapgarh and Mirzapur etc. Fall of more than 4m is observed in isolated patches of G.B. Nagar, Mathura, Kasganj, Jalaun, Jhansi, Banda and Pratapgarh districts. Annual water level fluctuation in

unconfined aquifer (May 2023 -May2024) is shown in Figure- 8 and Percentage of wells showing rise and fall in WL for unconfined aquifer (May2023 to May2024) in Figure – 7.




Figure-8: Annual water level fluctuation in unconfined aquifer (May 2023 -24)




Figure-9: Decadal water level fluctuation in unconfined aquifer (May 2014 -May 2023 with respect to 2024)

Decadal Fluctuation of Water Level in Unconfined Aquifer

Decadal Mean May (2014-2023) to May 2024)

Rise in Water Levels:

Out of 554 analyzed wells, the rise in water level of less than 2m is recorded in 28.52%/wells. 2 to 4 m in 3.24 % wells and more than 4m in 0.9% of the wells. Water level rise of less than 2m is seen in Bijnor, Aligarh, Firozabad, Kanpur Dehat, Jalaun, Hamirpur, Jhansi, Mahoba,

Banda, Basi, Siddharth Nagar, Balrampur, Shravasi, Bahraich, Gonda and Chaundali regions. Water level rise of 2 to 4 m is observed mainly in isolated patches of Bijnor, Agra, Lalitpur, Hamirpur, Sonbhadra and Lucknow districts etc and rise of more than 4m is significantly observed in isolated patches of Agra, Amroha, Lalitpur, Lucknow and Varanasi districts.

Fall in Water Levels:

Out of the 554 analyzed wells, 55.77% of the area shows, fall in water levels of less than 2m while 8.3% in the range of 2 to 4m and remaining 3.24% wells registered water level fall of more than 4m. Fall of less than 2 m is observed in major parts of Eastern Uttar Pradesh such as Sonbhadra, Ghazipur, Ballia, Mau, Jaunpur, Deoria, Ayodhya and Central parts of Uttar Pradesh such as Kannauj, Hardoi, Sitapur, Farrukhabad and Kasganj districts etc. Fall of 2 to 4m is observed in isolated patches of Shamli, Ghaziabad, Gautam Budh Nagar, Bulandshahar, Sambhal Budaan, Hardoi, Chitrakoot, Pratapgarh, Mirzapur and Sonbhadra districts etc. Fall more than 4m is observed in isolated patches of Shamli, Gautam Budh Nagar, Bulandshahar, Hardoi, Raebareli, Chitrakoot, Pratapgarh, Mirzapur and Sonbhadra districts etc. Decadal Water level fluctuation form (May 2014- 2023) with respect to May 2024 is shown in the Figure – 9 and percentage of wells showing

rise and fall in WL for unconfined Aquifer (Decadal Fluctuation (2014-2023) w.r.t. May 2024) is shown in Figure-10.




Figure-10: Percentage of wells showing rise and fall in WL in unconfined Aquifer (Decadal Fluctuation (2014-2023) w.r.t. May 2024)

5.2 DEEPER AQUIFER (CONFINED/SEMI-CONFINED)

5.2.1 DEPTH TO WATER LEVEL

Depth To Piezometric Level in Confined/Semi-Confined Aquifer (May 2024)

Analysis of piezometric level data of 205 wells shows piezometric levels vary between 0.57 mbgl (Manu) to 39.11 mbgl (Agra district). Piezometric level of less than 2 mbgl is recorded in 0.88% of wells, between 2 to 5 mbgl (17.70%) of wells, between 5 to 10 mbgl in

31.42% of wells, between 10 to 20 mbgl in 38.50% of wells, between 20-30 mbgl in 9.29% of wells, between 30 – 45 mbgl in 2.21% and none of the wells have registered piezometric level more than 40 mbgl. Percentage of wells in different water level ranges for Confined aquifer is shown in Figure-11. Shallow piezometer level of less than 2 mbgl is noticed in isolated patches of Azamgarh and Mau districts of the State. Piezometric level of 2 to 5 mbgl mainly observed in parts of Bahrach, Barabanki, Gonda, Shravasti, Balrampur, Sidhdharh Nagar, Maharajanji, Kushinagar, Gorakhpur etc. districts of the State.




Figure-11: Percentage of wells in different water level ranges in Confined aquifer

Eastern to North-Eastern part of the State falls under piezometric level of 5 to 10 mbgl with significant area of Ambedkar Nagar, Azamgarh, Bahraich, Sitapur, Kheri, Meerut districts etc. Piezometric level of 10 to 20 mbgl is observed mostly in North western parts of the state that is in Saharanpur, Sharnli, Muzaffarnagar, Baghpat, Meerut, Ghaziabad, Bulandshahar, Bijnor, Moradabad, Sambhal, Bareilly, Budaur, Pilibhit, Shahjahanpur, Unnao, Pratapgarh, Varanasi etc. districts. The peizometric level of 20-30 mbgl is observed in the isolated patches of Baghpat, Banda, Fatehpur, Hamirpur, Hathras, Moradabad, Muzaffarnagar, Sambhal, Sharnli, Varanasi etc and peizometric level of 30 – 45 mbgl observed mainly in areas of Agra, Hamirpur, Mahura, Muzaffarnagar district.

5.2.3 ANNUAL FLUCTUATION IN PIEZOMETRIC LEVEL

Annual Fluctuation of Piezometric Level in Confined / Semi-confined Aquifer (May 2023 to May 2024)

Rise in piezometric levels:

Out of 96 wells, the rise of less than 2m piezometric level is recorded in 7.3% wells and more than 2m – 4m in 1.04% of the wells. Piezometric level rise more than 4m in 7.3% wells. Piezometric level rise of less than 2m is seen in the districts, of Chitrakoot, Fatehpur, Rampur, Sambhal, districts. Peizometric level rise from 2 – 4m is seen

significantly in Unnao district. Piezometric level rise greater than 4m is seen in parts of Banda, Chitrakoot, Fatehpur, Hamirpur, Mahoba, Rampur and Sambhal districts.

Fall in Piezometric Levels: Out of 96 analyzed wells 75% of wells shows fall in piezometric levels of less than 2m while 4.16% wells registered piezometric level for fall of more than 2m. Fall of less than 2m is mainly observed in parts of Ambedkar Nagar, Amroha, Fatehpur, Ballia, Banda, Baghpat, Chitrakoot, Gorakhpur, Mau, Meerut, Sidharth Nagar, Sitapur, Unnao etc. districts. Piezometric level fall of 2 – 4m in minor parts of Ambedkar Nagar, Bulandshahar, Banda and Moradabad districts. Piezometric level fall of >4m is seen in Banda, Fatehpur and Mahoba districts. Percentage of wells showing rise and fall in piezometer level of confined aquifer (May 2023 to May 2024) is shown in the Figure – 12.




Figure12: Percentage of wells showing rise and fall in piezometric level in unconfined (May 2023 to May 2024)

Decadal Fluctuation of Piezometric Level in Confined/Semi-confined Aquifer (Decadal Mean May (2014-2023) to May 2024)

Rise in piezometric levels:

Out of 6 wells, that have registered rise in piezometric levels, 16.67 % have recorded less than 2m and remaining 16.67% wells registered piezometric level rise of more than 4m. Fall of less than 2m is observed 50% and fall of 2 – 4m range is seen in 16.67% and none of the districts is observed in fall of greater than 4m decadal fluctuation. Out of 6 wells rise of water level fluctuation less 2m is seen in Rampur district, whereas rise of more than 4m is typically seen in Amroha district. Fall of less than 2m is observed in all districts mainly in parts of Ghaziabad, Meerut districts, which covered 50% of the area

shows fall of more than 4 m. Percentage of wells showing rise and fall in piezometric level in confined/semi- confined Aquifer (Decadal Mean May (2014-2023 to May2024) in Figure – 13.




Figure-13: Percentage of wells showing rise and fall in piezometric level in confined/semi- confined Aquifer (Decadal Mean May (2014-2023 to May 2024).

6.0. SUMMARY

As a component of the National Ground Water Monitoring Programme, the CGWB, NR, Lucknow conducts monitoring of the ground water conditions on a quarterly basis: in January, pre-monsoon May, August, and post-monsoon November. Additionally, a yearly assessment of ground water quality is performed in May. As of May 31, 2024, the Northern Region-Central Ground Water Board supervises

1001 dug wells and 463 piezometers. This comprehensive effort aims to portray the variations in the states ground water conditions across different aquifers. In May 2024, around 75% of the state's area exhibited a depth to waterlevel within 10 meters below ground level for unconfined aquifers. And around 49% for confined. Deeper water levels of more than 20m cover 4% area of the State covering mainly Agra, Gautam Buddha Nagar, Lucknow, Baghpat, Hamirpur, Jhansi districts.

The ground water level in Uttar Pradesh during May 2024 has been significantly influenced by rainfall patterns from March 2024 to May 2024. This period witnessed a departure of -28.19% from the normal, classifying most of the region in the category of deficit rainfall. This has result in fall in the water level in many districts of UP. Annual water level comparison with previous year May-2023 to May-2024 has shown fall in about 63% area of the state in unconfined aquifer and 84% in confined aquifer because of the rainfall deficit in 2024. Around 32% of the area experienced rise of water level in decadal mean water level fluctuation of May, 2013-2024, with respect to May, 2024, in unconfined aquifer whereas 49% of the area experienced rise in decadal mean water level of May 2014-2023 with respect to May, 2024 in confined aquifer. The monsoon in 2024 witnessed significant fluctuations in rainfall pattern across the state. The evident fall in

annual ground water level during May 2024 in Uttar Pradesh can be attributed to a substantial deficit in rainfall when comparing 2024 to 2023. Rainfall distribution varied with major region of Uttar Pradesh receiving deficit in rainfall while region of Lalitpur experienced comparatively excess rainfall.